

# Highly Functionalized Terpyridines as Competitive Inhibitors of AKAP–PKA Interactions\*\*

Gesa Schäfer, Jelena Milić, Adeeb Eldahshan, Frank Götz, Kerstin Zühlke, Christian Schillinger, Annika Kreuchwig, Jonathan M. Elkins, Kamal R. Abdul Azeez, Andreas Oder, Marie C. Moutty, Nanako Masada, Monika Beerbaum, Brigitte Schlegel, Sylvia Niquet, Peter Schmieder, Gerd Krause, Jens Peter von Kries, Dermot M. F. Cooper, Stefan Knapp, Jörg Rademann, Walter Rosenthal, and Enno Klussmann\*

Protein kinase A (PKA) is a ubiquitous kinase that phosphorylates a broad variety of substrates. A-kinase anchoring proteins (AKAPs) confer specificity to PKA signaling by tethering the kinase to distinct cellular compartments, thereby limiting the access of PKA to a defined pool of its substrates.<sup>[1]</sup> Interactions between AKAPs and PKA play key roles in a plethora of physiologically relevant processes such as arginine-vasopressin (AVP) mediated water reabsorption in renal principal cells; AVP triggers PKA phosphorylation of the water channel aquaporin-2 (AQP2). AQP2 then redistributes from intracellular vesicles into the plasma membrane, thereby facilitating water reabsorption from primary urine. The redistribution only occurs if PKA interacts with AKAPs.<sup>[2]</sup> Dysregulation of cellular processes that depend on AKAP–PKA interactions causes or is associated with diseases.<sup>[1b,3]</sup> For example, in heart failure, elevated AVP levels contribute to the excessive water retention by the above-described mechanism. Cardiac myocyte contractility is decreased in the failing heart; the control of contractility crucially depends on AKAP–PKA interactions.<sup>[4]</sup>

PKA holoenzyme is a tetramer consisting of a dimer of regulatory (RI $\alpha$ , RI $\beta$ , RII $\alpha$ , or RII $\beta$ ) and two catalytic (C) subunits each bound to an R protomer. Upon binding of cAMP to the R subunits, the C subunits dissociate and

phosphorylate their substrates. Interactions of PKA with AKAPs are mediated by the dimerization/docking (D/D) domains of R subunit dimers and the RII binding domain (RBD) of AKAPs. Dimerized D/D domains form a hydrophobic pocket that directly interacts with the 14–25 amino acid long  $\alpha$ -helical RBD.<sup>[5]</sup> Synthetic peptides derived from RBDs of different AKAPs bind R subunits with nanomolar affinity, for example, AKAP18 $\delta$ -L314E from AKAP18 $\delta$  ( $K_D = 4$  nM; see Figures S1 and S2 as well as Table S2 in the Supporting Information).<sup>[6]</sup> Such peptides effectively inhibit AKAP–PKA interactions.<sup>[1]</sup> For example, membrane-permeable versions of AKAP18 $\delta$ -L314E<sup>[7]</sup> and Ht31<sup>[2a]</sup> (from AKAP-Lbc) abolish the AVP-induced redistribution of AQP2 in cultured principal cells. Thus disruption of AKAP–PKA interactions even appears beneficial for the treatment of diseases, such as heart failure, that are associated with AVP-dependent excessive water retention.<sup>[3]</sup>

As a consequence of their generally low membrane permeability and stability, peptides have limitations with regard to their use in cell and animal studies and for drug development. Small molecules and nonpeptide helix mimetics are considered as alternatives to peptides, and thus the small molecule FMP-API-1 was developed that inhibits AKAP–PKA interactions. However, FMP-API-1 also activates PKA,

[\*] Dr. G. Schäfer,<sup>[†]</sup> Dr. J. Milić,<sup>[†]</sup> Dr. A. Eldahshan, Dr. F. Götz, Dr. K. Zühlke, A. Oder, Dr. M. C. Moutty, S. Niquet, Dr. J. P. von Kries, Prof. Dr. W. Rosenthal, Priv.-Doz. Dr. E. Klussmann  
Max Delbrück Center for Molecular Medicine (MDC)  
Robert-Rössle-Strasse 10, 13125 Berlin (Germany)  
E-mail: enno.klussmann@mdc-berlin.de

C. Schillinger, A. Kreuchwig, M. Beerbaum, B. Schlegel, Dr. P. Schmieder, Dr. G. Krause, Prof. J. Rademann  
Leibniz-Institut für Molekulare Pharmakologie  
Robert-Rössle-Strasse 10, Berlin (Germany)

Dr. J. M. Elkins, K. R. Abdul Azeez, Prof. Dr. S. Knapp  
Nuffield Department of Clinical Medicine, Structural Genomics Consortium and Target Discovery Institute University of Oxford  
Old Road Campus Research Building  
Roosevelt Drive, Oxford, OX3 7DQ (UK)

N. Masada, Prof. D. M. F. Cooper  
Department of Pharmacology  
Tennis Court Rd., Cambridge (UK)

Prof. J. Rademann  
Free University Berlin, Institut für Pharmazie  
Königin-Luise-Strasse 2 + 4, 14195 Berlin (Germany)

[†] These authors contributed equally to this work.


[\*\*] This work was supported by the Deutsche Forschungsgemeinschaft (DFG KL1415/4-2, RA895/5-2). J.E., K.A.A., and S.K. were supported by the Oxford Structural Genomics Consortium (SGC), a charity (number 1097737) receiving funds from AbbVie, Boehringer Ingelheim, the Canada Foundation for Innovation, the Canadian Institutes for Health Research, Genome Canada, GlaxoSmithKline, Janssen, Lilly Canada, the Novartis Research Foundation, the Ontario Ministry of Economic Development and Innovation, Pfizer, Takeda, and the Wellcome Trust [092809/Z/10/Z]. AKAP = A-kinase anchoring protein, PKA = protein kinase A.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201304686>.

© 2013 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

which suggests that it causes side effects in complex systems such as whole animals.<sup>[8]</sup>


Nonpeptide helix mimetics including terphenyl and terpyridine scaffolds<sup>[9]</sup> mimic  $\alpha$ -helices by presenting amino acid derived side chains from a rodlike axis and are expected to combine the desired properties of peptides, that is, specificity/target selectivity, and those of small molecules, that is, stability and membrane permeability. We developed poly-pyridines as helix mimetics of peptide AKAP18 $\delta$ -L314E,<sup>[10]</sup> taking into account the favorable solubility, conformational orientation, and biological activity of helix mimetics of this class. The starting point was an interaction model in which hydrophobic residues (L301, L304, L308) of the peptide interact with the hydrophobic bottom of the D/D domain dimer of RII $\alpha$  and where the hydrophilic side chains such as of E300 of the peptide interact with hydrophilic residues such as Q4 at the peripheral rim of the cavity formed by the D/D dimer.<sup>[6]</sup> Several terpyridines, **1a–f** (Schemes 1 and 4), were



**Scheme 1.** Planned synthesis of terpyridines **1a–f** by a sequence of Suzuki coupling reactions.

designed in silico as synthetic targets. The hydrophobic phase of the AKAP18 $\delta$ -L314E helix in molecule **1b** corresponds to the *meta*-methyl group of the second pyridine ring, the condensed cyclopentyl ring of the third pyridine moiety, and the following two benzyl rings. Thus, they interact with the hydrophobic core (yellow in Figure 4 and Figure S1 in the Supporting Information) of the binding cavity. Other substituents address hydrophilic residues of the rim at the D/D domain such as the carboxy group of the external pyridine ring that mimics the side chain of E300 and is likely to interact with Q4 of the D/D domain.

The synthesis of the potential ligands **1a–f** involved two successive Suzuki cross-coupling reactions of 2,5-dibromopyridines **3a,b** (Scheme 2). The key to this strategy was the




**Scheme 2.** Synthesis of bipyrindyl bromides **6a,b**, the building blocks for the final coupling reaction. TMP = 2,2,6,6-tetramethylpiperidine.

distinctive reactivity of the 2-bromo and 5-bromo substituents, which allowed for the regioselective Suzuki coupling of the dibromides **3a,b**.<sup>[11]</sup>

The *ortho*-borylation of cyanopyridines **5** was adapted for the synthesis of the first pyridyl building block—the diisopropylboronic acid ester **2**. An *ortho*-directed metalation reaction (DoM) involves the cyano function, which reacts with the first equivalent of LiTMP as base and most likely forms a lithium amidinate that serves as the donor-metalation group (DMG) and directs deprotonation to the C3-site of the pyridine ring by the second equivalent of base.<sup>[12]</sup> A third equivalent of the base was required for effective metalation, presumably to transform the aliphatic ester into the enolate and thereby also protecting the neighboring benzylic position from deprotonation. The addition of four equivalents of the base increased the yield without any side reaction. Since the isolation of the diisopropylboronic ester was not possible and isolation of the boronic acid was associated with protodeboronation, we esterified the acid in situ by adding pinacol to the reaction mixture.<sup>[13]</sup> Synthesis of ester **2** required adding pinacol and heating the reaction mixture to 40 °C (Scheme 2).

The Pd-catalyzed coupling of 2,5-dibromo-3-methylpyridines **3a,b** and crude boronic acid ester **2** was carried out in the presence of potassium phosphate and yielded bipyrindines **6a,b** in low amounts. When boronic acid ester **2** was added slowly through a dropping funnel over 90 min, the yield increased to 74% (Scheme 2).

The cyclopentapyridine moiety of building blocks **4a,b** was synthesized by a Rh<sup>II</sup>-catalyzed [2+2+2] cycloaddition of 1,6-heptadiyne (**7**) with succinonitrile (**8**; Scheme 3).<sup>[14]</sup> Reducing the nitrile moiety with borane-dimethyl sulfide led to amine **10**. Monobenzylated amine **10-Bn** was obtained by reductive amination followed by dibenylation with Et<sub>3</sub>N as the base. The mixed lithium base aggregates approach was used for the regioselective halogenation at the C1-position of cyclopentapyridine **11a**.<sup>[15]</sup> No conversion of the starting material occurred when *n*-hexane was used as the solvent for the lithiation;<sup>[15]</sup> only the addition of diethyl ether as a co-solvent during the metalation process led to the formation of **4a,b**. CBr<sub>4</sub> and C<sub>2</sub>Cl<sub>6</sub> were used as electrophiles. Since the




**Scheme 3.** Synthesis of cyclopentapyridyl chlorides **4a,b**.

chlorination was more efficient, the final cross-coupling reaction was optimized using pyridyl chlorides **4a,b**.

The synthesis of cyclopentapyridine **11b** requires the separate introduction of the two different benzyl moieties. The secondary amine was obtained under harsh base conditions using  $\text{NaOH}$ , whereby the substituted benzyl group was introduced in a microwave-assisted aqueous  $\text{N}$ -alkylation.<sup>[16]</sup> Surprisingly, the employment of four equivalents of  $\text{Li}$  base was not sufficient for the lithiation of pyridine **4a** (59% yield); increasing amounts of the base raised the yield of chloride **4b** fourfold to 78%.

We developed a microwave-assisted “one-pot” procedure for the Suzuki–Miyaura coupling reaction for synthesis of terpyridines **12a,b**. For this, the borylation and  $\text{C–C}$  coupling reactions were optimized in terms of solvent,  $\text{Pd}^0$  catalyst, phosphine ligand, and reaction temperature. The best yields (**12a** = 46%, **12b** = 50%) were obtained with  $[\text{Pd}_2(\text{dba})_3]$  as the  $\text{Pd}^0$  source in combination with  $\text{XPhos}$  as a ligand and using dioxane as solvent. The use of aqueous potassium phosphate as a base was more effective than potassium carbonate. The borylation step resulted in complete consumption of the starting material when  $\text{KOAc}$  was used as the base, whereas the cross-coupling demanded the addition of a stronger base.<sup>[15]</sup> The last step of the synthesis was a hydrolysis of 2-cyanoterpyridines **12a,b**. Unexpectedly, the base-catalyzed hydrolysis (Scheme 4, conditions a) led to only partial hydrolysis of the nitrile function. The amidoterpyridylpropionic acids **1a** and **1c** were obtained from the ethyl



**Scheme 4.** Synthesis of the terpyridines **12a**, **12b**, and **12c** by microwave-assisted, “one-pot” Suzuki–Miyaura coupling followed by base- or acid-catalyzed hydrolysis to the terpyridylpropionic acids **1a–f**. Reagents and conditions: a)  $\text{NaOH}$ ,  $\text{EtOH}/\text{H}_2\text{O}$ , 85°C, 4 h; b) 6 M  $\text{HCl}$ ,  $\text{EtOH}$ , 95°C, 1 h; c)  $\text{NaOH}$ ,  $\text{EtOH}/\text{H}_2\text{O}$ , 85°C, 1 h.  $\text{dba} = \text{trans-}, \text{trans-}$ -dibenzylideneacetone,  $\text{MW} = \text{microwaves}$ .


esters **12a** and **12b**, respectively. The cyanoterpyridylpropionic acid **1e** was obtained when the hydrolysis time was reduced to one hour (Scheme 4, conditions c). Acid-catalyzed hydrolysis furnished the expected dicarboxylic acids **1b** and **1d** (Scheme 4, conditions b).

The binding of the new compounds to the  $\text{D}/\text{D}$  domain was evaluated in  $^{15}\text{N}$ -SOFAS-TM-QC experiments.<sup>[17]</sup> Compounds **1a–f** caused similar shifts or disappearances of signals as the peptides derived from AKAP18 $\delta$ , thus indicating that they bind to the same site on the  $\text{D}/\text{D}$  domain (see the Supporting Information).


Isothermal titration calorimetry revealed that **1b** and **1f** bind to the  $\text{D}/\text{D}$  domain with an estimated  $K_D = 148 \mu\text{M}$  and  $K_D = 31 \mu\text{M}$ , respectively (Figure 1). The binding is mainly driven by a large positive entropy change, presumably because of the release of water molecules from the binding interface. Compound **1d** bound to the  $\text{D}/\text{D}$  with a  $K_D$  value similar to that of **1f** (10–60  $\mu\text{M}$ ). However, the calorimetric data were of lower quality because of the small enthalpy change associated with this binding event (see Figure S4 in the Supporting Information).

The influence of our compounds on AKAP–PKA interactions was determined quantitatively in homogeneous time-resolved fluorescence (HTRF) assays. HTRF measurements (at pH 7) with the  $\text{D}/\text{D}$  domain of  $\text{RII}\alpha$  and AKAP18 $\alpha$  revealed that **1b** and **1e** decreased the FRET signal and thus inhibited the interaction ( $\text{IC}_{50} = 38 \mu\text{M}$  and 138  $\mu\text{M}$ , respectively; Figure 2). As expected, the peptide AKAP18 $\delta$ -L314E inhibited the interaction at a nanomolar concentration (12.7 nM),<sup>[6]</sup> while the inactive control peptide AKAP18 $\delta$ -PP did not affect the interaction (Figure 2). The inhibitory potency of **1a** and **1f** was too low to determine  $\text{IC}_{50}$  values.

Next, we examined the ability of **1b** to interfere with AKAP–PKA interactions in cells. HEK293 cells possess a negative feedback loop for terminating prostaglandin  $\text{E}_1$  ( $\text{PGE}_1$ ) induced cAMP synthesis, a pathway that depends on




**Figure 1.** Isothermal titration calorimetric measurements for determination of the  $K_D$  values for interactions of **1b** and **1f** with the D/D domain of RII $\alpha$ . Measurements consisted of successive 10  $\mu$ L injections of the D/D domain into a less concentrated solution of compound in a MicroCal VP-ITC instrument, with 1.0 mM D/D domain and 0.05 mM **1b**, or 0.6 mM D/D domain and 0.03 mM **1f**. The data were fitted to a one-site binding model.



**Figure 2.** Compounds **1b** and **1e** and peptide AKAP18 $\delta$ -L314E inhibit the interaction of the D/D domain of RII $\alpha$  with AKAP18 $\alpha$ . AKAP18 $\alpha$  lacks the 10 N-terminal amino acid residues which encode the membrane-targeting domain. Peptide AKAP18 $\delta$ -PP represents the RII binding domain of AKAP18 $\delta$ , but contains two proline residues which prevent interaction with the D/D domain. This peptide does not affect the interaction.<sup>[6]</sup> Homogeneous time-resolved fluorescence (HTRF) assays were carried out using the proteins in concentrations of 50 nM each.


AKAP–PKA interactions. Stimulation of the cells with PGE<sub>1</sub> leads to activation of adenylyl cyclase and the synthesis of cAMP; cAMP activates PKA, which phosphorylates cAMP phosphodiesterase PDE4D and thereby increases PDE activity and thus inhibits cAMP accumulation. PDE4D is phosphorylated by PKA bound to the AKAP gravin.<sup>[18]</sup> In addition, AKAP150 tethers PKA to adenylyl cyclase and facilitates PKA phosphorylation of the cyclase and thus its inactivation.<sup>[19]</sup> Disruption of AKAP–PKA interactions with peptides (Ht31), the small molecule FMP-API-1, or by PKA inhibition with H89 prevents the PGE<sub>1</sub>-dependent negative feedback and maintains high adenylyl cyclase activity in the



**Figure 3.** HEK293 cells were transfected to express cAMP-dependent cyclic nucleotide-gated channels (CNGC) and loaded with the Ca<sup>2+</sup> indicator Fura-2. The cells were left untreated, incubated with the PKA inhibitor H89, FMP-API-1, or compounds **1a** or **1b** (each 20  $\mu$ M) for 30 min. The cells were stimulated with prostaglandin E<sub>1</sub> (PGE<sub>1</sub>) to trigger cAMP synthesis at a time point of 60 s. cAMP opens the CNGCs, and then Ca<sup>2+</sup> enters the cells and binds to Fura-2. This binding generates fluorescence signals, which were imaged. Numbers in brackets indicate the numbers of individual cells measured.

presence of PGE<sub>1</sub>.<sup>[8,18,19]</sup> **1b** also inhibits the feedback inhibition, thus indicating that it interferes with AKAP–PKA interactions in cells (Figure 3). Although **1a** binds to the D/D domain (see Figure S3 in the Supporting Information), it did not interfere with the feedback inhibition, presumably because it does not reach sufficient concentrations in cells. The terpyridines do not inhibit the catalytic activity of PKA.

We have developed a set of terpyridines as  $\alpha$ -helix mimetics of the PKA anchoring disruptor peptide AKAP18 $\delta$ -L314E. Similar to the original peptide, the compounds bind to the D/D domain of RII subunits of PKA. One of them, **1b**, is the first nonpeptidic agent that binds to the D/D domain of RII subunits and inhibits AKAP–PKA interactions in vitro as well as in vivo. A model of the interaction is depicted in Figure 4. Evaluation of the structure–activity relationship (SAR) indicates that a COOH group as R<sup>2</sup> in **1b** is critically involved in defining its inhibitory potency; NH<sub>2</sub>, as in **1a**, is not tolerated at this position. Both H and OMe seem



**Figure 4.** Model of the interaction between terpyridine **1b** (cyan) and the D/D domain of human regulatory RII $\alpha$  subunits of PKA. The primary structure of the D/D domain is SHIQIPGLTELLQGYT-VEVLRQQPPDLVEFAVEYFTRLREAR (gene bank entry accession NP\_004148); the residues of the pocket are shown with side chains and a translucent surface (yellow: hydrophobic, blue: hydrophilic). The binding pocket is rather hydrophobic (I5, P6, L9, T10, L13, T17, L21) and contains the possible hydrogen-bond donors Q4 and Q14 at the rim (blue); Q4 and Q14 are predicted to interact with the carboxy groups of the terpyridine moiety of **1b**. For details see the Supporting Information.




## Communications



### Peptide Inhibitors

G. Schäfer, J. Milić, A. Eldahshan, F. Götz,  
K. Zühlke, C. Schillinger, A. Kreuchwig,  
J. M. Elkins, K. R. Abdul Azeez, A. Oder,  
M. C. Moutty, N. Masada, M. Beerbaum,  
B. Schlegel, S. Niquet, P. Schmieder,  
G. Krause, J. P. von Kries,  
D. M. F. Cooper, S. Knapp, J. Rademann,  
W. Rosenthal,  
E. Klussmann\* ————— ■■■■-■■■■

Highly Functionalized Terpyridines as  
Competitive Inhibitors of AKAP–PKA  
Interactions



**A good fit:** Interactions between A-kinase anchoring proteins (AKAPs) and protein kinase A (PKA) play key roles in a plethora of physiologically relevant processes whose dysregulation causes or is associated with diseases such as heart failure. Terpyridines have been developed as  $\alpha$ -helix mimetics for the inhibition of such interactions and are the first biologically active, nonpeptidic compounds that block the AKAP binding site of PKA.