

CHD2 variants are a risk factor for photosensitivity in epilepsy

Elizabeth C. Galizia,^{1,2,*} Candace T. Myers,^{3,*} Costin Leu,^{1,2,*} Carolien G. F. de Kovel,⁴ Tatiana Afrikanova,⁵ Maria Lorena Cordero-Maldonado,⁵ Teresa G. Martins,⁵ Maxime Jacmin,⁵ Suzanne Drury,⁶ V. Krishna Chinthapalli,^{1,2} Hiltrud Muhle,⁷ Manuela Pendziwiat,⁷ Thomas Sander,⁸ Ann-Kathrin Ruppert,⁸ Rikke S. Møller,^{9,10} Holger Thiele,⁸ Roland Krause,⁵ Julian Schubert,¹¹ Anna-Elina Lehesjoki,^{12,13} Peter Nürnberg,⁸ Holger Lerche¹¹ for the EuroEPINOMICS CoGIE Consortium,[#] Aarno Palotie,^{14,15,16} Antonietta Coppola,^{1,2,17} Salvatore Striano,¹⁷ Luigi Del Gaudio,¹⁷ Christopher Boustred,⁶ Amy L. Schneider,¹⁸ Nicholas Lench,⁶ Bosanka Jovic-Jakubi,^{19,20} Athanasios Covanis,²¹ Giuseppe Capovilla,²² Pierangelo Veggiotti,^{23,24} Marta Piccioli,²⁵ Pasquale Parisi,²⁶ Laura Cantonetti,²⁷ Lynette G. Sadleir,²⁸ Saul A. Mullen,²⁹ Samuel F. Berkovic,¹⁸ Ulrich Stephani,⁷ Ingo Helbig,⁷ Alexander D. Crawford,⁵ Camila V. Esguerra,^{30,31} Dorothee G. A. Kasteleijn-Nolst Trenité,⁴ Bobby P. C. Koeleman,^{4,\$} Heather C. Mefford,^{3,\$} Ingrid E. Scheffer^{18,29,\$} and Sanjay M. Sisodiya^{1,2,\$}

*,\$These authors contributed equally to this work.

#For details of the EuroEPINOMICS CoGIE Consortium see Appendix 1.

Photosensitivity is a heritable abnormal cortical response to flickering light, manifesting as particular electroencephalographic changes, with or without seizures. Photosensitivity is prominent in a very rare epileptic encephalopathy due to *de novo* CHD2 mutations, but is also seen in epileptic encephalopathies due to other gene mutations. We determined whether CHD2 variation underlies photosensitivity in common epilepsies, specific photosensitive epilepsies and individuals with photosensitivity without seizures. We studied 580 individuals with epilepsy and either photosensitive seizures or abnormal photoparoxysmal response on electroencephalography, or both, and 55 individuals with photoparoxysmal response but no seizures. We compared CHD2 sequence data to publicly available data from 34 427 individuals, not enriched for epilepsy. We investigated the role of unique variants seen only once in the entire data set. We sought CHD2 variants in 238 exomes from familial genetic generalized epilepsies, and in other public exome data sets. We identified 11 unique variants in the 580 individuals with photosensitive epilepsies and 128 unique variants in the 34 427 controls: unique CHD2 variation is over-represented in cases overall ($P = 2.17 \times 10^{-5}$). Among epilepsy syndromes, there was over-representation of unique CHD2 variants (3/36 cases) in the archetypal photosensitive epilepsy syndrome, eyelid myoclonia with absences ($P = 3.50 \times 10^{-4}$). CHD2 variation was not over-represented in photoparoxysmal response without seizures. Zebrafish larvae with *chd2* knockdown were tested for photosensitivity. *Chd2* knockdown markedly enhanced mild innate zebrafish larval photosensitivity. CHD2 mutation is the first identified cause of the archetypal generalized photosensitive epilepsy syndrome, eyelid myoclonia with absences. Unique CHD2 variants are also associated with photosensitivity in common epilepsies. CHD2 does not encode an ion channel, opening new avenues for research into human cortical excitability.

Received November 17, 2014. Revised December 15, 2014. Accepted January 7, 2015. Advance Access publication March 18, 2015

© The Author (2015). Published by Oxford University Press on behalf of the Guarantors of Brain.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

- 1 NIHR Biomedical Research Centre Department of Clinical and Experimental Epilepsy, UCL Institute of Neurology, National Hospital for Neurology and Neurosurgery, Queen Square, London, UK
- 2 Epilepsy Society, Bucks, UK
- 3 Department of Paediatrics, University of Washington, USA
- 4 Department of Medical Genetics Research, University Medical Centre Utrecht, The Netherlands
- 5 Luxembourg Centre for Systems Biomedicine, University of Luxembourg, Esch-sur-Alzette, Luxembourg
- 6 North East Thames Regional Genetics Laboratories, Great Ormond Street Hospital for Children NHS Foundation Trust, London, UK
- 7 Department of Neuropaediatrics, University Medical Centre Schleswig-Holstein and Christian-Albrechts-University of Kiel, Kiel, Germany
- 8 Cologne Centre for Genomics, University of Cologne, Cologne, Germany
- 9 Danish Epilepsy Centre, Dianalund, Denmark
- 10 Institute for Regional Health Services, University of Southern Denmark, Odense, Denmark
- 11 Department of Neurology and Epileptology, Hertie Institut for Clinical Brain Research, Tübingen, Germany
- 12 Folkhälsan Institute of Genetics and Neuroscience Centre, University of Helsinki, Helsinki, Finland
- 13 Research Programs Unit, Molecular Neurology, University of Helsinki, Helsinki, Finland
- 14 Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton, Cambridgeshire, UK
- 15 Institute for Molecular Medicine Finland, University of Helsinki, Helsinki, Finland
- 16 Program in Medical and Population Genetics and Genetic Analysis Platform, The Broad Institute of MIT and Harvard, Cambridge, USA
- 17 Epilepsy Centre, Neurology Department, Federico II University of Naples, Naples, Italy
- 18 Department of Medicine, University of Melbourne, Austin Health, Melbourne, Australia
- 19 Department of Child Neurology, Paediatric Clinic, Clinical Centre Nis, Serbia
- 20 Department of Paediatric Neurology, Paediatric Clinic, Al Sabah Hospital, Kuwait
- 21 Neurology Department, The Children's Hospital Agia Sophia, Athens, Greece
- 22 Epilepsy Centre 'C. Poma Hospital', Mantova, Italy
- 23 Department of Child Neurology and Psychiatry C. Mondino National Neurological Institute, Via Mondino, 2, 27100, Pavia, Italy
- 24 Brain and Behaviour Department, University of Pavia, Pavia, Italy
- 25 Neurophysiopathology Unit, San Filippo Neri Hospital, Rome, Italy
- 26 Child Neurology, NESMOS Department, Faculty of Medicine and Psychology, Sapienza University, Rome, Italy
- 27 Neurorehabilitation Unit, Department of Neuroscience and Neurorehabilitation, IRCCS, Bambino Gesù' Children's Hospital, Rome, Italy
- 28 Department of Paediatrics and Child Health, School of Medicine and Health Sciences, University of Otago, Wellington, New Zealand
- 29 Florey Institute of Neurosciences and Mental Health, and Department of Paediatrics, University of Melbourne, Royal Children's Hospital, Melbourne, Australia
- 30 Chemical Neuroscience Group, Biotechnology Centre of Oslo, University of Oslo, Oslo, Norway
- 31 Laboratory for Molecular Biodiscovery, University of Leuven, Leuven, Belgium

Correspondence to: Sanjay M Sisodiya,
Department of Clinical and Experimental Epilepsy,
UCL Institute of Neurology,
Queen Square,
London,
WC1N 3BG,
UK
E-mail: s.sisodiya@ucl.ac.uk

Correspondence may also be addressed to:
Ingrid E. Scheffer,
Epilepsy Research Centre,
Austin Health, 245 Burgundy St, Heidelberg,
Victoria 3081, Australia
E-mail: scheffer@unimelb.edu.au

Heather C. Mefford,
Department of Pediatrics,
Division of Genetic Medicine,
1959 NE Pacific St., Box 356320,
Seattle, WA 98195, USA
E-mail: hmefford@uw.edu

Bobby P. C. Koeleman,
 Medical Genetics,
 University Medical Center Utrecht,
 Universiteitsweg 100,
 3584 CG, The Netherlands
 E-mail: b.p.c.koeleman@umcutrecht.nl

Keywords: photosensitive; seizure; eyelid myoclonia with absences

Abbreviations: dpf = day post-fertilization; EMA = eyelid myoclonia with absences; ExAC = Exome Aggregation Consortium; GGE = genetic generalized epilepsy

Introduction

Photosensitivity is a heritable abnormal cortical response to flickering light, often manifesting as EEG changes called a photoparoxysmal response (Walter *et al.*, 1946). Photoparoxysmal response may occur with seizures, and in normal subjects, or with neuropsychiatric disorders (So *et al.*, 1993). The photoparoxysmal response is age-dependent: prevalence in healthy children is between 1.4 and 8.3%, dropping to <1% in adults (Gregory *et al.*, 1993; Quirk *et al.*, 1995; Kasteleijn-Nolst Trenite *et al.*, 2003; Verrotti *et al.*, 2012). Photosensitive epilepsy is a reflex epilepsy, with seizures triggered by visual stimuli. A population-based study in Great Britain determined that the annual incidence of epilepsy with photoparoxysmal response was 1.1 per 100 000 in the overall population, and 5.7 per 100 000 between 7 and 19 years of age (Quirk *et al.*, 1995). About 40% of people with photosensitive epilepsy only have seizures on exposure to visual stimuli. Photosensitive seizures also feature in specific epilepsy syndromes, with other seizure types, and in non-syndromic epilepsies. Examples include juvenile myoclonic epilepsy (Tauer *et al.*, 2005; Koeleman *et al.*, 2013; Taylor *et al.*, 2013), other genetic generalized epilepsies (GGE) (Taylor *et al.*, 2013), idiopathic photosensitive occipital epilepsy, and other focal (Taylor *et al.*, 2004; Lu *et al.*, 2008), symptomatic occipital, and progressive myoclonic, epilepsies. The archetypal photosensitive syndrome is eyelid myoclonia with absences (EMA), a GGE characterized by rapid eyelid jerks and upward eyeball deviation on eye closure: photosensitivity is an essential feature (Sadleir *et al.*, 2012).

The photoparoxysmal response is highly heritable (Waltz and Stephani, 2000; Tauer *et al.*, 2005; Taylor *et al.*, 2013). The genetics are complex: no single gene has been implicated despite linkage to several loci and formal meta-analysis (Tauer *et al.*, 2005; De Kovel *et al.*, 2010; Verrotti *et al.*, 2012). Photosensitive epilepsies also have complex genetic architecture (Sadleir *et al.*, 2012; Taylor *et al.*, 2013), with several linked loci (De Kovel *et al.*, 2010). Photosensitivity is a trait found in many syndromes, inheritable separately from epilepsy (Newmark and Penry, 1979). It is unclear whether isolated photoparoxysmal response is a risk factor for epilepsy (De Kovel *et al.*, 2010; Verrotti *et al.*, 2012).

Photosensitivity occurs in some epileptic encephalopathies, such as Dravet syndrome due to mutation in *SCN1A* and encephalopathy associated with mutation in *CHD2* (Carvill *et al.*, 2013). Published data do not allow determination of whether the photosensitivity in these conditions is due to the underlying gene mutation or to the epileptic encephalopathy *per se*. *CHD2* encodes chromodomain helicase DNA-binding protein 2, involved in transcriptional regulation. Additional attention was drawn to *CHD2* as a candidate photosensitive epilepsy gene as the only shared gene within several reported overlapping copy number variants of the chromosome 15q26.1 region associated with complex phenotypes including epilepsy with photosensitivity. Eight patients with *de novo* deletions of 15q26 encompassing part or all of *CHD2* have been reported (Veredice *et al.*, 2009; Dhamija *et al.*, 2011; Capelli *et al.*, 2012; Lund *et al.*, 2013; Mullen *et al.*, 2013; Chénier *et al.*, 2014). We and others subsequently showed 6/500 epileptic encephalopathy cases had *de novo* *CHD2* mutations (Carvill *et al.*, 2013; Epi4K Consortium *et al.*, 2013; Suls *et al.*, 2013; Lund *et al.*, 2014), and recently showed that clinical photosensitivity was prominent in the rare *CHD2*-associated myoclonic encephalopathy (Thomas *et al.*, 2015).

These findings led us to hypothesize that *CHD2* disruption would be associated with common forms of photosensitive epilepsy or photosensitivity manifesting as a photoparoxysmal response alone.

Materials and methods

Written informed consent was obtained from patients or parents/guardians for minors or those with intellectual disability. The study was approved by relevant institutional ethics committees.

We defined photosensitive epilepsy as the presence of a photoparoxysmal response (Kasteleijn-Nolst Trenite *et al.*, 2012) with a history of epilepsy, or seizures reproducibly induced by flickering light. The photoparoxysmal response *per se* was not an essential inclusion requirement in every patient with epilepsy because age, state (e.g. sleep deprivation) and antiepileptic medication affect its detectability. To test the effect of *CHD2* variation beyond the epileptic encephalopathies alone, we included a broad range of epilepsy types. Recruitment was from nine countries (see Supplementary

material for details) (Tauer *et al.*, 2005; Lu *et al.*, 2008; Taylor *et al.*, 2013). The cohort included 36 patients with EMA: all had photoparoxysmal response. We sequenced *CHD2* in 580 people with photosensitive epilepsy and 55 people with photoparoxysmal response but no history of seizures. All patients were of European ancestry. The phenotypic distribution is given in Table 1.

We evaluated data from two additional exome-sequenced cohorts of GGE patients, to determine the role of *CHD2* variation in GGE *per se*, independent of photoparoxysmal response. Not all patients in these cohorts had been formally assessed for photoparoxysmal response. These two groups were the Complex Genetics of Idiopathic Epilepsies Consortium (CoGIE) cohort of 238 probands with familial GGE (Supplementary material), and a published cohort of 118 patients with GGE (Heinzen *et al.*, 2012).

Targeted sequencing of *CHD2* was undertaken either using Illumina TruSeq Custom Amplicon™ (TSCA) or molecular inversion probes (see Supplementary material for details). Whole exome sequencing (Supplementary material) was performed on five EMA samples. Coverage data for all experiments are provided in the Supplementary material. Only variants confirmed by a second method (Sanger sequencing or a second independent molecular inversion probe capture, see Supplementary material) were used in analyses.

The Exome Aggregation Consortium (ExAC) formed a large control population of disease and population genetic studies (ExAC, Cambridge, USA; URL: <http://exac.broadinstitute.org> accessed October 2014; non-Finnish European samples only used), giving the best available population frequency of *CHD2* variants of interest. Detailed phenotypic data are not available for these individuals; some might, if tested, have or have had photoparoxysmal response or a history of photosensitive seizures. These unselected cases are unlikely to harbour more than the best estimates of photoparoxysmal response prevalence in the general population (1.4%) (Kasteleijn-Nolst Trenite *et al.*, 2003).

We focused on unique variants, in our cohort and in ExAC: this is a well-established approach (Carvill *et al.*, 2013; Cnossen *et al.*, 2014; Wain *et al.*, 2014). We hypothesized

an over-representation of unique variants in our cohort compared with the phenotypically-unselected ExAC cohort. We defined unique variants as those that occurred in one individual only, in cases and controls (from ExAC) considered together, that were non-synonymous, splice-site or frameshift. We used several methods for prediction of the functional consequences of unique variants in cases (Supplementary material). We defined ‘rare’ variants as those with a minor allele frequency <1% in the non-Finnish European ExAC samples.

We undertook functional studies. To test functional consequences of *Chd2* loss in zebrafish, we used the *chd2* E2I2 morpholino reported previously (Suls *et al.*, 2013). Briefly, morpholino (12 ng) was microinjected into 1- to 2-cell-stage embryos of the AB (wild-type) strain. Embryos were raised in a dark incubator. At 1 day post-fertilization (dpf), embryos were prepared using the least possible amount of light. In parallel, control non-injected embryos from the same clutch of eggs were processed in the same manner. At 4 dpf, optic tectal field recordings were performed (Suls *et al.*, 2013) (Supplementary material). The first 10 s of recording were performed in minimal light in order to place the needle. Immediately following these first 10 s, recordings were performed in the dark for five minutes. At the end of this 5-min period, a very bright light was switched on (‘light ON’ state; six times the standard brightness level used for needle placement), and recording continued for 5 min. A paroxysm of high-frequency activity (200–500 Hz) with amplitude >3 times background level, either spontaneous or evoked by light, was defined as a polyspiking episode.

Statistics

We performed a two-tailed Fisher’s exact test to determine whether the burden of unique variants in our case cohorts was greater than expected compared to ExAC controls. We examined the frequency of all rare variants in the entire cohort, and the frequency of unique variants only separately in patients with EMA, patients with GGE excluding EMA, and patients with focal epilepsies. The threshold for significance was set at $P < 0.01$, applying Bonferroni correction for these five comparisons. For the single separate comparison of cases with photoparoxysmal response without epilepsy and ExAC, significance was set at $P < 0.05$. For zebrafish data, comparison of the parameters of spiking activity (dark versus light condition) for each treatment group was performed using the Mann-Whitney test.

Results

We identified 22 rare variants (Supplementary Table 1) in the cohort of patients with photosensitive epilepsy: 11 were unique (Table 2). There was a significant difference ($P = 2.17 \times 10^{-5}$) in unique variant frequency between cases (11/580 cases; 11/1160 alleles; 0.95%) and controls (128/68 854 alleles; 0.19%). The unique variants in the cases were all well covered in ExAC controls (Supplementary material). The 11 unique variants in cases were also absent from additional data sets: Exome Variant Server (<http://evs.gs.washington.edu/EVS/>), 1000 Genomes data set (<http://www.1000genomes.org/>), and dbSNP

Table 1 Distribution of cases by continental origin and broad syndromic classification

Cohort	Syndrome			PPR without epilepsy
	GGE	Focal	Other	
European	249	24	32	55
Australian	230*	35*	11	0
Total	479*	59*	43	55

European includes epilepsy cases from Germany (90), Italy (82), The Netherlands (75), Greece (34), Serbia (17), UK (5) and Denmark (2).

GGE = genetic generalized epilepsies, including GGE for which other information was not available, and, where classified, juvenile myoclonic epilepsy, juvenile absence epilepsy, childhood absence epilepsy, early-onset absence epilepsy, epilepsy with myoclonic atonic seizures, epilepsy with generalized tonic-clonic seizures only, and EMA. Focal includes all types of focal epilepsies, including idiopathic photosensitive occipital lobe epilepsy (IPOE). *One Australian patient evolved from a GGE to a focal epilepsy. Other includes Lennox-Gastaut syndrome, epilepsy due to tuberous sclerosis, epilepsy with electrical status epilepticus in sleep and epilepsies otherwise unclassified: none of these particular cases had unique *CHD2* variants.

(<http://www.ncbi.nlm.nih.gov/SNP>). There was no difference in the overall burden of rare *CHD2* variants in cases compared to controls [22/1160 alleles (1.90%) versus 1236/68854 alleles (1.80%) respectively; $P = 0.74$]. We provide data on the frequency of variants in *CHD2* in cases and controls according to various thresholds in the Supplementary Table 2. Figure 1 shows all previously-reported variants and all unique variants identified in our cases.

We investigated the predicted deleteriousness of the 11 unique variants in the cases (Table 2). Eight of 11 unique variants (73%) had scaled CADD scores > 10 , placing them in the top 10% most deleterious single nucleotide variants; as a group, the 11 variants had a mean scaled CADD score of 32.6, ranking higher than 99.95% of all possible human single nucleotide variants (Kircher et al., 2014).

Next, we analysed variation by epilepsy type. The archetypal photosensitive GGE syndrome EMA had the highest frequency of unique variants, found in 3/36 patients, more than expected compared to ExAC controls (3/72 alleles versus 128/68 854 alleles) ($P = 3.50 \times 10^{-4}$). As a *post hoc* comparison, the frequency of unique variants (4.2%) in the small EMA group is considerably greater than in our overall cohort excluding EMA (0.74%) ($P = 0.026$).

Notably, two of three EMA variants were frameshift, compared to 9/128 unique variants in ExAC. One EMA variant was shown to have arisen *de novo*, strengthening its role in causation of the phenotype.

For all GGE excluding EMA, we found no significant difference compared to ExAC (4/888 alleles versus 128/68 854 alleles, $P = 0.089$). We also did not find significant over-representation in focal epilepsies compared with ExAC (2/118 alleles versus 128/68 854 alleles; $P = 0.021$). One case was included in both GGE and focal epilepsy cohorts, as the phenotype evolved from early-onset absence epilepsy to idiopathic photosensitive occipital epilepsy (Patient 11, Table 2). One of 55 (1.82%) individuals with photoparoxysmal response but no seizures had a unique *CHD2* variant (Table 2 and Fig. 1): this did not represent over-representation compared to ExAC (1/110 alleles versus 128/68 854 alleles; $P = 0.186$). This case has not developed epilepsy by the age of 18 years. We provide 99% confidence intervals (CI) (accounting for multiple comparisons) for all these comparisons in Table 3.

To investigate whether *CHD2* may be associated with the broader phenotype of GGE rather than photosensitive epilepsies specifically, we tested whether rare variants in *CHD2* were enriched in patients with GGE, with or

Table 2 Patients found to have unique mutations in *CHD2* and their clinical phenotypes

Case ID	Position (NCBI.37)	Consequence	cDNA change	Protein change	Computational Analysis Score (PolyPhen-2; SIFTindel; SIFT; splice-site inference)	CADD scores (PHRED scaled)	Syndromic diagnosis	Comments
1	15:93545502	Frameshift deletion	c.4233_4236del	p.E1412Gfs*64	Deleterious (0.858)	44	GGE	
2	15:93487750	Splice site	c.1153+5G>A	NA	No change in donor site	8.124	Unclassified	
3	15:93541780	Missense	c.C3937G	p.R1313G	Probably damaging (0.98)	16.9	Unclassified	
4	15:93543742	Missense	c.G4009T	p.A1337S	Benign (0.001)	8.728	IPOE	
5	15:93496808	Splice site	c.1719+5G>A	NA	Loss of donor site	15.74	Unclassified	Learning disability
6	15:93528855	Missense	c.G3365C	p.S1122T	Benign (0.01)	4.373	GGE	
7	15:93540316	Frameshift deletion	c.3725delA	p.K1245Nfs*4	Deleterious (0.858)	43	EMA	Autism; nephrolithiasis; migraine; scoliosis
8	15:93545442	Frameshift insertion	c.4173dupA	p.Q1392Tfs*17	Deleterious (0.85)	38	EMA	<i>De novo</i> mutation
9	15:93482909	Missense	c.C653T	p.P218L	Probably damaging (0.99)	21.3	EMA	Inherited from unaffected mother
10	15:93543767	Missense	c.G4034A	p.R1345Q	Possibly damaging (0.8)	33	JME	
11	15:93563244	Nonsense	c.C4909T	p.R1637X	Probably damaging (nonsense)	49	Phenotype evolved from early-onset absence epilepsy to IPOE	<i>De novo</i> mutation
i	15:93552396	Missense	c.G4435A	p.V1479M	Probably damaging (0.996)	27.9	PPR; febrile seizures only; no epilepsy	

IPOE = idiopathic photosensitive occipital epilepsy; JME = juvenile myoclonic epilepsy; PPR = photoparoxysmal response.

without photoparoxysmal response. Of 238 CoGIE GGE probands (Supplementary material), none had unique *CHD2* variants (not seen in ExAC or our cases). There were no unique mutations in *CHD2* in a previously-published cohort of 118 patients with GGE (Heinzen *et al.*, 2012).

To test functional consequences of *Chd2* loss in zebrafish, we used the *chd2* E212 morpholino reported previously (Suls *et al.*, 2013). As described, *chd2* morpholino-injected larvae displayed body curvature, excessive body pigmentation, and developmental delay (Suls *et al.*, 2013). This phenotype was observed after 50% knock-down of *chd2*. All non-treated larvae appeared normal. Recordings were obtained from 15 morpholino-injected larvae and 10 sibling controls. In comparison to 7 dpf larvae (Afrikanova *et al.*, 2013), spikes from 4 dpf larvae were shorter in duration and displayed a higher frequency of oscillations in polyspike complexes. Due to these differences, spontaneous spiking in controls was not excluded, but also quantified. We analysed duration of discharges, number of discharges under light conditions, cumulative duration of spiking activity, and cumulative discharge frequency distribution. Representative recordings are shown in Fig. 2.

In line with the previous findings (Suls *et al.*, 2013), the morpholino-injected larvae showed spontaneous abnormal burst discharges. There was a preferential occurrence during the light ON state (17 discharges in the dark versus 59 in the light). In the morpholino-injected group, 14/15 larvae had discharges during the light ON state; 7/15 larvae had spiking only during the 5-min light ON state, and 10/15 showed spiking activity within the first 3–5 s after the light ON. The average duration of any event (spike or polyspike discharge) in the morpholino-injected group fell during the light ON state (Fig. 3A), attributable to the fact that morpholino-injected larvae also displayed spontaneous polyspike discharges in the dark: the events

under light conditions were more heterogeneous (i.e. spontaneous polyspikes plus light-induced spiking), explaining reduced average duration. The average number of events/larva significantly increased in the morpholino-injected group in the light opposed to the dark period; this was not seen in the control group (Fig. 3B). A similar pattern was observed for cumulative duration of spiking activity (Fig. 3C): morpholino-injected larvae showed a steep increase in polyspike discharges in the light ON state, not observed for controls. The larvae from the non-injected control group also reacted to the light ON state by displaying an initial locomotor response, with 7/10 displaying short spontaneous burst activity within 2–13 s after the light was switched on. However, the overall distribution of event duration is different from that of morpholino-injected larvae (Fig. 3D): the controls' curve lies to the left of the morpholino-injected curve, indicating that the proportion of longer discharges is higher in the morpholino-injected group.

Table 3 Odds ratio for association with unique variants in *CHD2* by phenotype, with 99% CI

	P-value (Fisher's exact; 2-tailed)	Odds ratio	Lower bound of 99% CI	Upper bound of 99% CI
Whole photosensitive epilepsy cohort	2.17×10^{-5}	5.18	2.29	11.74
EMA alone	3.50×10^{-4}	24.36	5.06	117.38
GGE excluding EMA	0.089	2.44	0.65	9.08
Focal epilepsies	0.021	9.40	1.45	61.01
Cases with PPR only	0.186	4.96	0.36	67.74

The associations with photosensitive epilepsy overall and with EMA alone are significant, as documented in the text. PPR = photoparoxysmal response.

Distribution of *CHD2* variants




Figure 1 Schematic of *CHD2* illustrating its functional (chromo, DEXDc, DNA-binding and ATP helicase) domains, the location of previously-reported variants and the unique variants in both cases and controls identified in this study.

Discussion

We show an enrichment of unique variants in *CHD2* with photosensitivity in the common epilepsies overall, identifying *CHD2* as a photosensitive epilepsy gene. We also examined the distribution of unique variants by syndrome. *CHD2* is also the first gene to be discovered for EMA, the archetypal photosensitive epilepsy syndrome. In *CHD2* encephalopathy, though published phenotypes can be difficult to interpret, the seizure type of absence seizures with eyelid myoclonia, rather than the epilepsy syndrome, is seen in as many as 8/23 (35%) patients with *de novo* *CHD2* mutation or deletion (Veredice *et al.*, 2009; Dhamija *et al.*, 2011; Capelli *et al.*, 2012; Carvill *et al.*, 2013; Chénier *et al.*, 2014; Lund *et al.*, 2014). Together, these results suggest that *CHD2* is an important contributor to both the absence seizures with eyelid myoclonia seizure type and EMA epilepsy syndrome. For other epilepsy syndromes, *CHD2* variation over-representation in the photosensitive GGE or the mixed cohort of photosensitive focal epilepsies failed to meet the corrected threshold for significance. A single unique *CHD2* variant was found in one patient with photoparoxysmal response without seizures. In view of the comparatively small sizes of these syndrome cohorts, we can only confidently exclude effects with odds ratios greater than the upper limit for the 99% confidence intervals given in Table 3. Further studies in larger cohorts of these phenotypes would seem warranted.

Previous studies of photoparoxysmal response support a model of significant genetic heterogeneity and an overall complex genetic architecture (Sadleir *et al.*, 2012; Verrotti *et al.*, 2012; Taylor *et al.*, 2013): indeed, none of the

several linkage regions contain *CHD2*. Our findings confirm heterogeneity and complexity in the genetics of photosensitivity, but also suggest a single gene may contribute to photosensitivity in some cases. Two mutations we detected are recurrent: p.Glu1412Glyfs*64, previously reported in epileptic encephalopathy with marked photosensitivity (Carvill *et al.*, 2013); and p.Gln1392Thrfs*17, in Lennox-Gastaut syndrome with photosensitivity (Lund *et al.*, 2014). The unique variants detected are, as a group, predicted to be amongst the most deleterious variants possible (Kircher *et al.*, 2014) and *CHD2* is amongst the genes least tolerant of functional variation (Petrovski *et al.*, 2013; Residual Variation Intolerance Score 2.37).

CHD2 does not encode an ion channel, opening up new avenues for research into cortical excitability. *CHD2* is one of nine genes from a highly-conserved protein family with a unique domain combination: two N-terminal chromatin-organization modifier (chromo), SNF2-related helicase/ATPase and DNA-binding domains (Woodage *et al.*, 1997; Schuster *et al.*, 2002; Kulkarni *et al.*, 2008). *Chd2* knockdown zebrafish have multiple developmental abnormalities, abnormal movements and epileptiform discharges (Suls *et al.*, 2013). Disruption of *Chd2* in mice causes embryonic death in some heterozygote pups and a complex phenotype including growth retardation and lordokyphosis (Marfella *et al.*, 2006; Kulkarni *et al.*, 2008): epilepsy has not yet been described. Interestingly, the reported human mutations do not cluster to accessory domains of the protein and no obvious pattern has emerged. Recent data demonstrated that the N-terminal region of *CHD2* plays an inhibitory role, reducing DNA affinity and ATPase activity which may confer specificity, while the C-terminus enhances DNA binding and stimulates ATPase activity




Figure 2 Representative tectal field recordings of 4-dpf zebrafish larvae. Background fragment of non-treated wild-type control in the dark (A); reaction of a non-injected fish to light ON - movement artefacts (wavy background) and a very short spike were observed (B); response to light ON of the morpholino-injected larvae: significantly more spiking activity is seen (C). The scale is the same for all three fragments.

(Liu *et al.*, 2015). Additional studies investigating protein interacting partners and post-translational modifications of CHD2 will be necessary to understand how abnormal CHD2 leads to photosensitive epilepsy.

Our zebrafish data show that partial (50%) loss of *chd2* function causes photosensitivity. Although Suls *et al.* (2013) showed *chd2* knockdown could cause seizures, photosensitivity was not studied. Although normal zebrafish show complex sensitivity to light (Moore and Whitmore, 2014), and untreated larvae show minor sensitivity to sudden exposure to light, morpholino-injected larvae show significantly more spiking activity on sudden light exposure. Photosensitivity on constant, rather than only flickering, light exposure has been described in humans (Oguni *et al.*, 2001). The functional consequences of each of the human mutations we detected is not known,

but some at least very probably lead to loss of function, as caused by partial *chd2* knockdown that results in markedly enhanced photosensitivity in zebrafish. Together, these data strongly suggest that some human CHD2 mutations cause photosensitivity.

There are potential limitations of our work. Different sequencing platforms were used for the various studied groups. However, we note that all unique variants in cases were confirmed by a second method, whereas for ExAC controls we used a liberal threshold to maximize sensitivity to unique variants, such that a proportion of variants selected from ExAC will be false positive: the net result of this overall conservative approach is only to reduce study power. The ExAC cohort is also the biggest relevant control data set available, and the most likely of any existing data set to provide an accurate estimate of the




Figure 3 Electrophysiological activity of zebrafish larvae with *chd2* knockdown and light ON stimulus. Zebrafish larvae (4 dpf) were kept in the dark (or darkened environment, if not possible otherwise) for all groups in Danieau's medium. Tectal field recordings were performed for the first 5 min in the dark and subsequently in light ON state for the following 5 min in morpholino-injected larvae ($n = 15$) and non-injected larvae ($n = 10$). A spiking episode, either spontaneous or evoked by light, was defined as a paroxysm of high-frequency (200–500 Hz) activity with the amplitude exceeding three times the background. Average duration of spiking events \pm SEM detected per condition is shown in **A**. Average number of events per fish \pm SEM is shown in **B**. Cumulative duration of spiking activity per fish as seconds \pm SEM is shown in **C**. Cumulative frequency distribution of spiking episodes is shown in **D**: morpholino-injected larvae show more activity than any of the non-injected controls, and a higher photosensitivity (curve shift to the right in the light compared to the dark recordings). * $P < 0.05$ and ** $P < 0.01$ Mann-Whitney test.

true frequency of unique variation in *CHD2* in a population not enriched for photosensitive epilepsy. Taking all these factors into account, the use of different platforms is very unlikely to have generated false positive results—indeed, we are more likely to have underestimated unique variant numbers in cases. It is also possible that our choice of statistical test may have missed a true association between rare variation in *CHD2* and GGE (irrespective of photoparoxysmal response or photosensitivity), and we did not test whether *CHD2* variation contributes to epilepsy more broadly: we therefore cannot exclude the possibility that rare *CHD2* variation contributes to epilepsy *per se*. Lack of parental samples meant we could only confirm variants were *de novo* in two patients. Family samples were only available in one other case (Case 9): the variant was inherited from a clinically-unaffected mother in whom no EEG studies had been carried out.

Our results provide evidence for a specific gene in a particular trait in epilepsy. Understanding the genetic basis of the photosensitivity trait is a first step to elucidating the biology that underlies photoparoxysmal response and its relation to epilepsy. Human photosensitive epilepsy paradigms have facilitated epilepsy treatment discoveries (French *et al.*, 2014); understanding photoparoxysmal response biology may increase the value of these paradigms. Our findings may also provide new directions for understanding human cortical excitability.

Acknowledgements

We thank all the participants and their families. We thank the NE Thames Regional Genetics Service and the Institute of Neurology for the use of the laboratory facilities. The authors would like to thank the Exome Aggregation Consortium and the groups that provided exome variant data for comparison. A full list of contributing groups can be found at <http://exac.broadinstitute.org/about>. We thank Dr S. Balestrini for technical help.

Funding

Funding was provided by the Wellcome Trust (SMS, grant 084730), the Henry Smith Charity (SMS), Action Medical Research (SMS), National Institutes of Health (HCM R56NS69605), National Health and Medical Research Council of Australia (SFB, IES), Health Research Council of New Zealand (LGS), The Netherlands National Epilepsy Fund (grant 04-08 to B.P.C.K), German Research Foundation (HE 5415 3-1) within the EuroEPINOMICS framework of the Eurocores program by the European Science Foundation (ESF), intramural funds of the University of Kiel, Germany. H.L., P.N. and T.S. received grants by the EuroEPINOMICS programme (German Research Council, DFG grant numbers: HL: LE1030/11-1; PN: NU50/8-1, TS: SA434/5-1) within the

EUROCORES framework of the European Science Foundation (ESF); the collection of the CoGIE cohort was also supported by the EuroEPINOMICS program. Patient collection in Europe was partly made possible thanks to the European FP6 Marie Curie Excellence Grant on Visual Sensitivity MEXCT-CT-2005-024224. CVE is an Industrial Research Fund mandate holder of the KU Leuven. This work was partly undertaken at UCLH/UCL, which received a proportion of funding from the Department of Health's NIHR Biomedical Research Centres funding scheme.

Supplementary material

Supplementary material is available at *Brain* online.

References

- Afrikanova T, Serruys A-SK, Buenafe OEM, Clinckers R, Smolders I, de Witte PA, *et al.* Validation of the zebrafish pentylentetrazol seizure model: locomotor versus electrographic responses to antiepileptic drugs. *PloS One* 2013; 8: e54166.
- Capelli LP, Krepischi ACV, Gurgel-Giannetti J, Mendes MF, Rodrigues T, Varela MC, *et al.* Deletion of the *RMGA* and *CHD2* genes in a child with epilepsy and mental deficiency. *Eur J Med Genet* 2012; 55: 132–4.
- Carvill GL, Heavin SB, Yendle SC, McMahon JM, O'Roak BJ, Cook J, *et al.* Targeted resequencing in epileptic encephalopathies identifies *de novo* mutations in *CHD2* and *SYNGAP1*. *Nat Genet* 2013; 45: 825–30.
- Chénier S, Yoon G, Argiropoulos B, Lauzon J, Laframboise R, Ahn JW, *et al.* *CHD2* haploinsufficiency is associated with developmental delay, intellectual disability, epilepsy and neurobehavioural problems. *J Neurodev Disord* 2014; 6: 9.
- Cnossen WR, te Morsche RHM, Hoischen A, Gilissen C, Chrispijn M, Venselaar H, *et al.* Whole-exome sequencing reveals *LRP5* mutations and canonical Wnt signaling associated with hepatic cystogenesis. *Proc Natl Acad Sci USA* 2014; 111: 5343–8.
- De Kovel CGF, Pinto D, Tauer U, Lorenz S, Muhle H, Leu C, *et al.* Whole-genome linkage scan for epilepsy-related photosensitivity: a mega-analysis. *Epilepsy Res* 2010; 89: 286–94.
- Dhamija R, Brenningstall G, Wong-Kisiel L, Dolan M, Hirsch B, Wirrell E. Microdeletion of chromosome 15q26.1 in a child with intractable generalized epilepsy. *Pediatr Neurol* 2011; 45: 60–2.
- Epi4K Consortium, Epilepsy Phenome/Genome Project, Allen AS, Berkovic SF, Cossette P, Delanty N, Dlugos D, Eichler EE, *et al.* *De novo* mutations in epileptic encephalopathies. *Nature* 2013; 501: 217–21.
- French JA, Krauss GL, Kasteleijn D, DiVentura BD, Bagiella E. Effects of marketed antiepileptic drugs and placebo in the human photosensitivity screening protocol. *Neurotherapeutics* 2014; 11: 412–18.
- Gregory RP, Oates T, Merry RT. Electroencephalogram epileptiform abnormalities in candidates for aircrew training. *Electroencephalogr Clin Neurophysiol* 1993; 86: 75–7.
- Heinzen EL, Depondt C, Cavalleri GL, Ruzzo EK, Walley NM, Need AC, *et al.* Exome sequencing followed by large-scale genotyping fails to identify single rare variants of large effect in idiopathic generalized epilepsy. *Am J Hum Genet* 2012; 91: 293–302.
- Kasteleijn-Nolst Trenité D, Rubboli G, Hirsch E, Martins da Silva A, Seri S, Wilkins A, *et al.* Methodology of photic stimulation revisited: updated European algorithm for visual stimulation in the EEG laboratory. *Epilepsia* 2012; 53: 16–24.

Kasteleijn-Nolst Trenite D, Silva L, Maureza M. Prevalence of photoparoxysmal EEG responses in normal children and adolescents in Teofle Otoni, Brazil; 2001–2002. *Epilepsia* 2003; 44 (Suppl 8): 48.

Kircher M, Witten DM, Jain P, O’Roak BJ, Cooper GM, Shendure J. A general framework for estimating the relative pathogenicity of human genetic variants. *Nat Genet* 2014; 46: 310–5.

Koeleman BPC, de Kovel CGF, Kasteleijn-Nolst Trenité DGA. Photoparoxysmal EEG response and genetic dissection of juvenile myoclonic epilepsy. *Epilepsy Behav* 2013; 28 (Suppl 1): S69–71.

Kulkarni S, Nagarajan P, Wall J, Donovan DJ, Donnell RL, Ligon AH, et al. Disruption of chromodomain helicase DNA binding protein 2 (CHD2) causes scoliosis. *Am J Med Genet A* 2008; 146A: 1117–27.

Liu JC, Ferreira CG, Yusufzai T. Human CHD2 Is a Chromatin Assembly ATPase Regulated by Its Chromo- and DNA-binding Domains. *J Biol Chem* 2015; 290: 25–34.

Lu Y, Waltz S, Stenzel K, Muhle H, Stephani U. Photosensitivity in epileptic syndromes of childhood and adolescence. *Epileptic Disord* 2008; 10: 136–43.

Lund C, Brodtkorb E, Oye A-M, Røsby O, Selmer KK. CHD2 mutations in Lennox-Gastaut syndrome. *Epilepsy Behav* 2014; 33: 18–21.

Lund C, Brodtkorb E, Røsby O, Rødningen OK, Selmer KK. Copy number variants in adult patients with Lennox-Gastaut syndrome features. *Epilepsy Res* 2013; 105: 110–17.

Marfella CGA, Ohkawa Y, Coles AH, Garlick DS, Jones SN, Imbalzano AN. Mutation of the SNF2 family member Chd2 affects mouse development and survival. *J Cell Physiol* 2006; 209: 162–71.

Moore HA, Whitmore D. Circadian rhythmicity and light sensitivity of the zebrafish brain. *PLoS One* 2014; 9: e86176.

Mullen SA, Carvill GL, Bellows S, Bayly MA, Trucks H, Lal D, et al. Copy number variants are frequent in genetic generalized epilepsy with intellectual disability. *Neurology* 2013; 81: 1507–14.

Newmark ME, Penry JK. Photosensitivity and epilepsy: a review. New York: Raven Press; 1979.

Oguni H, Hayashi K, Awaya Y, Fukuyama Y, Osawa M. Severe myoclonic epilepsy in infants—a review based on the Tokyo Women’s Medical University series of 84 cases. *Brain Dev* 2001; 23: 736–48.

Petrovski S, Wang Q, Heinzen EL, Allen AS, Goldstein DB. Genic intolerance to functional variation and the interpretation of personal genomes. *PLoS Genet* 2013; 9: e1003709.

Quirk JA, Fish DR, Smith SJ, Sander JW, Shorvon SD, Allen PJ. Incidence of photosensitive epilepsy: a prospective national study. *Electroencephalogr Clin Neurophysiol* 1995; 95: 260–7.

Sadleir LG, Vears D, Regan B, Redshaw N, Bleasel A, Scheffer IE. Family studies of individuals with eyelid myoclonia with absences. *Epilepsia* 2012; 53: 2141–8.

Schuster EF, Stöger R. CHD5 defines a new subfamily of chromodomain-SWI2/SNF2-like helicases. *Mamm Genome* 2002; 13: 117–19.

So EL, Ruggles KH, Ahmann PA, Olson KA. Prognosis of photoparoxysmal response in nonepileptic patients. *Neurology* 1993; 43: 1719–22.

Suls A, Jaehn JA, Kecskés A, Weber Y, Weckhuysen S, Craiu DC, et al. De novo loss-of-function mutations in CHD2 cause a fever-sensitive myoclonic epileptic encephalopathy sharing features with Dravet syndrome. *Am J Hum Genet* 2013; 93: 967–75.

Tauer U, Lorenz S, Lenzen KP, Heils A, Muhle H, Gresch M, et al. Genetic dissection of photosensitivity and its relation to idiopathic generalized epilepsy. *Ann Neurol* 2005; 57: 866–73.

Taylor I, Berkovic SF, Scheffer IE. Genetics of epilepsy syndromes in families with photosensitivity. *Neurology* 2013; 80: 1322–9.

Taylor I, Marini C, Johnson MR, Turner S, Berkovic SF, Scheffer IE. Juvenile myoclonic epilepsy and idiopathic photosensitive occipital lobe epilepsy: is there overlap? *Brain J Neurol* 2004; 127: 1878–86.

Thomas RH, Zhang LM, Carvill GL, Archer JS, Heavin SB, Mandelstam SA, et al. CHD2 myoclonic encephalopathy is frequently associated with self-induced seizures. *Neurology* 2015. pii: 10.1212/WNL.0000000000001305.

Veredice C, Bianco F, Contaldo I, Orteschi D, Stefanini MC, Battaglia D, et al. Early onset myoclonic epilepsy and 15q26 microdeletion: observation of the first case. *Epilepsia* 2009; 50: 1810–15.

Verrotti A, Beccaria F, Fiori F, Montagnini A, Capovilla G. Photosensitivity: epidemiology, genetics, clinical manifestations, assessment, and management. *Epileptic Disord Int Epilepsy J Videotape* 2012; 14: 349–62.

Wain LV, Sayers I, Soler Artigas M, Portelli MA, Zeggini E, Obeidat M, et al. Whole exome re-sequencing implicates CCDC38 and cilia structure and function in resistance to smoking related airflow obstruction. *PLoS Genet* 2014; 10: e1004314.

Walter WG, Dovey VJ, Shipton H. Analysis of the electrical response of the human cortex to photic stimulation. *Nature* 1946; 158: 540.

Waltz S, Stephani U. Inheritance of photosensitivity. *Neuropediatrics* 2000; 31: 82–5.

Woodage T, Basrai MA, Baxevanis AD, Hieter P, Collins FS. Characterization of the CHD family of proteins. *Proc Natl Acad Sci USA* 1997; 94: 11472–7.

Appendix I

EuroEPINOMICS CoGIE Consortium

Aarno Palotie, Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton, Cambridgeshire, UK; Institute for Molecular Medicine Finland, University of Helsinki, Helsinki, Finland; Program in Medical and Population Genetics and Genetic Analysis Platform, The Broad Institute of MIT and Harvard, Cambridge, USA;

Anna-Elina Lehesjoki, Folkhälsan, Institute of Genetics, Helsinki, Finland; Research Programs Unit, Molecular Neurology, University of Helsinki, Helsinki, Finland; Neuroscience Center, University of Helsinki, Helsinki, Finland;

Ann-Kathrin Ruppert, Cologne Center for Genomics, University of Cologne, Cologne, Germany; Auli Siren, Outpatient Clinic for Persons with Intellectual Disability, Tampere University Hospital, Tampere, Finland;

Bobby Koeleman, Department of Medical Genetics, Division of Biomedical Genetics, University Medical Center Utrecht, Utrecht, Netherlands;

Dennis Lal, Cologne Center for Genomics, University of Cologne, Cologne, Germany;

Federico Zara, Laboratory of Neurogenetics, Pediatric Neurology and Muscular Diseases Unit, Department of Neurosciences, Gaslini Institute, Genova, Italy;

Felicitas Becker, Department of Neurology and Epileptology, Hertie Institute for Clinical Brain Research, University of Tübingen, Tübingen, Germany;

Hande Caglayan, Department of Molecular Biology and Genetics, Bogazici University, Istanbul, Turkey;

Helle Hjalgrim, Danish Epilepsy Centre, Dianalund, Denmark Institute for Regional Health Services, University of Southern Denmark, Odense, Denmark;

Hiltrud Muhle, University Medical Center Schleswig-Holstein, Christian-Albrechts University, Kiel, Germany;

Holger Lerche, Department of Neurology and Epileptology, Hertie Institute for Clinical Brain Research, University of Tübingen, Germany;

Holger Thiele, Cologne Center for Genomics, University of Cologne, Cologne, Germany;

Ingo Helbig, University Medical Center Schleswig-Holstein, Christian-Albrechts University, Kiel, Germany;

Janine Altmüller, Cologne Center for Genomics, University of Cologne, Cologne, Germany;

Julian Schubert, Department of Neurology and Epileptology, Hertie Institute for Clinical Brain Research, University of Tübingen, Tübingen, Germany;

Kamel Jabbari, Cologne Center for Genomics, University of Cologne, Cologne, Germany;

Kate Everett, Human Genetics Research Centre, St George's University of London, London, UK;

Pasquale Striano, Pediatric Neurology and Muscular Diseases Unit, Department of Neurosciences, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health, University of Genoa, G. Gaslini Institute, Genoa, Italy;

Patrick May, Luxembourg Centre for Systems Biomedicine, University of Luxembourg, Esch-sur-Alzette, Luxembourg;

Peter Nürnberg, Cologne Center for Genomics, University of Cologne, Cologne, Germany; Center for Molecular Medicine Cologne (CMMC), University of Cologne, Cologne, Germany; Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases (CECAD), University of Cologne, Cologne, Germany;

Rikke Møller, Danish Epilepsy Centre, Dianalund, Denmark; Institute for Regional Health Services, University of Southern Denmark, Odense, Denmark;

Rima Nabbout, Centre de Reference Epilepsies Rares, Inserm U1129, Neuropediatrics Department, Necker-Enfants Malades Hospital, APHP, Paris Descartes University, CEA, Orsay, France;

Roland Krause, Luxembourg Centre for Systems Biomedicine, University of Luxembourg, Esch-sur-Alzette, Luxembourg;

Rudi Balling, Luxembourg Centre for Systems Biomedicine, University of Luxembourg, Esch-sur-Alzette, Luxembourg;

Stephanie Baulac, Institut National de la Santé et de la Recherche Médicale U975, Centre de Recherche de l'Institut du Cerveau et de la Moelle Epinière, Hôpital Pitié-Salpêtrière, Paris, France; Centre National de la Recherche Scientifique, Centre de Recherche de l'Institut du Cerveau et de la Moelle Epinière, Hôpital Pitié-Salpêtrière, Paris, France; Université Pierre et Marie Curie (Paris VI), UMR_S 975, Paris, France;

Thomas Sander, Cologne Center for Genomics, University of Cologne, Cologne, Germany;

Wolfram Kunz Department of Epileptology and Life & Brain Center, University of Bonn Medical Center, Bonn, Germany;

Yvonne Weber, Department of Neurology and Epileptology, Hertie Institute for Clinical Brain Research, University of Tübingen, Tübingen, Germany.