

Cell Reports, Volume 41

Supplemental information









**Pan-claudin family interactome analysis
reveals shared and specific interactions**









Lorena Suarez-Artiles, Tilman Breiderhoff, Rossana Girardello, Hannes Gonschior, Sophie Rodius, Antoine Lesur, Ulf Reimer, Evelyn Ramberger, Daniel Perez-Hernandez, Dominik Müller, Philipp Mertins, and Gunnar Dittmar

Supplemental information

Pan-claudin family interactome analysis reveals shared and specific interactions

Lorena Suarez-Artiles, Tilman Breiderhoff, Rossana Girardello, Hannes Gonschior, Sophie Rodius, Antoine Lesur, Ulf Reimer, Evelyn Ramberger, Daniel Perez-Hernandez, Dominik Müller, Philipp Mertins, and Gunnar Dittmar

YFP_CLDN1 vs eGFP**YFP_CLDN2 vs eGFP****YFP_CLDN3 vs eGFP****YFP_CLDN4 vs eGFP****YFP_CLDN5 vs eGFP****YFP_CLDN6 vs eGFP****YFP_CLDN7 vs eGFP****YFP_CLDN8 vs eGFP**

YFP_CLDN9 vs eGFP**YFP_CLDN10 vs eGFP****YFP_CLDN11 vs eGFP****YFP_CLDN12 vs eGFP****YFP_CLDN14 vs eGFP****YFP_CLDN15 vs eGFP****YFP_CLDN16 vs eGFP****CFP_CLDN17 vs eGFP**













YFP_CLDN18.1 vs eGFP**YFP_CLDN18.2 vs eGFP****CFP_CLDN19a vs eGFP****YFP_CLDN19b vs eGFP****YFP_CLDN20 vs eGFP****YFP_CLDN22 vs eGFP****YFP_CLDN23 vs eGFP****YFP_CLDN24 vs eGFP**

Figure S2, related to Figure 2. Volcano plots showing the results from all CoIP experiments. LFQ intensity values of YFP/CFP-Claudin pull-downs were compared against the GFP control using moderated t-test. First significance level $\text{adj-pval} < 0.05$ (grey line), 2nd level cutoff of adj-pval that leaves only 5% of the interactions identified in the GFP control (blue line).












Figure S3, related to Figure 2. Confocal microscopy images of MDCK-C7 stable cell lines overexpressing recombinant YFP-/CFP- claudins. Anti-ZO-1 (61-7300 Invitrogen) and Alexa Fluor® 647 were used for tight junction staining.




Figure S4, related to Figure 3. Filtering strategy for SLiM-dependent interactors identified by PRISMA. **A.** Schematic representation of the SLiM-based intensity profile pattern used to filter significant PRISMA interactions. **B, C.** Example of the intensity profile of filtered (highly abundant unspecific proteins) and SLiM-dependent (CCT/TriC subunits) significant claudin interactors. **D.** Heatmap showing all the significant interactors identified by PRISMA with the SLiM-dependent interactors on top (based on their intensity profile) and the interactions that were removed after the filtering process.




Figure S5, related to Figure 3. Interacting partners of the unstructured C-termini of claudins identified by PRISMA. A. Bar plot corresponding to the normalized intensities of the significant interactions (p -value <0.05 and intensity binding profile) identified per peptide spot. **B.** Heatmap showing the interaction patterns of the significant proteins identified interacting with unmodified peptides derived from the C-terminal region of human claudins.



A**B**

Figure S6, related to Figure 4. Proximity ligation assay confirms interaction between claudin-23 and the mitochondrial protein MCU. **A.** Immunofluorescence images of PLA between MCU and claudin-23 or claudin-3. Caco2 cells were transiently transfected with YFP-tagged claudin-23 or claudin-3. MCU mitochondrial protein interacts with claudin-23 but not with claudin-3. **B.** Dot plot representing the number of PLA signals per transfected cell. Statistically significant results ($P < 0.05$) confirm those of the YFP-tagged claudin-23 CoIP.