

G_q-mediated arrhythmogenic signaling promotes atrial fibrillation

Felix Hohendanner, M.D. Ph.D.^{1,2,3}; Ashok Prabhu¹, Nicola Wilck, M.D.^{2,3,4,5,6}; Verena Stangl, M.D.^{2,3,7}; Burkert Pieske, M.D.^{1,2,3}; Karl Stangl, M.D.^{2,3,7}; Till F. Althoff, M.D.^{2,3,7,8,9}

¹ Charité – University Medicine Berlin, Campus Virchow-Klinikum, Department of Cardiology and German Heart Center, Augustenburger Platz 1, 13353 Berlin, Germany

² DZHK (German Centre for Cardiovascular Research), partner site Berlin, 13316 Berlin, Germany

³ Berlin Institute of Health at Charité – Universitätsmedizin Berlin, 10117 Berlin, Germany

⁴ Max Delbrück Center for Molecular Medicine in the Helmholtz Association (MDC), 13125 Berlin, Germany

⁵ Experimental and Clinical Research Center (ECRC), a cooperation of Charité - Universitätsmedizin Berlin and Max Delbrück Center for Molecular Medicine (MDC), 13125 Berlin, Germany

⁶ Department of Nephrology and Medical Intensive Care Medicine, Charité - Universitätsmedizin Berlin, 10117 Berlin, Germany

⁷ Charité – University Medicine Berlin, Department of Cardiology and Angiology, Charité Campus Mitte, Charitéplatz 1, 10117 Berlin, Germany


⁸ Arrhythmia Section, Cardiovascular Institute (ICCV), Hospital Clínic, Universitat de Barcelona, C/Villarroel N° 170, 08036 Barcelona, Catalonia, Spain

⁹ Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), 08036 Barcelona, Catalonia, Spain


Correspondence: althoff@clinic.cat

Short Title: G_q-signaling in atrial fibrillation

Supplementary Figures



Supplementary Figure S1. Per-animal analyses of Ca^{2+} signaling. Quantification of maximal cytosolic (A) and nuclear Ca^{2+} release (B), time to 50% of maximal cytosolic Ca^{2+} release (TF50) (C), the time constant of Ca^{2+} decay/removal (TAU) (D), subcellular Ca^{2+} spark frequency (E), as well as arrhythmic Ca^{2+} wave frequency (F) and propagation velocity (G). The total number of animals per group was $n=5$. * $p<0.05$ vs. WT.



Supplementary Figure S2. SR Ca²⁺ content in WT and KO mice as obtained with caffeine (top). Ca²⁺ spark frequency as corrected for SR Ca²⁺ content (ratio, a.u.).

*p<0.05 vs. KO.