



# Dietary protein and blood pressure: an umbrella review of systematic reviews and evaluation of the evidence

Heiner Boeing<sup>1</sup> · Anna M. Amini<sup>2</sup> · Julia Haardt<sup>2</sup> · Annemarie Schmidt<sup>2</sup> · Heike A. Bischoff-Ferrari<sup>3</sup> · Anette E. Buyken<sup>4</sup> · Sarah Egert<sup>5</sup> · Sabine Ellinger<sup>5</sup> · Anja Kroke<sup>6</sup> · Stefan Lorkowski<sup>7,8</sup> · Sandrine Louis<sup>9</sup> · Katharina Nimptsch<sup>10</sup> · Matthias B. Schulze<sup>11,12</sup> · Alexandra Schutkowski<sup>13</sup> · Lukas Schwingshackl<sup>14</sup> · Roswitha Siener<sup>15</sup> · Armin Zittermann<sup>16</sup> · Bernhard Watzl<sup>9</sup> · Gabriele I. Stangl<sup>13</sup> on behalf of the German Nutrition Society

Received: 14 July 2023 / Accepted: 22 January 2024  
© The Author(s) 2024

## Abstract

**Introduction** This umbrella review aimed to investigate the evidence of an effect of dietary intake of total protein, animal and plant protein on blood pressure (BP), and hypertension (PROSPERO: CRD42018082395).

**Methods** PubMed, Embase and Cochrane Database were systematically searched for systematic reviews (SRs) of prospective studies with or without meta-analysis published between 05/2007 and 10/2022. The methodological quality and outcome-specific certainty of evidence were assessed by the AMSTAR 2 and NutriGrade tools, followed by an assessment of the overall certainty of evidence. SRs investigating specific protein sources are described in this review, but not included in the assessment of the overall certainty of evidence.

**Results** Sixteen SRs were considered eligible for the umbrella review. Ten of the SRs investigated total protein intake, six animal protein, six plant protein and four animal vs. plant protein. The majority of the SRs reported no associations or effects of total, animal and plant protein on BP (all “possible” evidence), whereby the uncertainty regarding the effects on BP was particularly high for plant protein. Two SRs addressing milk-derived protein showed a reduction in BP; in contrast, SRs investigating soy protein found no effect on BP. The outcome-specific certainty of evidence of the SRs was mostly rated as low.

**Discussion/conclusion** This umbrella review showed uncertainties whether there are any effects on BP from the intake of total protein, or animal or plant proteins, specifically. Based on data from two SRs with milk protein, it cannot be excluded that certain types of protein could favourably influence BP.

**Keywords** Dietary protein · Blood pressure · Hypertension · Umbrella review · Grading of evidence · Meta-analyses

## Abbreviations

AMSTAR 2	A Measurement Tool to Assess Systematic Reviews 2
AS	Amino acids
BP	Blood pressure
CI	Confidence interval
CVD	Cardiovascular diseases
DASH	Dietary approaches to stop hypertension
DBP	Diastolic blood pressure
MAs	Meta-analysis/meta-analyses
RCT	Randomised controlled trial
SBP	Systolic blood pressure
SR	Systematic review

## Introduction

Hypertension is an important modifiable risk factor for cardiovascular, cerebrovascular and chronic kidney diseases, and the leading underlying cause of global mortality and disability [1, 2]. It is suggested that 62% of cerebrovascular diseases and almost 50% of the ischaemic heart diseases are attributable to elevated BP, which will affect almost one-third of the adult population worldwide [2].

The American Heart Association categorised the systolic BP (SBP) and diastolic BP (DBP) into four ranges: normal (SBP < 120 mmHg and DBP < 80 mmHg), elevated (SBP 120 to 129 mmHg and DBP < 80 mmHg), stage 1 hypertension (SBP 130–139 mmHg or DBP 80 to 89 mmHg) and stage 2 hypertension (SBP ≥ 140 mmHg or DBP ≥ 90 mmHg) [3]. The regulation of BP is controlled

Extended author information available on the last page of the article

by several complex mechanisms, such as baroreceptors, the activity of sympathetic nervous system, the renin–angiotensin–aldosterone system, antidiuretic hormone, natriuretic peptides and the nitric oxide system [4]. High BP which has emerged on the basis of medical conditions, such as renal diseases or endocrine disorders, is referred to as secondary hypertension. However, the most common form of hypertension is primary hypertension that is caused by a combination of genetic and lifestyle factors, such as obesity, physical inactivity, smoking and unhealthy diets [5]. In 2019, a network meta-analysis (MA) identified the Dietary Approaches to Stop Hypertension (DASH) diet, which favours a high intake of fruits and vegetables, low-fat dairy products, whole grains and low sodium, to be the most effective dietary strategy to reduce BP [6]. A recently published umbrella review, including 341 meta-analyses of randomised controlled trials (RCTs) and 70 meta-analysed observational studies, found high-quality evidence for a BP-lowering effect of the DASH diet. Additionally, the umbrella review demonstrated beneficial BP effects linked with the consumption of Mediterranean dietary patterns, which is characterised by low sodium, and moderate alcohol intake [7]. Notably, this umbrella review also included a few SRs on protein, revealing that high-protein diets were associated with BP-lowering effects in RCTs of low quality, but not in those of moderate quality. However, there is currently no published umbrella review focussing exclusively on the link between dietary proteins and BP.

Based on data showing that certain proteins may serve as a source of antihypertensive peptides [8], the hypothesis that dietary proteins can modulate BP appears biologically plausible. Most studies in the field of bioactive peptides have been published on milk peptides; among them, several peptides have been identified which can inhibit the angiotensin-converting enzyme (ACE) and lower BP [9, 10]. In addition to peptides, certain amino acids (AS) have been linked with mechanisms controlling BP and beneficial effects on elevated BP. For example, an MA of RCTs on the effect of L-arginine supplementation demonstrated a significant reduction in SBP of  $-6.40$  mmHg and DBP of  $-2.64$  mmHg and identified the effective dosage of L-arginine for SBP reduction to be  $\geq 4$  g per day [11].

It is therefore tempting to speculate that the intake of high-protein diets or proteins from plant and/or animal origin can modulate BP. The current umbrella review addressed the level and certainty of evidence derived from SRs concerning whether dietary intake of protein, and proteins from plant and animal sources in general are capable of modifying BP or hypertension risk in the general adult population. Further, proteins from specific food sources were also reviewed but not evaluated for evidence. The present umbrella review will contribute to the upcoming evidence-based guideline for

protein intake of the German Nutrition Society considering different pathologies.

## Methods

We conducted an umbrella review (PROSPERO: CRD42018082395) following the methodological protocol published by Kroke et al. [12]. This protocol was developed as part of the evaluation of protein intake and various health-related outcomes and was also used for BP. In preparing this manuscript, we followed the guidelines of reporting outlined in the PRISMA 2020 checklist [13]. The literature search, selection of SRs, data extraction and evaluation of the methodological quality and outcome-specific certainty of evidence was conducted independently by two authors (AMA, AnS). Any disagreements were resolved by discussion to reach consensus.

## Literature search

The systematic literature search was conducted in PubMed, Embase and Cochrane Database of Systematic Reviews for SRs published between 05/2007 and 10/2022 to cover a period of at least 10 years. The initial database search was conducted in 05/2017 and was updated on 6 October 2022 due to elapsed time reasons. The search strategies regarding study type (SRs), proteins (exposure or intervention) and BP in general, as well as SBP, DBP and hypertension, are presented in Supplementary Material S1. In addition to the SRs found in this context, reference lists of included SRs were screened for further SRs of relevance.

## Literature selection

Titles and/or abstracts of the results of the literature searches were screened according to pre-defined inclusion and exclusion criteria [12] in order to identify potentially eligible SRs. The full texts of potentially eligible records were retrieved and assessed for final eligibility.

SRs had to address the general adult population (without lactating women or top athletes) as inclusion criteria and were eligible for the umbrella review if they analysed one of the following study designs: SR with or without MA of prospective studies in humans (RCTs, prospective cohort studies, case-cohort studies or nested case–control studies). If an SR also included case–control studies or cross-sectional studies, those studies or MAs predominantly including those studies ( $\geq 50\%$  of all studies) were not considered. The SRs had to address the association/effect between protein intake and SBP, DBP or the incidence of hypertension. All SRs that exclusively meta-analysed studies with whole foods were excluded. From SRs addressing studies with whole foods,

only the studies which addressed proteins were considered in this umbrella review.

## Data extraction

The following data from each included SR were extracted: the first author's surname, year of publication, study type (e.g. SR with MA of RCTs), study duration(s), study population, intervention/exposure(s), outcome(s), effect estimate(s) including 95% confidence intervals (CIs), p-value(s) and heterogeneity estimate(s). Corresponding and first authors were contacted in case of insufficient data. Where results were reported from multiple analysis methods (e.g. MA conducted with both end of study and change values), we extracted all available results into Table 1. Subsequently, for the purpose of rating the overall certainty of evidence, content experts (HB and GIS) determined the selection of data to be utilised. Further, the utilised single original studies in each SR are listed in Supplementary Material S2, subdivided by study type and intervention/exposure.

## Assessments of methodological quality and outcome-specific certainty of evidence

The methodological quality of included SRs was assessed using a modified version of the "A Measurement Tool to Assess Systematic Reviews 2" tool (AMSTAR 2) [14], and the modifications are described in detail in our methodological protocol [12]. This version contains 14 evaluation items grading the methodology of SRs on a scale from high quality to critically low quality according to the presence of critical and non-critical methodological weaknesses (Supplementary Material S3). SRs graded as "critically low" by AMSTAR 2 were excluded from the evaluation of the overall certainty of evidence. The outcome-specific certainty of evidence of included SRs with and without MA was assessed using the NutriGrade scoring tool [15] (Supplementary Material S4). It utilises a numerical scoring system, and four categories rate the potential outcome-specific certainty of evidence: high, moderate, low and very low. The NutriGrade scoring tool was modified for the assessment of SRs without MA, and the adaptations are described in detail elsewhere [12]. For SRs reporting more than one relevant exposure or outcome, a separate assessment by NutriGrade was conducted.

## Rating of the overall certainty of evidence

The overall certainty of the evidence was assessed according to Kroke et al. [12] and is described in Supplementary Material S5. Kroke et al. [12] proposed using specific criteria for grading, including result concordance, existing biological plausibility, methodological quality and outcome-specific certainty of evidence. The assessment was performed for the intake of total protein, as well as proteins derived from animal and plant sources. SRs which addressed specific protein

sources, but not protein intake or animal and plant proteins in general, were included in this review, but not used to assess the overall certainty of evidence. The rating of the overall certainty of evidence was conducted independently by three authors (HB, AMA, GIS). Any disagreements were resolved by discussion to reach consensus.

## Results

The study selection process is outlined in the flow diagram depicted in Fig. 1. The literature search within the three databases identified 6850 potentially eligible publications, which were reduced to 5901 articles when duplicates were removed. 5730 publications were excluded due to irrelevant titles and abstracts. In total, 171 articles were subjected to full text screening. Out of the 171, 155 were found not to be eligible due to different reasons, which are shown in Supplementary Material S6. Most of the reasons of non-eligibility referred to irrelevant exposures or lack of exposure-outcome investigations fitting the research question. Three SRs were excluded because of a "critically low" AMSTAR 2 rating [16–18] (Supplementary Material S7). All three SRs used only one database for their literature search. Additionally, Altorf-van der Kuil et al. [16] and Tielemans et al. [18] failed to conduct an adequate risk of bias assessment and Dong et al. [17] failed to provide a list of excluded studies. A total of 16 SRs were considered eligible to be addressed in this umbrella review. Details of these studies (outcomes, rating according to methodological quality, outcome-specific certainty of evidence) are found in Table 1, which subdivides the SRs into total protein, animal and plant protein and those which compared animal with plant proteins. The detailed results of the assessment of the methodological quality are shown in Supplementary Material S7 and of the outcome-specific certainty of evidence in Supplementary Material S8. The duration of the underlying primary RCTs ranged from one week to two years. Approximately 10% had a duration of one to four weeks, while around 20% had a study duration of at least one year. The sample size of these underlying primary RCTs ranged between seven and 419 participants, with approximately 10% having fewer than 22 participants, around 20% having more than 100 participants, and approximately 10% having more than 150 participants. There were only eight underlying primary cohort studies, with a follow-up duration of 1.5–11.3 years. Their sample size ranged from 272 to 80,426 participants. Two cohort studies investigated fewer than 1000 participants, while five cohort studies had participant numbers ranging between 1361 and 5880. One cohort study had a relatively large number of participants, specifically 80,426.

Table 1 Characteristics of the included systematic reviews

References	Study type, study duration/follow-up	Study population	Exposition	Protein intake	Outcome	Pooled effect estimates (95% CI)	Heterogeneity estimator	NutriGrade rating	AMSTAR 2 rating
<i>(A) Total protein studies</i>									
Rehbolz et al. [19]	SR with MA of 29 RCTs published between 1980 and 04/11 Study duration: 1–24 wk	<i>n</i> = 2546 Both sexes Mean age: 26–62 yr (age range: 18–80 yr)	Protein vs. carbohydrates	Supplement intervention studies: 20–54 g/d vs. 0–8 g/d (control dose NP for all studies) Feeding/diet intervention studies: 19.5–95.0 En% vs. 11.5–50.0 En%	SBP  DBP	Net change: – 1.76 mmHg (– 2.33, – 1.20) <i>P</i> < 0.001  Net change: – 1.15 mmHg (– 1.59, – 0.71) <i>P</i> < 0.001	<i>I</i> <sup>2</sup> = 0%  <i>I</i> <sup>2</sup> = 0%	Moderate  Moderate	Low
Santesso et al. [20]	SR with MA of RCTs published before 08/11 Study duration: 28–365 d	Both sexes Mean age: 26–54 yr Healthy or with hypertension, overweight, obesity, hyperlipidemia or metabolic syndrome <i>n</i> = 1337	Higher- vs. lower-protein diets	25–35 En% vs. 15–24 En%	SBP  DBP	SMD: – 0.07 (– 0.21, 0.07) <i>P</i> = 0.33 SMD: – 0.03 (– 0.15, 0.09) <i>P</i> = 0.63	<i>I</i> <sup>2</sup> = 24%  <i>I</i> <sup>2</sup> = 9%	Low  Low	High
	MA conducted with end of study values: 21 RCTs				SBP	SMD: – 0.21 (– 0.32, – 0.09) <i>P</i> = 0.0004	<i>I</i> <sup>2</sup> = 0%	Low	
	MA conducted with change values: 15 RCTs	<i>n</i> = 1186			DBP	SMD: – 0.18 (– 0.29, – 0.06) <i>P</i> = 0.003	<i>I</i> <sup>2</sup> = 2%	Low	
Wycherley et al. [21]	SR with MA of 5 RCTs published before 05/11 Study duration: 4–16 wk	<i>n</i> = 230 Both sexes Mean age: 26 to ~ 62 yr Healthy or with overweight, obesity, type 2 diabetes or polycystic ovary syndrome	(High-protein, energy-restricted, low-fat diet) vs. (standard-protein, energy-restricted, low-fat diet) at least 10% difference in protein intake	28–33 En% (1.14–1.40 g/kg BW) vs. 16–21 En% (0.68–0.88 g/kg BW), this info was only provided for 4 out of 5 studies	SBP  DBP	WMD: – 2.09 mmHg (– 5.01, 0.83) <i>P</i> = 0.16 WMD: – 0.72 mmHg (– 2.67, 1.23) <i>P</i> = 0.47	<i>I</i> <sup>2</sup> = 0%  <i>I</i> <sup>2</sup> = 0%	Low  Low	Low
Pedersen et al. [22]	SR of 4 studies (1 RCTs, 2 cohort studies and 1 SR with MA) Studies published between 01/00 and 12/11 Study duration: NP	<i>n</i> = 41,170 Both sexes Healthy or with prehypertension or stage 1 hypertension	Total protein	This info was only available for 1 out of the 4 included studies: 25 vs. 15 En%	SBP, DBP and hypertension	All studies: non-significant association/effect	NA	Low <sup>a</sup> , SBP and DBP Moderate: hypertension	Moderate
Schwingshackl and Hoffmann [23]	SR with MA of 11 RCTs published before 08/12 Study duration: 1–2 yr	<i>n</i> = 1414 (SBP) <i>n</i> = 1402 (DBP) Both sexes Mean age: 41 to > 60 yr Healthy or with type 2 diabetes	High protein (≥ 25 En%) vs. low protein (≤ 20 En%) with both low fat (≤ 30 En%)	25–40 En% vs. 10–20 En%	SBP  DBP	WMD: – 1.61 mmHg (– 3.45, 0.23) <i>P</i> = 0.09 WMD: – 0.42 mmHg (– 1.37, 0.54) <i>P</i> = 0.39	<i>I</i> <sup>2</sup> = 41%  <i>I</i> <sup>2</sup> = 0%	Low  Low	High  Low
Clifton et al. [24]	SR with MA of 19 weight loss/maintenance RCTs published before 08/13 Study duration: 52–104 wk	<i>n</i> = 2650 Both sexes Mean age: 40–63 yr Healthy or with type 2 diabetes or polycystic ovary syndrome	High protein, low carbohydrate weight loss diet vs. control diet	25–60 En% vs. 10–30 En%	SBP  DBP	SMD: – 0.022 (– 0.13, 0.09) <i>P</i> = 0.69 SMD: 0.11 OR – 0.062 (– 0.23, 0.01) <i>P</i> = 0.104 OR 0.08 <sup>b</sup>	<i>I</i> <sup>2</sup> = 33%  <i>I</i> <sup>2</sup> = 52%	Low  Low	Moderate  Low

**Table 1** (continued)

References	Study type, study duration/follow-up	Study population	Exposition	Protein intake	Outcome	Pooled effect estimates (95% CI)	Heterogeneity estimator	NutriGrade rating	AMSTAR 2 rating
Mousavi et al. [25]	SR with MA of 5 cohort studies Studies published until 04/20 Follow-up: 2–11 yr	$n = 93,496$ (5,620 cases) Both sexes Age range: 18–65 yr General population	Total protein	NP		RR: 1.01 (0.90, 1.14) dose-response analysis (3 cohort studies): RR: 0.99 (0.83, 1.18) per 5% increase in energy intake from total protein $P_{\text{linear}} = 0.96$	$I^2 = 47\%$	Moderate	Moderate
Lonnie et al. [26]	SR without MA of 1 RCT RCT published before 03/20 Study duration: 3 wk	$n = 48$ Both sexes Mean age 58 yr With hypertension and overweight/obesity	Protein mix (pea protein, soya protein, milk protein and egg white protein) vs. maltodextrin vs. sucrose	NP	Postprandial SBP	Lower SBP with maltodextrin compared to protein mix No difference in SBP with sucrose compared to protein mix	NA	Very low	Low
Vogtschmidt et al. [27]	SR with MA of 25 RCTs RCTs published before 11/20 Study duration: 4–52 wk	$n = 1813$ Both sexes Mean age: 26–71 yr Healthy or with prehypertension, hypertension, overweight, obesity, insulin resistance, hyperlipidemia, hyperinsulinemia or polycystic ovary syndrome	Higher- vs. lower-protein diets (hypo- and isocaloric)	20–36 En% vs. 14–23 En%	SBP	Lower DBP with maltodextrin compared to protein mix No difference in DBP with sucrose compared to protein mix SMD: $-0.12$ ( $-0.21, -0.02$ ) $P$ NP	$I^2 = 0\%$	Low	High
Hengeveld et al. [28]	SR without MA of 4 RCTs RCTs published before 04/20 Study duration: 12 wk–2 yr	$n = 312$ Both sexes Mean age: 67–74 yr Older adults from the general population Healthy or with overweight, obesity or sarcopenia	High protein vs. low protein (2 RCTs with concomitant exercise in control and intervention group)	1.0–1.5 g/kg BW/d vs. 0.8–1.1 g/kg BW/d	SBP DBP	None of the RCTs showed an effect None of the RCTs showed an effect WMD: $-1.16$ mmHg ( $-2.13, -0.20$ ) $P$ NP	$I^2 = 0\%$	Low	Moderate
<i>(B) Animal protein studies</i>									
Rebbholz et al. [19]	SR with MA of 15 RCTs RCTs published between 1980 and 04/11 Study duration: NP	$n = NP$ Both sexes Mean age: NP	Animal protein (meat, fish, poultry, milk, casein or whey) vs. carbohydrates	Supplement intervention studies: 24.4–54.0 g/d vs. 0–2.8 g/d (control dose NP for all studies) Feeding/diet intervention studies: 19.5–95.0 En% vs. 15.2–50.0 En%	SBP	Net change: $-2.54$ mmHg ( $-3.55, -1.53$ ) $P < 0.001$	$I^2 = 0\%$	Moderate	Low
					DBP	Net change: $-0.95$ mmHg ( $-1.72, -0.19$ ) $P = 0.014$	$I^2 = 0\%$	Moderate	Moderate

Table 1 (continued)

References	Study type, study duration/follow-up	Study population	Exposition	Protein intake	Outcome	Pooled effect estimates (95% CI)	Heterogeneity estimator	NutriGrade rating	AMSTAR 2 rating
Pedersen et al. [22]	SR of 2 cohort studies published between 01/00 and 12/11 Study duration: NP	<i>n</i> = 7594 Both sexes Healthy	Animal protein	NP	SBP, DBP and hypertension	All studies: non-significant association	NA	Low: SBP, Low: DBP, Moderate: hypertension	Moderate
Chalvon-Demersay et al. [37]	SR without MA of 4 prospective studies 03/16 Follow-up: 1.5–11 yr	<i>n</i> = 4157 Both sexes Mean age: 42–70 yr Healthy or with hypertension, overweight, obesity, diabetes or hypercholesterolemia	Animal protein	NP	SBP & DBP	3 studies: no effect with animal protein 1 study: inverse relation between animal protein and BP	NA	Low <sup>c</sup>	Moderate
Hidayat et al. [35] <sup>d</sup>	SR with MA of 7 RCTs published before 05/16 Study duration: 4 wk–2 yr	<i>n</i> = 412 Both sexes Mean age: 23–61 yr, Healthy or with prehypertension, overweight, obesity, hypercholesterolemia or metabolic syndrome	Milk protein (low-fat milk, whey protein isolate, whey or casein, reduced-fat fortified milk or whey protein isolate and sodium caseinate) vs. (carbohydrates, usual diet, no supplementation or placebo)	70 mg–82 g extra milk protein/d	SBP	WMD: – 3.33 mmHg (– 5.62, – 1.03) <i>P</i> NP	<i>I</i> <sup>2</sup> = 0%	Low	High
Badelj et al. [36] <sup>e</sup>	SR with MA of 18 RCTs Literature search period: NP Study duration: 4–96 wk	<i>n</i> = 885 Both sexes Mean age: 23–74 yr With overweight or obesity	Whey protein (powder, supplement, isolate, hydrolyse, concentrate) vs. food protein, soy protein, (calcium) casein, casein + carbohydrates, glucose + lycopene, soy, low-fat conventional yogurt, placebo, control diet	0.7–110 g whey protein/d vs. 0–110 g non-whey protein/d	DBP	WMD: – 1.08 mmHg (– 3.38, – 0.22) <i>P</i> NP	<i>I</i> <sup>2</sup> = 0%	Low	Low
Mousavi et al. [25]	SR with MA of 5 cohort studies published until 04/20 Follow-up: 2–11 yr	<i>n</i> = 93,496 (5,620 cases) Both sexes Age range: 18–65 yr General population	Animal protein		DBP	WMD: – 5.69 mmHg (– 6.70, – 4.68) <i>P</i> < 0.001	<i>I</i> <sup>2</sup> = 99%	Low	Moderate
(C) Plant protein studies					Hypertension	RR: 1.03 (0.89, 1.19) Dose-response analysis (2 cohort studies): RR: 0.92 (0.60, 1.42) per 5% increase in energy intake from animal protein	<i>I</i> <sup>2</sup> = 68%	Moderate	Moderate
Rebbholz et al. [19]	SR with MA of 9 RCTs published between 1980 and 04/11 Study duration: NP	<i>n</i> = NP Both sexes Mean age: NP	vegetable protein (soy or other plant protein) vs. carbohydrates	Supplement intervention studies: 20–40 g/d vs. 0–8 g/d (control dose NP for all studies) feeding studies: 22.8–23.7 En% vs. 11.5–12.2 En%	SBP	Net change: – 2.27 mmHg (– 3.36, – 1.18) <i>P</i> < 0.001	<i>I</i> <sup>2</sup> = 5%	Moderate	Low

**Table 1** (continued)

References	Study type, study duration/follow-up	Study population	Exposition	Protein intake	Outcome	Pooled effect estimates (95% CI)	Heterogeneity estimator	NutriGrade rating	AMSTAR 2 rating
Pedersen et al. [22]	SR of 3 studies (2 cohort studies and 1 SR with MA) Studies published between 01/00 and 12/11 Study duration: NP SR without MA of 4 prospective studies 03/16 Follow-up: 1.5–11 yr	<i>n</i> = 9202 Both sexes Healthy or with prehypertension or stage I hypertension	Plant protein	This info was only available for 1 out of the 3 included studies: 18–66 g extra soy protein/d	DBP  SBP, DBP and hypertension	Net change: – 1.26 mmHg (– 2.26, – 0.26) <i>P</i> = 0.014  All studies: inverse association	<i>I</i> <sup>2</sup> = 22%  NA	Moderate  Low: SBP Low: DBP High: hypertension	Moderate
Chalvon-Demersay et al. [37]	SR without MA of 4 prospective studies 03/16 Follow-up: 1.5–11 yr	<i>n</i> = 4157 Both sexes Mean age: 42–70 yr Healthy or with hypertension, overweight, obesity, diabetes or hypercholesterolemia	Plant protein	NP	SBP and DBP	All studies: inverse relation between plant protein and BP	NA	Moderate <sup>d</sup>	Moderate
Mousavi et al. [25]	SR with MA of 5 cohort studies Studies published until 04/20 Follow-up: 2–11 yr	<i>n</i> = 93,496 (5620 cases) Both sexes Age range: 18–65 yr General population	Plant protein	NP	Hypertension	RR: 0.87 (0.74, 1.01) Dose-response analysis (2 cohort studies): RR: 0.76 (0.65, 0.90) per 5% increase in energy intake from plant protein	<i>I</i> <sup>2</sup> = 68%	Moderate	Moderate
Mohammadifard et al. [38] <sup>g</sup>	SR with MA of 2 RCTs RCTs published before 06/20 Study duration: 8–12 wk	<i>n</i> = 92 Females only Mean age: 48–64 yr With metabolic syndrome	Soya protein vs. control (“control” or DASH diet)	30–35 g/d	SBP	WMD: – 0.04 mmHg (– 0.33, 0.25)	<i>I</i> <sup>2</sup> = 62%	Very low	Low
Mosallanezhad et al. [39] <sup>h</sup>	SR without MA of 4 RCTs RCTs published before 06/20 Study duration: 5–24 wk	<i>n</i> = 605 Both sexes Mean age: 51–61 yr Healthy or with hypertension or hypercholesterolemia	Soya protein vs. control	NP	DBP  SBP	WMD: 0.13 mmHg (– 0.16, 0.42)  2 out of 4 comparisons: no difference between intervention and control 2 out of 4 comparisons: lower SBP with soya protein	<i>I</i> <sup>2</sup> = 25%  NA	Very low  Moderate	Moderate
(D) Animal vs. plant protein studies	SR with MA of 12 RCTs RCTs published between 1980 and 04/11 Study duration: 4–12 wk	<i>n</i> = 870 Both sexes Mean age: 36–65 yr (age range: 18–79 yr)	Vegetable protein (soy or other plant protein) vs. animal protein (meat, fish, poultry, milk, casein or whey)	Supplement intervention studies: 30–50 g/d vs. 40–50 g/d (control dose NP for all studies) Feeding/diet intervention studies: 12.6–21.2 En% vs. 12.9–20.7 En%	DBP	Net change: – 0.10 mmHg (– 2.31, 2.11) <i>P</i> NP	<i>I</i> <sup>2</sup> = 97%	Low	Low



Table 1 (continued)

References	Study type, study duration/follow-up	Study population	Exposition	Protein intake	Outcome	Pooled effect estimates (95% CI)	Heterogeneity estimator	NutriGrade rating	AMSTAR 2 rating
Chalvon-Demersay et al. [37]	SR without MA of 10 RCTs published before 03/16 Study duration: 4 wk–1 yr	$n = 1199$ Both sexes Mean age: 21–68 yr Healthy or with prehypertension, stage I hypertension, overweight, obesity or hypercholesterolemia	Animal vs. plant protein	1.5–90 g (additional) protein/d	DBP  SBP and DBP	Net change: – 0.24 mmHg (– 1.58, 1.10) $P$ NP 7 RCTs: no differential effect of plant- vs. animal-sourced proteins 3 RCTs: inconsistent effects	$I^2 = 78\%$  NA	Low  Moderate	Moderate
Lonnie et al. [26]	SR without MA of 3 RCTs published before 03/20 Study duration: 1 d–8 wk	$n = 105$ Both sexes Age range: 18–80 yr Healthy or with hypertension, overweight, obesity or hypercholesterolemia	Extracted plant protein (lupin or pea isolate) vs. animal protein (milk and/or egg white)	25 g/d or 0.6 g/kg BW/d	SBP	3 out of 4 comparisons: no difference between intervention and control 1 out of 4 comparisons: higher SBP with egg white protein compared to pea protein	NA	Low	Low
Bryant et al. [40]	SR without MA of 2 RCTs published between 01/00 and 09/21 Study duration: 28 d–8 wk	$n = 101$ Both sexes Mean age: 49–60 yrs All with hypercholesterolemia	Lupin protein isolate vs. milk protein isolate (isocaloric)	25 g/d	SBP	Both RCTs: no differences between intervention and control 1 out of 4 comparisons: higher DBP with egg white protein compared to pea protein	NA	Low	Moderate
					DBP	Both RCTs: no differences between intervention and control	NA	Low	Low

AMSTAR 2 A Measurement Tool to Assess Systematic Reviews 2,  $BP$  blood pressure,  $BW$  body weight,  $CI$  confidence interval,  $d$  day(s),  $DBP$  diastolic blood pressure,  $En\%$  percentage of energy intake,  $MA$  meta-analysis,  $mmHg$  millimetre of mercury,  $NA$  not applicable,  $NP$  not provided,  $RCT$  randomised controlled trial,  $RR$  relative risk,  $SBP$  systolic blood pressure,  $SMD$  standardised mean difference,  $SR$  systematic review,  $wk$  week(s),  $WMD$  weighted mean difference,  $yr$  year(s)

<sup>a</sup>A total of 4 NutriGrade assessments was conducted and all received a rating of low: RCTs: SBP and total protein, RCTs: DBP and total protein, cohorts: SBP and total protein, cohorts: DBP and total protein

<sup>b</sup>The manuscript states two different values for the effect estimator and the  $P$  value. The author was contacted for clarification, but no answer was received

<sup>c</sup>A total of 2 NutriGrade assessments were conducted (DBP and animal protein and SBP and animal protein), both received a rating of low

<sup>d</sup>This SR does not address animal protein in general, but specific protein sources and was therefore not used to assess the overall certainty of evidence

<sup>e</sup>This SR does not address animal protein in general, but specific protein sources and was therefore not used to assess the overall certainty of evidence

<sup>f</sup>A total of 2 NutriGrade assessments was conducted (DBP and plant protein and SBP and plant protein), both received a rating of moderate

<sup>g</sup>This SR does not address plant protein in general, but specific protein sources and was therefore not used to assess the overall certainty of evidence

<sup>h</sup>This SR does not address plant protein in general, but specific protein sources and was therefore not used to assess the overall certainty of evidence



## Total protein studies

Ten SRs addressed total protein intake and BP [19–28] (Table 1A). Two of these SRs included prospective cohort studies and investigated the association between total protein intake and BP or hypertension [22, 25]. The other SRs analysed only RCTs [19–21, 23, 24, 26–28]. The SRs with RCTs usually included individuals of both sexes and healthy subjects, but also accepted original studies that were conducted with individuals with overweight, hypertension and diabetes. High-protein diets used in the RCTs typically contained more than 25 energy% (E%) proteins. The quantity of proteins in the control diets mostly ranged from 10 to 20 E%. The study of Rebholz et al. [19] analysed only RCTs in which carbohydrates were replaced by protein.

The two SRs including prospective studies did not find associations between protein and BP or hypertension [22]. The SR of Pedersen et al. [22] included two prospective cohort studies [29, 30], but also a SR with MA [31], that analysed cross-sectional studies, but also two prospective cohort studies, one with young adults [32] and another with children [33]. The prospective cohort study with young adults found inverse, but predominantly non-significant associations between protein intake and BP [32]. The SR of Mousavi et al. [25] meta-analysed five cohort studies and did not find a significant association between total protein intake and risk of hypertension.

Nine SRs of RCT reviewed or meta-analysed the effect of total protein intake on BP [19–24, 26–28]. Rebholz et al. [19] found a reduction in SBP and DBP when carbohydrates were replaced by protein. The SR of Santesso et al. [20] observed no effect of protein on BP when comparing the final values between the intervention and control group (21 RCTs), but reported a significant reduction in BP following protein intake (15 RCTs) when comparing the final and baseline values. The SR of Wycherley et al. included five RCTs on BP and did not find significant effects of high-protein diets on SBP and DBP [21]. Schwingshackl and Hoffmann [23], conducting an MA of 11 RCTs addressing a similar research question, reached the same conclusion, revealing no effect of protein intake on SBP or DBP, neither in the MA of all RCTs, nor in the subgroup of high-quality RCTs [23]. No conclusive effects of dietary proteins on BP were found in the SR of Pedersen et al. [22]. An insufficient control of ethnicity and body weight were given as reasons for this conclusion [22]. Clifton et al., who meta-analysed 19 long-term (> 12 months) weight loss RCTs, did not observe a significant effect of dietary protein in exchange for carbohydrates

on SBP and DBP [24]. The SR of Lonnie et al. [26] included only one RCT that investigated the effect of total protein (mixture of pea protein, soy protein, milk protein, egg white protein) compared to maltodextrin or sucrose on postprandial BP [34]. This study found no protein effect on postprandial SBP. In contrast, DBP was significantly increased 60 min postprandial compared to maltodextrin, but not compared to sucrose intervention. The SR of Vogtschmidt et al. [27] included an MA of 25 RCTs and found high-protein diets (protein range: 20–36 E%) accompanied by a greater reduction in SBP than low-protein diets (14–23 E%), whereas the protein effect on DBP did not reach statistical significance. The SR of Hengeveld et al. [28], included four RCTs on protein intake and BP. None of these studies found an effect of an increased protein intake on SBP and DBP. The authors remarked critically that three of the four RCTs did not reach a sufficient statistical power to demonstrate an effect on BP.

The methodological quality assessed with AMSTAR 2 was graded as high for three SRs (all SRs of RCTs) [20, 23, 27], moderate for four SRs [22, 24, 25, 28] and low for three SRs [19, 21, 26] (Table 1A). The methodological quality of the SRs was independent of the publication date. The NutriGrade assessment included 23 entries separated according to the outcome investigated. Most of the ratings regarding the outcome-specific certainty of the evidence were low ( $n=17$ ). Only four exposure-outcome assessments were rated as moderate, and two assessments as very low. The list of studies being used in the SRs that demonstrate the potential study overlap is shown in Supplementary Material S2. The majority of RCTs on total protein were used also once ( $n=44$ ), while 29 RCTs were utilised multiple times (up to five times), mostly published between 2000 and 2005. Regarding cohort studies, there was only minor overlap (Supplementary Material S2B).

## Animal protein studies

Six SRs addressed animal protein and BP, with two of them analysing RCTs with milk proteins [35, 36]. Of the remaining four SRs on animal protein, three included cohort studies [22, 25, 37], and one analysed RCTs that replaced carbohydrates by animal protein [19]. Pedersen et al. [22] who analysed two cohort studies did not find an association between animal protein intake and BP. The SR of Chalvon-Demersay et al. [37] found in three out of four prospective studies no link between animal protein and BP, and an inverse association in one study. The formally well-performed quantitative

SR with MA of Mousavi et al. [25] including five cohort studies did not find animal protein intake associated with the risk of hypertension. The only SR with RCTs specifically addressing animal protein found that animal protein replacing carbohydrates led to significant reductions in SBP and DBP [19]. There are also two SRs of RCTs that addressed subtypes of animal proteins, in particular, proteins from milk. These were the SR of Hidayat et al. [35], who meta-analysed seven RCTs that investigated the effect of milk protein, in particular, whey protein and casein, on BP, and the SR of Badely et al. [36], who meta-analysed 18 RCTs that investigated the effect of whey protein on BP. Both SRs found a reduction in SBP and DBP following milk protein intake. The findings of these SRs are important with respect to health implications and dietary recommendations, but are not within the scope of the current review that aimed to investigate animal proteins in general. Thus, both SRs were not included in the evaluation of the overall certainty of evidence.

The outcome-specific certainty of evidence (NutriGrade rating) was rated four times as moderate and eight times as low, and the methodological quality (AMSTAR 2 rating) once as high [35], three times as moderate [22, 25, 37] and twice as low [19, 36]. The list of original studies being used in the SRs is shown in Supplementary Material S2C and S2D. The studies used by Pedersen et al. [22] were also used by Chalvon-Demersay et al. [37] and Mousavi et al. [25], but there was no overlap of studies in the SRs of Chalvon-Demersay et al. [37] and Mousavi et al. [25].

### Plant protein studies

Six SRs addressed plant proteins and BP [19, 22, 25, 37–39] (Table 1C). One SR analysed RCTs that replaced carbohydrates by plant proteins [19], three SRs analysed observational studies including cohort studies [22, 25, 37] and two SRs analysed soy products which included also studies with soy protein [38, 39]. According to the procedure of SRs with milk proteins, SRs which exclusively addressed soy were not used for the evaluation of the overall certainty of evidence of plant proteins on BP. Pedersen et al. [22] concluded that their SR provided evidence for an inverse relationship between plant protein and BP. This conclusion was based on two cohort studies, that found an inverse relationship between plant protein and BP and a MA of RCTs addressing soy protein, which show a BP-lowering effect of these plant proteins [17]. This MA was excluded from this umbrella review due to its low methodological quality [17]. In the SR of Chalvon-Demersay

et al. [37], the four cohort studies showed an inverse relationship between plant protein intake and SBP and DBP, respectively. Mousavi et al. [25] concluded in their SR that plant protein intake was not associated with risk of hypertension, although a subset of dose–response studies observed an inverse relationship between plant proteins and BP. Rebholz et al. [19] observed an inverse relation in RCTs when carbohydrates were replaced by plant proteins.

In the SR of Mohammadifard et al. [38] which addressed health effects of soy in subjects with the clinical diagnosis of metabolic syndrome, only two RCTs focussed on the effects of soy protein on BP. These two RCTs did not find any effect of soy protein on SBP and DBP, and were in line with the overall findings of consumption of soy products on BP in this SR. The SR on soy conducted by Mosallanezhad et al. included four RCTs on soy protein [39], two of them showed a BP-lowering effect and the other two observed no effect on BP.

The outcome-specific certainty of evidence was rated twice as very low, two times as low, seven times as moderate and once as high. The methodological quality was considered to be moderate in three [22, 25, 37] of the four SRs and low in one [19]. There had been a moderate overlap of studies included in the SRs (Supplementary Material S2E and S2F).

### RCTs with animal vs. plant protein

Four eligible SRs (Table 1D) included studies that compared the effects of animal proteins with plant protein intake on BP [19, 26, 37, 40]. The SR of Rebholz et al. [19] included 12 RCTs that compared plant with animal proteins on BP, but found no differences between these two protein sources. No differences between plant and animal proteins on BP were observed in the SR of Chalvon-Demersay et al. [37] either, which included 10 RCTs, most of them with soy protein as the plant protein source. The SR of Lonnie et al. [26] included three relevant RCTs, two of them did not show differences between animal and plant proteins on SBP and DBP, and one RCT found higher postprandial BP values following egg white protein consumption compared to plant protein intake. Bryant et al. [40] analysed two RCTs in their SR on subjects with hypercholesterolemia and found that lupin protein isolates and milk protein did not differ in their effect on SBP and DBP.

The ratings regarding NutriGrade and AMSTAR 2 referred to all four SRs and, with one exception, achieved a rating of low regarding outcome-specific certainty of

evidence and a split between low and moderate regarding methodological quality (Table 1D). There was some overlap of studies being used in the SRs (Supplementary Material S2G).

### Grading of the overall certainty of the evidence

Twelve SRs were used to grade the evidence of whether total protein and the subtypes animal and plant protein affect BP (Table 2). Four SRs [35, 36, 38, 39] were excluded from the evidence grading as they examined specifically milk and soy proteins.

Most of the ten SRs on total protein and BP reached at least a moderate methodological quality, a low rating of the outcome-specific certainty of evidence, and found no effect on BP (Table 2). Thus, the overall certainty of evidence regarding the BP-modulating influence of total protein was rated “possible” for having no effect on BP.

The four SRs on animal protein were mostly of moderate methodological quality and outcome-specific certainty of evidence, and the vast majority of the SRs found no effect of animal protein on BP (Table 2B). An exception was the SR by Rebholz et al. [19], which analysed a specific research question, namely, the replacement of carbohydrates with total protein or animal and plant protein. They found that such replacement reduced BP at all instances (total protein, animal and plant protein) (Table 2A–C). Overall, we concluded for animal protein that there is “possible” overall certainty of evidence for no effect.

Concerning the link between plant protein and BP, the majority of SRs analysing cohort studies showed an inverse association between plant protein intake and BP (Table 2C). However, this finding was counterbalanced by the four SRs of RCTs showing no relationship when comparing animal and plant protein directly, and was given greater weight for the assessment of the overall certainty of evidence than the SRs of cohort studies. Therefore, we concluded that there is no relation. Most of the SRs received moderate ratings in terms of methodological quality and outcome-specific certainty of evidence. Consequently, we graded the causal link between plant protein and BP as “possible” for no effect.

### Discussion

This umbrella review systematically evaluated the evidence on the role of dietary protein on BP or hypertension, yielding “possible” evidence for no link between total protein, animal protein and plant protein intake and BP. However, SRs which

analysed exclusively special types of proteins such as milk or soy proteins were not included in the grading of the overall certainty of evidence. These proteins may have effects on BP that are different from those of total, animal or plant proteins in general. The grading of evidence regarding the role of dietary protein on BP in the current umbrella review is in line with findings of a recently published umbrella review on the role of diet in the prevention and management of hypertension [7]. This umbrella review indicated that the evidence for dietary protein overall, as well as for animal or vegetable protein, is of low quality.

The grading of the overall certainty of evidence depends not only on the conduct of the SRs but also on the availability and quality of the original studies. Both aspects are critical with respect to dietary protein and BP. Out of the 16 eligible SRs, six did not perform a formal MA [22, 26, 28, 37, 39, 40], and two SRs utilised only a subset of studies (see method section, Supplementary Material S7). The vast majority of SRs were rated “low” in terms of their outcome-specific certainty of evidence, despite the predominantly moderate or high methodological quality of most SRs. The reasons for the low outcome-specific certainty of evidence grading in most SRs resulted from the low number of included primary studies, which can cause publication bias and/or heterogeneity, and the existence of a potential conflict of interest (Supplementary Materials S7 and S8). In terms of the quality of original studies, we noted that many existing dietary cohort studies that addressed associations between diets and pathologies, evaluated data regarding BP change or incidence of hypertension, but failed to address specifically the association between protein intake and BP. BP is difficult to analyse in an observational setting due to the many factors affecting BP, and it is challenging to define the clinical diagnosis of primary hypertension because of the widespread use of antihypertensive medication and the presence of other diseases. In addition, some cohort studies have failed to consider important confounding factors for BP, such as other dietary factors associated with protein consumption. The critical remarks on the conduct of observational studies can also be applied to RCTs. Many RCTs did not include information on the use and type of medication or did not consider the effects of antihypertensive drugs. For example, in the SR of Rebholz et al. only 15 of 32 RCTs included subjects without BP-lowering medication [19]. Another factor that has a strong effect on BP is weight reduction. Many SRs included weight loss studies, although weight reduction is known to lower BP [41–43] and could dominate the presumed protein effect on BP.

In addition, another limitation is that the majority of studies included in the SRs used self-reports on dietary protein intake, although protein intake can be determined more precisely by renal nitrogen excretion [44, 45]. In view of the potential biases associated with self-reports of dietary intake, more studies using biomarkers or controlled protein applications are highly warranted. The latter aspect is important because many clinical trials studies used foods rich in protein and not protein isolates to address the protein effect on BP. Many trials advised the study participants to consume a diet high in protein from meat, fish, eggs or other animal sources (often referred to as an omnivorous diet), while the control diets were often more in line with a vegetarian diet. An umbrella review of MA of interventional and observational studies shows that individual food groups and dietary patterns can influence BP very differently due to their ingredients such as sodium, potassium, magnesium, plant compounds and fatty acids [7]. Thus, the use of proteins from food sources as an intervention measure instead of purified protein isolates could compromise the findings by providing additional bioactive compounds affecting BP, for example, isoflavones with soy protein. While such a phenomenon of confounding effects could in principle be addressed in RCTs, the seriousness of such bias is much greater in observational studies. In the past, information on bioactive compounds was not available in nutrient tables and thus could not be considered in the statistical analyses. Observational studies, particularly those addressing plant proteins without considering the potential effects of a high vegetable and fruit intake on BP [46] and for BP-modulating dietary bioactive compounds, have a high chance of confounding bias if showing an inverse association. In the grading of the evidence of an effect of plant protein on BP, we have thus weighted the RTCs comparing animal and plant proteins higher than findings of an inverse association in cohorts. Further, we excluded SRs from our evaluation of the evidence that addressed proteins from specific food sources.

In addition, RCTs typically involve a treatment group receiving additional dietary protein which is replaced totally or in part in the control group by similar quantities or energy-adjusted amounts of non-protein macronutrients such as carbohydrates or fat, or by providing different types of proteins in the intervention and control groups. Most studies investigated the effect of increasing dietary protein intake in the range of > 25 E%, but not the effect of reducing dietary protein intake from the current range of 15–20 E% to 10–12 E%, reflecting the protein requirement. Moreover, the replacement of other macronutrients by protein as being specifically analysed by Rebolhol et al. [19] raises the question


of whether the increasing protein intake or reducing, e.g. carbohydrates, causes the effect.

In contrast to total protein, animal protein and plant protein, proteins from specific food sources such as milk proteins are described to be efficient in BP reduction. Interestingly, SRs exclusively focussing on milk proteins, such as whey protein or casein hydrolysate, demonstrate favourable effects on BP [47, 48]. This is in line with a recently published umbrella review that found moderate quality evidence for a BP-lowering effect of lactotripeptides and a lower prevalence of hypertension associated with low-fat dairy, milk and fermented dairy consumption [7]. These results are in accordance with mechanistic data that identified specific peptides to be capable of modulating BP. A recent MA of 12 RCTs on food–protein–derived peptides found pooled effects of peptide intervention on SBP and DBP to be – 3.28 mm Hg and – 1.82 mm Hg, respectively [49]. Most peptides used in this MA were derived from milk and milk products, such as the casein-derived tripeptides valine–proline–proline and isoleucine–proline–proline, whey-derived lactokinins and fragments of  $\alpha$ - and  $\beta$ -lactoglobulin which can inhibit ACE, thereby reducing the synthesis of angiotensin II and vasoconstriction [50–52]. These findings emphasise the role of specific peptides in lowering BP, and the need for more studies addressing peptides, rather than proteins in total.

The strengths of this umbrella review are (1) the standardised methodical procedures to include the entire appropriate literature, (2) the systematic literature search in three literature databases that aimed to include all relevant SRs and (3) the evaluation of the methodological quality as well as the evaluation of the outcome-specific certainty of evidence of the included SRs.

A weakness of this umbrella review is that the overall certainty of evidence is mainly based on data of SRs that included very heterogeneous RCTs in terms of the study population, study design, protein intake and control interventions, while methodically well-conducted SRs of observational studies are under-represented. It is crucial to note that the RCTs included in the SRs, which may not have specifically measured BP as the primary outcome, pose the risk of being underpowered to detect BP-related effects. This circumstance is also apparent in the assessment of the outcome-specific certainty of the evidence. In fact, about half of the outcome-specific NutriGrade assessments received a score of 0 points for the precision domain, due to low sample size (e.g. < 400 participants for a meta-analysis of RCTs) and/or wide 95% CIs, indicating a potential power issue. Furthermore, the best tool for rating the outcome-specific certainty of evidence warrants

Fig. 1 Flow diagram



discussion. In our umbrella review, we chose NutriGrade, specifically developed to address the unique requirements of nutrition research [53]. Notably, the GRADE approach has undergone subsequent amendments and may emerge as the primary tool in the future. Generally, by considering all SRs from the last 10 years as commonly done in umbrella reviews earlier published studies are over-represented. We addressed this issue in Supplementary Material S2, where the original study overlap is explored. The analysis proved insightful, revealing that there was only a moderate overlap among the SRs regarding the primary studies. Finally, to mitigate the risk of overlooking relevant SRs published recently, an updated literature search was conducted in PubMed in November 2023, using our original search strategy. The search identified two additional SRs, of which one included only three RCTs, which were already considered in other SRs included in our umbrella review [54]. The second SR, specifically addressing milk

proteins, is discussed above [47]. Importantly, the findings from these additional SRs do not alter the key messages of our umbrella review.

## Conclusion

This umbrella review showed uncertainties regarding the link between BP and the intake of total protein, as well as animal or plant proteins specifically. The methodological quality of the SRs ranged from low to high, and the outcome-specific certainty of evidence was mostly low. Future high-quality RCTs using well-characterised study populations, defined quantities of proteins or valid assessments of protein intake and iso-energetic control interventions are warranted to provide high-quality evidence and a solid basis for recommendations on dietary protein and BP.

**Table 2** Overview of the systematic reviews used to grade the overall certainty of the evidence for the link between blood pressure and total, animal and plant protein

	Systematic review	Included study type	Protein and BP	NutriGrade rating	AMSTAR 2 rating
<i>(A) Total protein</i>					
	Hengeveld et al. [28]	RCTs	∅	SBP: low DBP: low	Moderate
	Vogtschmidt et al. [27]	RCTs	↓: SBP ∅: DBP	SBP: low DBP: low	High
	Lonnie et al. [26]	RCTs	∅	SBP: very low DBP: very low	Low
	Mousavi et al. [25]: high vs low analysis	Cohorts	∅	Hypertension: moderate	Moderate
	Clifton et al. [24]	RCTs	∅	SBP: low DBP: low	Moderate
	Pedersen et al. [22]	Cohorts and RCTs	∅	SBP: low DBP: low Hypertension: moderate	Moderate
	Schwingshackl and Hoffmann [23]	RCTs	∅	SBP: low DBP: low	High
	Wycherley et al. [21]	RCTs	∅	SBP: low DBP: low	Low
	Rebholz et al. [19]	RCTs	↓	SBP: moderate DBP: moderate	Low
	Santesso et al. [20]: analysis with end of study values	RCTs	∅	SBP: low DBP: low	High
	∑	<i>n</i> = 10 SRs on total protein: 9 ∅, 2 ↓ Majority of SRs reach at least a “low” NutriGrade rating Majority of SRs reach at least a “moderate” AMSTAR 2 rating			
<i>(B) Animal protein</i>					
	Mousavi et al. [25]: high vs low analysis	Cohorts	∅	Hypertension: moderate	Moderate
	Chalvon-Demersay et al. [37]	Cohorts	∅	SBP: low DBP: low	Moderate
	Pedersen et al. [22]	Cohorts and RCTs	∅	SBP: low DBP: low Hypertension: moderate	Moderate
	Rebholz et al. [19]	RCTs	↓	SBP: moderate DBP: moderate	Low
	∑	<i>n</i> = 4 SRs on animal protein: 3 ∅, 1 ↓ Majority of SRs reach at least a “low” NutriGrade rating Majority of SRs reach at least a “moderate” AMSTAR 2 rating			



Table 2 (continued)

	Systematic review	Included study type	Protein and BP	NutriGrade rating	AMSTAR 2 rating
(C) <i>Plant protein</i>	Mousavi et al. [25]: high vs low analysis	Cohorts	∅	Hypertension: moderate	Moderate
	Chalvon-Demersay et al. [37]	Cohorts	↓	SBP: moderate DBP: moderate	Moderate
	Pedersen et al. [22]	Cohorts	↓	SBP: low DBP: low	Moderate
	Rebholz et al. [19]	RCTs	↓	Hypertension: high SBP: moderate DBP: moderate	Low
	Σ	n = 4 SRs on plant protein: 1 ∅, 3 ↓ Majority of SRs reach at least a “moderate” NutriGrade rating Majority of SRs reach at least a “moderate” AMSTAR 2 rating			
(D) <i>Plant vs animal protein</i>	Systematic review	Included study type	Protein and BP	NutriGrade rating	AMSTAR 2 rating
	Bryant et al. [40]	RCTs	∅	SBP: low DBP: low	Moderate
	Lonnie et al. [26]	RCTs	∅	SBP: low DBP: low	Moderate
	Chalvon-Demersay et al. [37]	RCTs	∅	Moderate	Moderate
	Rebholz et al. [19]	RCTs	∅	SBP: low DBP: low	Low
Σ	n = 4 SRs on plant vs animal protein: 4 ∅ Majority of SRs reach at least a “low” NutriGrade rating Majority of SRs reach at least a “moderate” AMSTAR 2 rating				

AMSTAR 2 A Measurement Tool to Assess Systematic Reviews 2, BP blood pressure, DBP diastolic blood pressure, RCT randomised controlled trial, SBP systolic blood pressure, SR systematic review, ∅ no effect/association or inconsistent results, ↓ blood pressure decrease



**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s00394-024-03336-8>.

**Acknowledgements** We would like to thank all panel members of the guideline on protein intake for their contributions to the methodological approach and specifically to the manuscript on BP. The following scientists deserve special thanks for providing helpful remarks during guideline panel meetings and previous versions of this manuscript: Jürgen M. Bauer, Nicole Kalotai, Tilman Kühn, Andreas Lehmann, Thomas Remer and Dorothee Volkert.

**Author contributions** AMA and AnS conducted the systematic literature search, study selection, data extraction and assessment of methodological quality and outcome-specific certainty of evidence. HB and GIS evaluated the evidence and graded the overall certainty of evidence, which was finalised after discussion with all guideline panel members. HB, GIS, AMA, AnS, AIS and JH prepared the manuscript. All authors read, provided critical feedback of, and approved the manuscript.

**Funding** This research was funded by the German Federal Ministry of Food and Agriculture. The funder had no role in the decisions about data collection, analyses, interpretation of data, in writing the manuscript or in the decision to submit the article for publication.

**Data availability** Not applicable.

## Declarations

**Conflict of interest** A list of any possible conflicts of interest is provided in Supplementary Material S9.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.
















## References

- GBD 2019 Viewpoint Collaborators (2020) Five insights from the Global Burden of Disease Study 2019. *Lancet* 396:1135–1159. [https://doi.org/10.1016/S0140-6736\(20\)31404-5](https://doi.org/10.1016/S0140-6736(20)31404-5)
- Mills KT, Stefanescu A, He J (2020) The global epidemiology of hypertension. *Nat Rev Nephrol* 16:223–237. <https://doi.org/10.1038/s41581-019-0244-2>
- Carey RM, Whelton PK (2018) Prevention, detection, evaluation, and management of high blood pressure in adults: synopsis of the 2017 American College of Cardiology/American Heart Association Hypertension Guideline. *Ann Intern Med* 168:351–358. <https://doi.org/10.7326/M17-3203>
- Shahoud JS, Sanvictores T, Aeddula NR (2023) StatPearls. Physiology, arterial pressure regulation. StatPearls Publishing, St. Petersburg
- Whelton PK, Carey RM, Aronow WS et al (2018) 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension* 71:e13–e115. <https://doi.org/10.1161/HYP.000000000000065>
- Schwingshackl L, Chaimani A, Schwedhelm C et al (2019) Comparative effects of different dietary approaches on blood pressure in hypertensive and pre-hypertensive patients: a systematic review and network meta-analysis. *Crit Rev Food Sci Nutr* 59:2674–2687. <https://doi.org/10.1080/10408398.2018.1463967>
- Aljuraiban GS, Gibson R, Chan DS et al (2023) The role of diet in the prevention of hypertension and management of blood pressure: an umbrella review of meta-analyses of interventional and observational studies. *Adv Nutr*. <https://doi.org/10.1016/j.advnut.2023.09.011>
- Aluko RE (2015) Antihypertensive peptides from food proteins. *Annu Rev Food Sci Technol* 6:235–262. <https://doi.org/10.1146/annurev-food-022814-015520>
- Jauhiainen T, Korpela R (2007) Milk peptides and blood pressure. *J Nutr* 137:825S–S829. <https://doi.org/10.1093/jn/137.3.825S>
- Bhandari D, Rafiq S, Gat Y et al (2020) A review on bioactive peptides: physiological functions, bioavailability and safety. *Int J Pept Res Ther* 26:139–150. <https://doi.org/10.1007/s10989-019-09823-5>
- Shiraseb F, Asbaghi O, Bagheri R et al (2022) Effect of L-arginine supplementation on blood pressure in adults: a systematic review and dose-response meta-analysis of randomized clinical trials. *Adv Nutr* 13:1226–1242. <https://doi.org/10.1093/advances/nmab155>
- Kroke A, Schmidt A, Amini AM et al (2022) Dietary protein intake and health-related outcomes: a methodological protocol for the evidence evaluation and the outline of an evidence to decision framework underlying the evidence-based guideline of the German Nutrition Society. *Eur J Nutr* 61:2091–2101. <https://doi.org/10.1007/s00394-021-02789-5>
- Page MJ, Moher D, Bossuyt PM et al (2021) PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ* 372:n160. <https://doi.org/10.1136/bmj.n160>
- Shea BJ, Reeves BC, Wells G et al (2017) AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ* 358:j4008. <https://doi.org/10.1136/bmj.j4008>
- Schwingshackl L, Knüppel S, Schwedhelm C et al (2016) Perspective: NutriGrade: a scoring system to assess and judge the meta-evidence of randomized controlled trials and cohort studies in nutrition research. *Adv Nutr* 7:994–1004. <https://doi.org/10.3945/an.116.013052>
- Altorf-van der Kuil W, Engberink MF, Brink EJ et al (2010) Dietary protein and blood pressure: a systematic review. *PLoS ONE* 5:e12102. <https://doi.org/10.1371/journal.pone.0012102>
- Dong J-Y, Tong X, Wu Z-W et al (2011) Effect of soya protein on blood pressure: a meta-analysis of randomised controlled trials. *Br J Nutr* 106:317–326. <https://doi.org/10.1017/S0007114511000262>
- Tielemans SMAJ, Altorf-van der Kuil W, Engberink MF et al (2013) Intake of total protein, plant protein and animal protein in relation to blood pressure: a meta-analysis of observational and intervention studies. *J Hum Hypertens* 27:564–571. <https://doi.org/10.1038/jhh.2013.16>
- Rebholz CM, Friedman EE, Powers LJ et al (2012) Dietary protein intake and blood pressure: a meta-analysis of randomized

- controlled trials. *Am J Epidemiol* 176(Suppl 7):S27–S43. <https://doi.org/10.1093/aje/kws245>
20. Santesso N, Akl EA, Bianchi M et al (2012) Effects of higher-versus lower-protein diets on health outcomes: a systematic review and meta-analysis. *Eur J Clin Nutr* 66:780–788. <https://doi.org/10.1038/ejcn.2012.37>
  21. Wycherley TP, Moran LJ, Clifton PM et al (2012) Effects of energy-restricted high-protein, low-fat compared with standard-protein, low-fat diets: a meta-analysis of randomized controlled trials. *Am J Clin Nutr* 96:1281–1298. <https://doi.org/10.3945/ajcn.112.044321>
  22. Pedersen AN, Kondrup J, Børsheim E (2013) Health effects of protein intake in healthy adults: a systematic literature review. *Food Nutr Res* 57:21245. <https://doi.org/10.3402/fnr.v57i0.21245>
  23. Schwingshackl L, Hoffmann G (2013) Long-term effects of low-fat diets either low or high in protein on cardiovascular and metabolic risk factors: a systematic review and meta-analysis. *Nutr J* 12:48. <https://doi.org/10.1186/1475-2891-12-48>
  24. Clifton PM, Condo D, Keogh JB (2014) Long term weight maintenance after advice to consume low carbohydrate, higher protein diets—a systematic review and meta analysis. *Nutr Metab Cardiovasc Dis* 24:224–235. <https://doi.org/10.1016/j.numecd.2013.11.006>
  25. Mousavi SM, Jayedi A, Jalilpiran Y et al (2022) Dietary intake of total, animal and plant proteins and the risk of coronary heart disease and hypertension: a systematic review and dose-response meta-analysis of prospective cohort studies. *Crit Rev Food Sci Nutr* 62:1336–1349. <https://doi.org/10.1080/10408398.2020.1841730>
  26. Lonnie M, Laurie I, Myers M et al (2020) Exploring health-promoting attributes of plant proteins as a functional ingredient for the food sector: a systematic review of human interventional studies. *Nutrients* 12:2291. <https://doi.org/10.3390/nu12082291>
  27. Vogtschmidt YD, Raben A, Faber I et al (2021) Is protein the forgotten ingredient: effects of higher compared to lower protein diets on cardiometabolic risk factors. A systematic review and meta-analysis of randomised controlled trials. *Atherosclerosis* 328:124–135. <https://doi.org/10.1016/j.atherosclerosis.2021.05.011>
  28. Hengeveld LM, de Goede J, Afman LA et al (2022) Health effects of increasing protein intake above the current population reference intake in older adults: a systematic review of the Health Council of the Netherlands. *Adv Nutr* 13:1083–1117. <https://doi.org/10.1093/advances/nmab140>
  29. Stamler J, Liu K, Ruth KJ et al (2002) Eight-year blood pressure change in middle-aged men: relationship to multiple nutrients. *Hypertension* 39:1000–1006
  30. Alonso A, Beunza JJ, Bes-Rastrollo M et al (2006) Vegetable protein and fiber from cereal are inversely associated with the risk of hypertension in a Spanish cohort. *Arch Med Res* 37:778–786. <https://doi.org/10.1016/j.arcmed.2006.01.007>
  31. Liu L, Ikeda K, Sullivan DH et al (2002) Epidemiological evidence of the association between dietary protein intake and blood pressure: a meta-analysis of published data. *Hypertens Res* 25:689–695
  32. Liu K, Ruth KJ, Flack JM et al (1996) Blood pressure in young blacks and whites: relevance of obesity and lifestyle factors in determining differences. The CARDIA Study. *Coronary Artery Risk Development in Young Adults*. *Circulation* 93:60–66. <https://doi.org/10.1161/01.cir.93.1.60>
  33. Simons-Morton DG, Hunsberger SA, van Horn L et al (1997) Nutrient intake and blood pressure in the dietary intervention study in children. *Hypertension* 29:930–936. <https://doi.org/10.1161/01.hyp.29.4.930>
  34. Teunissen-Beekman KFM, Dopheide J, Geleijnse JM et al (2015) Dietary proteins improve endothelial function under fasting conditions but not in the postprandial state, with no effects on markers of low-grade inflammation. *Br J Nutr* 114:1819–1828. <https://doi.org/10.1017/S0007114515003530>
  35. Hidayat K, Du H-Z, Yang J et al (2017) Effects of milk proteins on blood pressure: a meta-analysis of randomized control trials. *Hypertens Res* 40:264–270. <https://doi.org/10.1038/hr.2016.135>
  36. Badely M, Sepandi M, Samadi M et al (2019) The effect of whey protein on the components of metabolic syndrome in overweight and obese individuals; a systematic review and meta-analysis. *Diabetes Metab Syndr* 13:3121–3131. <https://doi.org/10.1016/j.dsx.2019.11.001>
  37. Chalvon-Demersay T, Azzout-Marniche D, Arfsten J et al (2017) A systematic review of the effects of plant compared with animal protein sources on features of metabolic syndrome. *J Nutr* 147:281–292. <https://doi.org/10.3945/jn.116.239574>
  38. Mohammadifard N, Sajjadi F, Haghighatdoost F (2021) Effects of soy consumption on metabolic parameters in patients with metabolic syndrome: a systematic review and meta-analysis. *EXCLI J* 20:665–685. <https://doi.org/10.17179/excli2021-3348>
  39. Mosallanezhad Z, Mahmoodi M, Ranjbar S et al (2021) Soy intake is associated with lowering blood pressure in adults: a systematic review and meta-analysis of randomized double-blind placebo-controlled trials. *Complement Ther Med* 59:102692. <https://doi.org/10.1016/j.ctim.2021.102692>
  40. Bryant L, Rangan A, Grafenauer S (2022) Lupins and health outcomes: a systematic literature review. *Nutrients* 14:327. <https://doi.org/10.3390/nu14020327>
  41. Semlitsch T, Jeitler K, Berghold A et al (2016) Long-term effects of weight-reducing diets in people with hypertension. *Cochrane Database Syst Rev* 3:CD008274. <https://doi.org/10.1002/14651858.CD008274.pub3>
  42. Sabaka P, Dukat A, Gajdosik J et al (2017) The effects of body weight loss and gain on arterial hypertension control: an observational prospective study. *Eur J Med Res* 22:43. <https://doi.org/10.1186/s40001-017-0286-5>
  43. Neter JE, Stam BE, Kok FJ et al (2003) Influence of weight reduction on blood pressure: a meta-analysis of randomized controlled trials. *Hypertension* 42:878–884. <https://doi.org/10.1161/01.HYP.0000094221.86888.AE>
  44. Prentice RL (2020) Dietary assessment and opportunities to enhance nutritional epidemiology evidence. *Ann Intern Med* 172:354–355. <https://doi.org/10.7326/M19-3290>
  45. Bingham SA (2003) Urine nitrogen as a biomarker for the validation of dietary protein intake. *J Nutr* 133(Suppl 3):921S-924S. <https://doi.org/10.1093/jn/133.3.921S>
  46. Boeing H, Bechthold A, Bub A et al (2012) Critical review: vegetables and fruit in the prevention of chronic diseases. *Eur J Nutr* 51:637–663. <https://doi.org/10.1007/s00394-012-0380-y>
  47. Vajdi M, Musazadeh V, Zareei M et al (2023) The effects of whey protein on blood pressure: a systematic review and dose-response meta-analysis of randomized controlled trials. *Nutr Metab Cardiovasc Dis* 33:1633–1646. <https://doi.org/10.1016/j.numecd.2023.05.025>
  48. Zhou S, Xu T, Zhang X et al (2022) Effect of casein hydrolysate on cardiovascular risk factors: a systematic review and meta-analysis of randomized controlled trials. *Nutrients*. <https://doi.org/10.3390/nu14194207>
  49. Liao W, Sun G, Xu D et al (2021) The blood-pressure-lowering effect of food-protein-derived peptides: a meta-analysis of recent

- clinical trials. *Foods* 10:2316. <https://doi.org/10.3390/foods10102316>
50. Mohanty DP, Mohapatra S, Misra S et al (2016) Milk derived bioactive peptides and their impact on human health—a review. *Saudi J Biol Sci* 23:577–583. <https://doi.org/10.1016/j.sjbs.2015.06.005>
51. Perpetuo EA, Juliano L, Lebrun I (2003) Biochemical and pharmacological aspects of two bradykinin-potentiating peptides obtained from tryptic hydrolysis of casein. *J Protein Chem* 22:601–606. <https://doi.org/10.1023/b:jopc.0000008724.98339.ff>
52. Ibrahim HR, Ahmed AS, Miyata T (2017) Novel angiotensin-converting enzyme inhibitory peptides from caseins and whey proteins of goat milk. *J Adv Res* 8:63–71. <https://doi.org/10.1016/j.jare.2016.12.002>
53. Schwingshackl L, Schünemann HJ, Meerpohl JJ (2020) Improving the trustworthiness of findings from nutrition evidence syntheses: assessing risk of bias and rating the certainty of evidence. *Eur J Nutr*. <https://doi.org/10.1007/s00394-020-02464-1>
54. Lamberg-Allardt C, Bärebring L, Arnesen EK et al (2023) Animal versus plant-based protein and risk of cardiovascular disease and type 2 diabetes: a systematic review of randomized controlled trials and prospective cohort studies. *Food Nutr Res*. <https://doi.org/10.29219/fnr.v67.9003>

## Authors and Affiliations

Heiner Boeing<sup>1</sup>  · Anna M. Amini<sup>2</sup>  · Julia Haardt<sup>2</sup>  · Annemarie Schmidt<sup>2</sup>  · Heike A. Bischoff-Ferrari<sup>3</sup> · Anette E. Buyken<sup>4</sup> · Sarah Egert<sup>5</sup>  · Sabine Ellinger<sup>5</sup>  · Anja Kroke<sup>6</sup>  · Stefan Lorkowski<sup>7,8</sup>  · Sandrine Louis<sup>9</sup> · Katharina Nimptsch<sup>10</sup> · Matthias B. Schulze<sup>11,12</sup>  · Alexandra Schutkowski<sup>13</sup>  · Lukas Schwingshackl<sup>14</sup>  · Roswitha Siener<sup>15</sup>  · Armin Zittermann<sup>16</sup>  · Bernhard Watzl<sup>9</sup>  · Gabriele I. Stangl<sup>13</sup>  on behalf of the German Nutrition Society

✉ Anna M. Amini  
corresponding\_author@dge.de

<sup>1</sup> Department of Epidemiology (closed), German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany

<sup>2</sup> German Nutrition Society, Godesberger Allee 136, 53175 Bonn, Germany

<sup>3</sup> Department of Aging Medicine and Aging Research, University Hospital Zurich, University of Zurich, and City Hospital Zurich, Zurich, Switzerland

<sup>4</sup> Institute of Nutrition, Consumption and Health, Faculty of Natural Sciences, Paderborn University, Paderborn, Germany

<sup>5</sup> Institute of Nutritional and Food Science, University of Bonn, Bonn, Germany

<sup>6</sup> Department of Nutritional, Food and Consumer Sciences, Fulda University of Applied Sciences, Fulda, Germany

<sup>7</sup> Institute of Nutritional Sciences, Friedrich Schiller University Jena, Jena, Germany

<sup>8</sup> Competence Cluster for Nutrition and Cardiovascular Health (nutriCARD), Halle-Jena-Leipzig, Germany

<sup>9</sup> Department of Physiology and Biochemistry of Nutrition, Max Rubner-Institut, Karlsruhe, Germany

<sup>10</sup> Molecular Epidemiology Research Group, Max Delbrück Center for Molecular Medicine (MDC) in the Helmholtz Association, Berlin, Germany

<sup>11</sup> Department of Molecular Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany

<sup>12</sup> Institute of Nutritional Science, University of Potsdam, Potsdam, Germany

<sup>13</sup> Institute of Agricultural and Nutritional Sciences, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany

<sup>14</sup> Institute for Evidence in Medicine, Medical Center, Faculty of Medicine, University of Freiburg, Freiburg, Germany

<sup>15</sup> Department of Urology, University Stone Center, University Hospital Bonn, Bonn, Germany

<sup>16</sup> Clinic for Thoracic and Cardiovascular Surgery, Herz- und Diabeteszentrum Nordrhein Westfalen, Ruhr University Bochum, Bad Oeynhausen, Germany