

# Bace1 and Neuregulin-1 cooperate to control formation and maintenance of muscle spindles

Cyril Cheret<sup>1</sup>, Michael Willem<sup>2</sup>,  
Florence R Fricker<sup>3</sup>, Hagen Wende<sup>1</sup>,  
Annika Wulf-Goldenberg<sup>4</sup>,  
Sabina Tahirovic<sup>5</sup>, Klaus-Armin Nave<sup>6</sup>,  
Paul Saftig<sup>7</sup>, Christian Haass<sup>2,5,8</sup>,  
Alistair N Garratt<sup>9</sup>, David L Bennett<sup>3</sup>  
and Carmen Birchmeier<sup>1,\*</sup>

<sup>1</sup>Entwicklungsbiologie/Signaltransduktion, Max-Delbrück-Center for Molecular Medicine, Berlin, Germany, <sup>2</sup>Adolf-Butenandt-Institute - Biochemistry, Ludwig-Maximilians-University Munich, Munich, Germany, <sup>3</sup>University of Oxford, NDCN, Oxford, UK, <sup>4</sup>Experimental Pharmacology & Oncology Berlin-Buch GmbH, Berlin, Germany, <sup>5</sup>German Center for Neurodegenerative Diseases (DZNE) Munich, Munich, Germany, <sup>6</sup>Max Planck Institute of Experimental Medicine, Göttingen, Germany, <sup>7</sup>Biochemical Institute, Christian-Albrechts University Kiel, Kiel, Germany, <sup>8</sup>Munich Cluster for Systems Neurology (SyNergy), Berlin, Germany and <sup>9</sup>Center for Anatomy, Institute of Cell Biology and Neurobiology, Charité - Universitätsmedizin Berlin, Berlin, Germany

**The protease  $\beta$ -secretase 1 (Bace1) was identified through its critical role in production of amyloid- $\beta$  peptides (A $\beta$ ), the major component of amyloid plaques in Alzheimer's disease. Bace1 is considered a promising target for the treatment of this pathology, but processes additional substrates, among them Neuregulin-1 (Nrg1). Our biochemical analysis indicates that Bace1 processes the Ig-containing  $\beta$ 1 Nrg1 (IgNrg1 $\beta$ 1) isoform. We find that a graded reduction in IgNrg1 signal strength *in vivo* results in increasingly severe deficits in formation and maturation of muscle spindles, a proprioceptive organ critical for muscle coordination. Further, we show that Bace1 is required for formation and maturation of the muscle spindle. Finally, pharmacological inhibition and conditional mutagenesis in adult animals demonstrate that Bace1 and Nrg1 are essential to sustain muscle spindles and to maintain motor coordination. Our results assign to Bace1 a role in the control of coordinated movement through its regulation of muscle spindle physiology, and implicate IgNrg1-dependent processing as a molecular mechanism.**

*The EMBO Journal* (2013) 32, 2015–2028. doi:10.1038/emboj.2013.146; Published online 21 June 2013

**Subject Categories:** development; neuroscience

**Keywords:** Bace1; muscle spindle; Nrg1; proprioception

\*Corresponding author. Entwicklungsbiologie/Signaltransduktion, Max-Delbrück-Center for Molecular Medicine, Robert Roessle Strasse 10, Berlin 13125, Germany. Tel.: +49 30 9406 2403; Fax: +49 30 9406 3765; E-mail: cbirch@mdc-berlin.de

Received: 5 February 2013; accepted: 29 May 2013; published online: 21 June 2013

## Introduction

Proteolysis of membrane-tethered molecules is critical for cellular communication. Sheddases, a group of membrane-bound proteases, cleave single-span membrane proteins at the extracellular surface. One of the most studied sheddases, the aspartyl protease Bace1 ( $\beta$ -secretase 1), is an important drug target for Alzheimer's disease. Bace1 cleaves the amyloid precursor protein (APP) and is responsible for generation of pathogenic A $\beta$  peptides (Luo *et al.*, 2001; Vassar *et al.*, 2009). In addition to APP, around 20 substrates have been identified for Bace1 (Willem *et al.*, 2006; Kandalepas and Vassar, 2012; Kuhn *et al.*, 2012; Zhou *et al.*, 2012). Knowledge of their physiological functions can help to monitor adverse effects caused by Bace1 inhibition in patients.

One Bace1 substrate is Neuregulin-1 (Nrg1), a trophic factor that signals through ErbB tyrosine kinase receptors to regulate nervous system development and regeneration (Falls, 2003; Willem *et al.*, 2006; Birchmeier, 2009; Fricker *et al.*, 2011). Nrg1 is a complex gene encoding more than 15 protein isoforms that are generated by a combination of alternative mRNA splicing and the use of several promoters. All subtypes of Nrg1 proteins present an EGF-like domain required for receptor binding and signalling. Depending on the presence of either an Ig-like or a cysteine-rich domain (CRD) in their amino-terminal (N-terminal) sequences, Nrg1 variants can be classified into IgNrg1 (type I and type II) and CRD-Nrg1 (type III). Few Nrg1 variants are secreted molecules (e.g., glial growth factor, GGF), most being membrane-bound proteins that can be shed by proteases (Falls, 2003). Bace1-dependent shedding of Nrg1 has been implicated in the control of myelination in the peripheral nervous system (Hu *et al.*, 2006; Willem *et al.*, 2006).

Coordinated body movement requires constant input of sensory information to elicit a concerted motor response. Muscle spindles are sensory organs dispersed throughout muscles of vertebrates, which detect muscle stretch, allowing thus the perception of body position (proprioception) important for coordinated movement (Maier, 1997). The muscle spindle is composed of a bundle of specialized (intrafusal) muscle fibres, and its formation is induced by contact between TrkC<sup>+</sup> sensory axons and muscle fibres (Ernfors *et al.*, 1994; Farinas *et al.*, 1994; Klein *et al.*, 1994; Tessarollo *et al.*, 1994; Walro and Kucera, 1999; Chen *et al.*, 2003). Further maturation takes place during late fetal and early postnatal phases when muscle spindles grow and become enclosed by the capsules (Hunt, 1990; Zelena and Soukup, 1993; Maier, 1997). Previous studies showed that the genetic deletion of neuronally produced Nrg1 or its receptor ErbB2 in muscle tissue prevent muscle spindle differentiation (Andrechek *et al.*, 2002; Hippenmeyer *et al.*, 2002; Leu *et al.*, 2003).

Here we show that formation, maturation and maintenance of muscle spindles depend on Bace1. We use mouse mutants to demonstrate that in the absence of Bace1, muscle spindle numbers are reduced and spindle maturation is impaired.

Moreover, we find that a graded reduction in IgNrg1 signal strength results in increasingly severe deficits in the formation and maturation of muscle spindles. Conversely, we observed supernumerary muscle spindles upon overexpression of a membrane-tethered Nrg1 variant (Ig-containing  $\beta 1$  Nrg1 (IgNrg1 $\beta 1$ )) in sensory neurons, an effect which strictly depended on the presence of Bace1. Strikingly, inhibition of Bace1 activity or ablation of Nrg1 expression in adult mice resulted in a massive reduction of the muscle spindle pool and impaired coordination of movement. Together, these data implicate Bace1-dependent processing of IgNrg1 in ontogenesis and long-term maintenance of muscle spindles.

## Results

### **Bace1 mutant mice display coordination defects**

During handling of Bace1<sup>-/-</sup> mutant mice, we noted that their movement was altered. Most notably, they appeared not to be able to hang to an inverted grid, a task requiring motor coordination and/or muscle strength (Coughenour *et al*, 1977; Landauer *et al*, 2003). A quantitative assessment demonstrated that wild-type littermates did hang to the grid three times longer than Bace1<sup>-/-</sup> mice; Bace1<sup>+/-</sup> and wild-type animals performed similarly (Figure 1A). We then assessed muscle coordination using gait analysis. The walking pattern was recorded as mice freely walked on a moving belt, and gait analysis monitored movement and position of individual paws using the Treadscan system (Beare *et al*, 2009). The walking pattern of wild-type mice (Figure 1B and C) shows coordinated alternation of paws, i.e., opposing movements of anterior and posterior limbs on the same side (homolateral coupling value  $\approx 0.5$ ; Figure 1D and Supplementary Figure S1A), and opposing movements of left and right limbs of the same axial levels (homologue coupling value  $\approx 0.5$ ; Figure 1E and Supplementary Figure S1B and C). Homolateral coupling of Bace1<sup>-/-</sup> mutants was severely affected and deviated considerably from wild-type animals. This reflects a lack of forelimb/hindlimb coordination and resulted in a swaying walking pattern (Figure 1C; quantified in D and Supplementary Figure S1A). In contrast, homologue coupling was little disturbed at either axial level, indicating correct left/right alternation (Figure 1C; quantified in E and Supplementary Figures S1B and C). Bace1 mutant mice display peripheral hypomyelination but little Schwann cell turnover, similar to Schwann cell-specific coErbB2 (Krox20<sup>cre</sup> ErbB2<sup>fllox/fllox</sup>) mutant mice (*g*-ratios P180: control, Bace1<sup>-/-</sup> and coErbB2:  $0.68 \pm 0.01$ ,  $0.75 \pm 0.01$  and  $0.80 \pm 0.01$ , respectively; *cf.* Garratt *et al*, 2000; Hu *et al*, 2006; Willem *et al*, 2006; Grossmann *et al*, 2009). In contrast to Bace1 mutants, we did not observe significant changes in motor coordination in coErbB2 mutants (Supplementary Figure S2A–D). Together, our data indicate that motor coordination is disrupted in Bace1<sup>-/-</sup> mutants, and this coordination deficit is not caused by hypomyelination.

### **Bace1 mutation affects formation and maintenance of muscle spindles**


Coordination of body movement requires functional proprioception, and muscle spindles are important proprioceptive organs governing the coupling of antagonistic muscles (the spindle structure is shown schematically in Figure 1F).

We quantified the amounts of muscle spindles in lower hindlimbs of newborn control and mutant mice at P0, using a combination of morphological criteria (large nuclei, presence of a capsule) and immunohistochemistry with antibodies against Egr3, a muscle spindle-specific transcription factor, collagen IV, a marker for muscle spindle outer capsules, and NF200, a neurofilament isoform expressed by sensory fibres contacting the spindles (Figure 1G and H; *cf.* Tourtellotte and Milbrandt, 1998; Tourtellotte *et al*, 2001; Hippenmeyer *et al*, 2002). Newborn Bace1<sup>-/-</sup> mice presented a pronounced reduction (45%;  $P < 0.001$ ) of the number of muscle spindles, and the reduction persisted until adulthood (Figure 1G, Table I). A small but significant reduction (14%,  $P = 0.01$ ) was observable in heterozygous Bace1<sup>+/-</sup> mutant mice. The overall morphology of persisting muscle spindles was unchanged in Bace1<sup>-/-</sup> mice at P0, but postnatal muscle spindle growth was impaired (Figure 1H, Table II, see also below). We conclude that Bace1 is required for the correct formation and maturation of muscle spindles.

### **Bace1 activity is required to sustain muscle spindles and to maintain motor coordination**

We next assessed whether Bace1 activity is needed to sustain mature muscle spindles. Adult (P180) wild type, heterozygous and homozygous Bace1 mutant mice were treated with the pharmacologic Bace1 inhibitor Ly2811376 for a period of 29 days (May *et al*, 2011). Ly2811376 inhibited Bace1 activity effectively *in vivo*, as assessed by monitoring the Nrg1 processing in the brain (Figure 2A). Ly2811376 treatment led to a regression of adult muscle spindles, notably a loss of 40% of muscle spindles in wild type and heterozygous Bace1 mutant animals, compared to corresponding vehicle-treated groups (Figure 2B; Supplementary Figure S2E). Ly2811376 treatment did not further decrease the muscle spindle pool in homozygous Bace1 mutant mice, demonstrating that this effect was mediated through specific Bace1 inhibition (Figure 2B; Supplementary Figure S2E). Together, these genetic and pharmacological data indicate that Bace1 controls the maintenance of muscle spindles during adulthood, as well as their formation during development.

We next tested whether Bace1 inhibition in the adult affected motor coordination. A quantitative assessment of grip ability demonstrated that animals treated with a Bace1 inhibitor lost their footing 3–4 times faster than animals treated with the vehicle (Figure 2C). Gait analysis also demonstrated that the walking pattern of the mice was aberrant after long-term inhibition with Bace1 inhibitor (Figure 2D–F). Thus, the value of homolateral coupling deviated considerably from the one observed in vehicle-treated animals. This reflects a lack of forelimb/hindlimb coordination and resulted in a swaying walking pattern (Figure 2D; quantified in E and Supplementary Figure S1A). Homologue coupling was little disturbed after Bace1 inhibition (Figure 2D; quantified in F and Supplementary Figure S1B and C). We conclude that long-term treatment of adult mice with Bace1 inhibitor disrupts motor coordination. It is noteworthy that coordination was affected to similar extents in Bace1<sup>-/-</sup> and in Bace1 inhibitor-treated animals, indicating that Bace1 activity is continuously required for motor coordination.



**Figure 1** Bace1 is required for the correct formation of muscle spindles. (A) Performance of Bace1<sup>+/+</sup>, Bace1<sup>+/-</sup> and Bace1<sup>-/-</sup> mice (P180) in the inverted grid test. Horizontal bars show the average time the mice remain clinging to the grid. (B) Normal walking pattern of a mouse; the movement of homolateral (top) and homologous (bottom) limbs are indicated. (C) Representative walking patterns of control and Bace1<sup>-/-</sup> mice; displayed are series of four consecutive steps. (D, E) Homolateral (D) and homologue (E) coupling values for the movements of control (inner circle) and Bace1<sup>-/-</sup> mice (outer circle). A value of 0.5 defines coordinated movement of the paws. The grey area indicates the non-pathological interval (0.5 ± 0.1). (F) Schematic representation of a muscle spindle; indicated are the central intrafusal fibres of the spindle, which are surrounded by a capsule and (extrafusal) muscle fibres. (G) Quantification of muscle spindles in hindlimbs of control, heterozygous and homozygous Bace1 mutant mice at P0. (H) Immunohistological analysis of muscle spindles from control and Bace1 mutant mice at P0. Intrafusal fibres express Egr3, the nascent outer capsule displays a pronounced collagen IV staining, and contacting sensory fibres are NF200<sup>+</sup>. Scale bar, 10 μm (H).

**Table I** Number of muscle spindles in lower hindlimbs of newborn and adult mice

	Control	IgNrg1 <sup>Δ/+</sup>	Bace1 <sup>-/-</sup>	Bace1 <sup>-/-</sup> IgNrg1 <sup>Δ/+</sup>	co-IgNrg1	IgNrg1β1 <sup>0v</sup>	Bace1 <sup>-/-</sup> IgNrg1β1 <sup>0v</sup>
P0	8.8 ± 0.3	6.8 ± 0.3**	4.8 ± 0.6***	2.6 ± 0.4***	1.0 ± 0.2***	14.1 ± 1.2***	5.0 ± 0.2***
P30	10.0 ± 0.6	8.1 ± 0.5*	5.6 ± 0.5***	3.2 ± 0.2***	1.8 ± 0.3***	18.8 ± 0.5***	5.9 ± 0.2***

Values are mean ± s.e.m. of 5–6 animals per genotype and age, expressed as number of muscle spindles per hindlimb section. Significance of the differences between numbers observed in control and mutants is indicated.

### Bace1 processes Nrg1 isoforms

Various Nrg1 isoforms exist (Figure 3A) that can take over distinct functions. Nrg1 isoforms containing an Ig domain

(IgNrg1) are produced by proprioceptive neurons and control the induction of the muscle spindle (Hippenmeyer *et al*, 2002). Bace1 is expressed broadly in sensory neurons

**Table II** Diameter of muscle spindles and intrafusal content at birth and during adulthood

	Control	Bace1 <sup>-/-</sup>	Bace1 <sup>-/-</sup> IgNrg1 <sup>Δ/+</sup>	co-IgNrg1	IgNrg1β1 <sup>OV</sup>	Bace1 <sup>-/-</sup> IgNrg1β1 <sup>OV</sup>
<i>Spindle diameter (μm)</i>						
P0	26.5 ± 0.9	23.7 ± 1.4 <sup>NS</sup>	23.4 ± 0.9 <sup>NS</sup>	25.9 ± 1.1 <sup>NS</sup>	26.8 ± 1.5 <sup>NS</sup>	22.7 ± 0.5 <sup>NS</sup>
P30	40.6 ± 0.6	32.5 ± 0.8 <sup>***</sup>	33.3 ± 0.6 <sup>***</sup>	29.4 ± 1.0 <sup>***</sup>	42.1 ± 0.6 <sup>NS</sup>	34.6 ± 1.2 <sup>**</sup>
<i>Intrafusal fibres/spindle</i>						
P0	4.0 ± 0.2	3.9 ± 0.2 <sup>NS</sup>	3.1 ± 0.1 <sup>**</sup>	2.9 ± 0.1 <sup>***</sup>	4.3 ± 0.2 <sup>NS</sup>	4.0 ± 0.1 <sup>NS</sup>
P30	3.9 ± 0.1	3.9 ± 0.1 <sup>NS</sup>	3.4 ± 0.1 <sup>NS</sup>	3.1 ± 0.1 <sup>**</sup>	4.5 ± 0.2 <sup>***</sup>	3.8 ± 0.1 <sup>NS</sup>

Equatorial diameter and number of intrafusal fibres per muscle spindle was determined in 3–6 animals per genotype and per age and is expressed as mean ± s.e.m. Asterisks indicate significance of the differences observed when mutants were compared to control littermates.

(Willem *et al*, 2006). *In situ* hybridization combined with immunohistochemistry showed that Bace1 and IgNrg1 are co-expressed in NF200<sup>+</sup> large diameter sensory neurons at birth; thus proprioceptive neurons co-express Bace1 and IgNrg1 (Figure 3B). Quantification demonstrated that the vast majority of sensory neurons as well as IgNrg1<sup>+</sup> sensory neurons in dorsal root ganglia (DRG) co-expressed Bace1 (93.7 ± 1.2% and 99.8 ± 0.2%, respectively). Various IgNrg1 isoforms exist, called α1/2 and β1–4 that differ in the EGF-like domains and in sequences carboxy-terminal (C-terminal) thereof (Supplementary Figure S3A; Falls, 2003). We analysed their expression in DRG at P0 using semi-quantitative PCR (qPCR). This showed that among the IgNrg1 isoforms, β variants and particularly, the β1 isotype are expressed at highest levels (Figure 3C).

IgNrg1β1 isoforms contain a predicted Bace1 cleavage site (Figure 3D and Hu *et al*, 2008). However, IgNrg1 isoforms are produced by few sensory neurons, whereas CRD-Nrg1 is expressed broadly in sensory neurons (Figure 3B, *cf.* Meyer *et al*, 1997; Hippenmeyer *et al*, 2002). This hampers a direct biochemical analysis of IgNrg1 processing in sensory ganglia *in vivo*. We therefore analysed processing of HA-tagged β1 isoforms in cultured HEK293 cells, and tested whether the predicted Bace1 cleavage sites present in β1 variants of Ig- or CRD-containing Nrg1 are recognized. Full-length and processed C-terminal fragment of Nrg1β1 proteins containing CRD or Ig domains were observed in the absence of transfected Bace1 cDNA, using an antibody against the C-terminus of Nrg1 (Figure 3E). In the presence of Bace1, the full-length protein almost disappeared, whereas the processed C-terminal fragment accumulated (Figure 3E). Thus, Bace1 cleaves the Nrg1β1 sequence, regardless of whether it is present in a CRD or Ig isoform.

IgNrg1β2 and IgNrg1β3 represent a membrane-tethered and a secreted isoform, respectively, and are produced at lower levels than IgNrg1β1 in DRG (Figure 3A and C). We compared processing of the two membrane-tethered isoforms IgNrg1β1 and IgNrg1β2 in transfected HEK293 cells and analysed the release of their HA-tagged extracellular fragment, which contains the receptor-binding EGF domain (Figure 3F). IgNrg1β1 was constitutively processed, but co-transfection of Bace1 resulted in a further increase in the amount of the C-terminal and HA-tagged N-terminal fragments, as well as a decrease of full-length protein, indicating increased processing. In contrast, IgNrg1β2 was neither processed constitutively, nor in a Bace1-dependent manner (Figure 3F). The release of the N-terminal fragments was quantified using Nrg1 constructs carrying alkaline phosphatase in their extracellular domain (Figure 3G; secreted


alkaline phosphatase or SEAP, *cf.* Willem *et al*, 2006). Co-transfection of IgNrg1β1<sup>SEAP</sup> and Bace1 increased the amount of released alkaline phosphatase six-fold, which was abolished by the Bace1 inhibitor C3 but not by the metalloproteinase inhibitor GM6001 (Figure 3G; Willem *et al*, 2006; Freese *et al*, 2009). In contrast, released alkaline phosphatase from the corresponding IgNrg1β2<sup>SEAP</sup> was low and little affected by Bace1 activity. We conclude that the major Ig variant expressed in sensory neurons, Nrg1β1, is a substrate of Bace1.

We next tested whether endogenous levels of Bace1 can process IgNrg1β1. Hippocampal neurons were transfected with an IgNrg1β1 expression vector (Figure 3H and I). The majority of cellular IgNrg1β1 was processed, and only little full-length protein was observable. In the presence of the Bace1 inhibitor C3, IgNrg1β1 cleavage was reduced, resulting in accumulation of Nrg1 FL and slightly reduced levels of the 65-kDa C-terminal fragment (Figure 3H). HA-tagged Nrg1β1 in the supernatant was detected using HA-specific antibodies, which revealed mildly reduced quantities of the N-terminal fragment in case of Bace1 and metalloproteinase inhibition (Figure 3I). The combination of Bace1 and metalloproteinase inhibitors acted synergistically. In addition, the N-terminal fragment was immunoprecipitated using HA-specific antibodies and detected on western blot using HA and 4F10 antibodies. We took advantage of the 4F10 antibody, which recognizes a Nrg1β1-specific epitope only when exposed by Bace1 cleavage (Fleck *et al*, 2013). This revealed that the neuronal production of the Nrg1β1 4F10-specific epitope was impaired upon inhibition of Bace1, but not of metalloproteinases (Figure 3I). These data indicate that endogenous amounts of Bace1 suffice to cleave IgNrg1β1 in primary hippocampal neurons, and support the notion that in such neurons IgNrg1β1 is cleaved by Bace1 and metalloproteinases.

### ***IgNrg1 isoforms are required for motor coordination***

To directly define the function of IgNrg1 isoforms in motor coordination, we generated a new mutant allele in which one of the exons encoding Ig sequences is ‘floxed’ or constitutively deleted (IgNrg1<sup>fllox</sup> and IgNrg1<sup>Δ</sup>; Supplementary Figure S3B). IgNrg1<sup>Δ</sup> or cre-recombined IgNrg1<sup>fllox</sup> mutations specifically interfere with the production of IgNrg1, but not CRD-Nrg1 transcripts (see Materials and methods for more details). We then generated Wnt1<sup>cre/+</sup> IgNrg1<sup>fllox/Δ</sup> mice; Wnt1<sup>cre</sup> is expressed in neural crest cells that give rise to sensory neurons (Danielian *et al*, 1998; Le Douarin and Kalcheim, 1999). Semi-quantitative RT-PCR indicated that expression of IgNrg1 and particularly IgNrg1β1





**Figure 2** Bace1 activity regulates motor coordination and maintains muscle spindles during adulthood. (A) Western blot analysis of Bace1 expression and processing of endogenous Nrg1 in the mid-brain of adult (P180) wild type (wt) or Bace1<sup>-/-</sup> (B1<sup>-/-</sup>) mice treated with vehicle (Vh) or the Bace1 inhibitor Ly2811376 (Ly). Nrg1 FL denotes the 130 kDa Nrg1 species corresponding to full-length CRD-Nrg1, detected by an antibody directed against the Nrg1 C-terminal sequence. The amount of Nrg1 FL is markedly increased upon Bace1 inhibition or ablation, indicating reduced cleavage, but Bace1 expression is unaffected by Ly2811376 treatment. Calnexin is used as internal control. (B) Quantification of muscle spindles in the tibialis anterior muscle of control, heterozygous and homozygous Bace1 mutant mice (P180) treated with the Bace1 inhibitor Ly2811376. (C) Performance of Bace1<sup>+/+</sup>, Bace1<sup>-/-</sup> and vehicle- or Ly2811376-treated wild-type mice in the inverted grid test. Note that Bace1<sup>-/-</sup> and Ly2811376-treated mice perform similarly. (D) Representative walking patterns of vehicle- and Ly2811376-treated wild-type mice. (E, F) Homolateral (E) and homologue (F) coupling values for the movements of vehicle- (inner circle) and Ly2811376-treated (outer circle) wild-type mice. Bace1<sup>-/-</sup> mice (middle circle) are included for comparison.

transcripts were reduced by 90 and 50% in DRG of Wnt1<sup>cre/+</sup> IgNrg1<sup>flox/Δ</sup> (hereafter called co-IgNrg1 mutants) and heterozygous IgNrg1<sup>Δ/+</sup> newborn mice, respectively (Figure 4A; Supplementary Figure S4A). In contrast,

expression of CRD-Nrg1 transcripts was unaltered in DRG of co-IgNrg1 mice (Figure 4A).


Conditional IgNrg1 mutants displayed pronounced curled tails and uncoordinated tail movements, which we attributed to an almost complete depletion of the spindle pool in tail muscles (Supplementary Figure S4B). Besides, co-IgNrg1 mice lost their grip five times faster than their control littermates in the inverted grid test (Figure 4B). Moreover, gait analysis showed that co-IgNrg1 mutants, like the Bace1 mice described above, suffered from pronounced deficits in forelimb/hindlimb coordination (Figure 4C and D). In addition, a pronounced deficit in left/right coordination was apparent in hindlimbs, resulting in frequent hopping movement (Figure 4E, summarized in Figure 7I). We conclude that, similar to Bace1 mutants, coordinated movement is severely disrupted in co-IgNrg1 mice.

We next analysed muscle spindles in IgNrg1 mutants (Figure 5A; Supplementary Figure S5A and Supplementary Table II). We observed a significant (30%,  $P=0.002$ ) reduction and a massive loss of spindles (84%,  $P<0.001$ ) in heterozygous IgNrg1<sup>Δ/+</sup> and co-IgNrg1 newborn mice, respectively. No obvious alteration was apparent in the morphology of Golgi tendon organs and Pacinian corpuscles in co-IgNrg1 mutants, nor did we detect changes in myelination of peripheral nerves (Supplementary Figure S5B and C,  $g$ -ratios P12:  $0.72 \pm 0.02$  and  $0.71 \pm 0.01$  in the sciatic nerve of control and co-IgNrg1 mice, respectively). When the heterozygous IgNrg1 mutation was introduced on a Bace1<sup>-/-</sup> background, a more pronounced loss of muscle spindles was observed, and only 25% of the normal numbers were present in Bace1<sup>-/-</sup> IgNrg1<sup>Δ/+</sup> mice (Figure 5A; Supplementary Figure S5A). This indicates that a graded reduction in IgNrg1 signal strength results in increasingly severe deficits in muscle spindle formation.


We further assessed the formation of muscle spindles in a transgenic mouse strain in which IgNrg1β1 is overexpressed in neurons under the control of the Thy1 promoter (type I IgNrg1β1, hereafter called IgNrg1β1<sup>ov</sup>; cf. Michailov *et al*, 2004 and Figure 5A; Supplementary Figure S5A). Such mice displayed a 50% increase in muscle spindle numbers at birth, and these supernumerary spindles persisted until adulthood (Figure 5A, Table I). The expansion of the muscle spindle pool was accompanied by increased survival of proprioceptive neurons during the selection phase (Figure 5B–D), which is possibly caused by expression of neurotrophin 3 (NT3) in the supernumerary muscle spindles. Strikingly, only in the presence of Bace1 supernumerary spindles were observed in IgNrg1β1<sup>ov</sup> transgenic mice. In particular, the muscle spindle pool of Bace1<sup>-/-</sup> IgNrg1β1<sup>ov</sup> was smaller than the one of control mice, and reached similar numbers to those observed in Bace1<sup>-/-</sup> mutants (Figure 5A). Increased survival of proprioceptive neurons during the selection phase was not observable in compound mutant Bace1<sup>-/-</sup> IgNrg1β1<sup>ov</sup> animals (Figure 5C and D). We conclude that transgenic IgNrg1β1 strictly requires Bace1 processing to exert its role in muscle spindle formation.

### Maturation of muscle spindles depends on Bace1 and Nrg1

Muscle spindles are induced before birth and mature during postnatal life, when they can be detected by a combination of morphological criteria and immunohistology (Maier, 1997;



**Figure 3** Nrg1 $\beta$ 1 is a substrate of Bace1. **(A)** Structure of Nrg1 isoforms containing cysteine-rich domain (CRD) or Ig-like (Ig) domains. Full-length (Nrg1 FL), N-terminal (Nrg1 NtF) and C-terminal (Nrg1 CtF) fragments analysed in **(E, F, H, I)** are indicated. Green arrows show the Bace1 cleavage site present in  $\beta$ 1 isoforms, yellow star the position of the HA-tag used for biochemical analysis; C-terminal transmembrane domain TM. **(B)** *In situ* hybridization (*Bace1*, *IgNrg1*) combined with immunohistology (neurofilament 200; NF200) demonstrates co-expression of *Bace1* and *IgNrg1* in sensory neurons. Left: *Bace1* is broadly expressed in sensory neurons. Right: *IgNrg1* and *Bace1* (insert) are co-expressed in NF200<sup>+</sup> sensory neurons. **(C)** qPCR of DRG mRNA encoding CRD-Nrg1 and  $\alpha/\beta$  IgNrg1 isoforms (P0). **(D)** Sequence alignment of  $\alpha/\beta$  Nrg1 isoforms; predicted Bace1 cleavage sites are indicated. Asterisk indicates stop codon in  $\beta$ 3. **(E, F)** Western blot analysis of Bace1-dependent processing of **(E)** Ig- and CRD-Nrg1 $\beta$ 1, **(F)** IgNrg1 $\beta$ 1 and IgNrg1 $\beta$ 2 in HEK293 cells. Antibodies are indicated ( $\alpha$ CNX: anti-calnexin). Full-length precursors (Nrg1 FL; IgNrg1: 110 kD, CRD-Nrg1: 130 kD) and processed C-terminal fragment (Nrg1 CtF: 65 kD) are detected in RIPA lysates using an antibody against the Nrg1 C-terminus; HA-tagged N-terminal fragments (Nrg1 NtF) are detected in the supernatant using anti-HA. Calnexin serves as loading control. NT, non-transfected. **(G)** Quantification of Bace1-dependent shedding of IgNrg1 $\beta$ 1<sup>SEAP</sup> and IgNrg1 $\beta$ 2<sup>SEAP</sup>. The N-terminal sequence of IgNrg1<sup>SEAP</sup> variants contains alkaline phosphatase whose enzymatic activity is detected in supernatants in the absence/presence of Bace1 cDNA, C3 or GM6001 inhibitors. **(H)** Western blot analysis of IgNrg1 $\beta$ 1 cleavage in primary neurons. Detected are full-length and C-terminal IgNrg1 $\beta$ 1 in RIPA lysates using an antibody recognizing the Nrg1 C-terminus. **(I)** The N-terminal fragment of IgNrg1 $\beta$ 1 was directly detected in supernatant by western blotting using anti-HA (upper panels). Alternately, supernatant was immunoprecipitated using anti-HA (middle and lower panels), and Nrg1 NtF fragments were detected using anti-HA and 4F10 antibodies; the latter identifies a Bace1-specific Nrg1 cleavage product. Asterisks indicate cross-reactive proteins. Scale bar, 50  $\mu$ m **(B)**.



**Figure 4** IgNrg1 isoforms control motor coordination. (A) Quantification of the expression of CRD and Ig Nrg1 isoforms in DRG neurons from control, IgNrg1<sup>Δ/Δ</sup>, co-IgNrg1 and IgNrg1β1<sup>Ov</sup> newborn mice using qPCR. (B) Performance of control and co-IgNrg1 mice (P180) in the inverted grid test. (C) Representative walking pattern of control and co-IgNrg1 mice. (D, E) Homolateral (D) and homologue (E) coupling values of limbs in control (inner circle) and co-IgNrg1 (outer circle) mice (P180).

Chen *et al*, 2003). Generally, similar changes in numbers of muscle spindles were observed in adult (P30) IgNrg1 and Bace1 mutant mice as those present at birth (Table I). However, we noted that muscle spindles from different mutant strains presented maturation deficits. Calbindin protein is detected only in mature intrafusal fibres (Chen *et al*, 2002; Gould *et al*, 2008). Calbindin staining was reduced in co-IgNrg1 mutants and, conversely, increased in IgNrg1β1<sup>Ov</sup> mice (Figure 6A, quantified in Supplementary Figure S6A). Similar changes were observed with S100 staining in intrafusal fibres of control, co-IgNrg1 and IgNrg1β1<sup>Ov</sup> mice (Supplementary Figure S6B). We estimated the diameter and determined the number of intrafusal fibres per muscle spindle and observed that the co-IgNrg1 or Bace1 mutations resulted in increased proportions of muscle

spindles characterized by a small diameter (<30 μm; Figure 6B) and few intrafusal fibres (≤3 fibres/spindle; Figure 6D). Conversely, neuronal overexpression of IgNrg1β1 resulted in higher proportions of muscle spindles with unusually large diameter (>50 μm; Figure 6C) and high intrafusal fibre numbers (>4 fibres/spindle; Figure 6E). The outer capsule morphology and the innervation of remaining muscle spindles appeared unchanged in all genotypes (Figure 6A). Thus, Bace1 as well as IgNrg1 govern the maturation of muscle spindles.


### Maintenance of muscle spindles requires continuous Nrg1 signalling

Bace1 inhibition in the adult results in motor coordination deficits and in a depletion of the muscle spindle pool (Figure 2B–F). We next tested whether Nrg1 is also continuously needed for motor coordination and muscle spindle maintenance. We used for this a previously established colony of cre-ER<sup>TM/+</sup> Nrg1<sup>flox/flox</sup> mice (Fricker *et al*, 2013) hereafter called coTxNrg1 mice. In such animals, a ubiquitously expressed tamoxifen-inducible cre induces recombination, and recombination deletes sequences encoding the EGF domain present in all Nrg1 isoforms. The mutation was introduced at P70, after somatic growth ended, and eliminated expression of Nrg1 transcripts in DRG of coTxNrg1 animals at P100 (see Figure 7A for an outline of the experiment and Figure 7B for quantification of Nrg1 transcripts). coTxNrg1 mice displayed profoundly impaired grip and did hang 3–4 times less long in the inverted grid test than control littermates. Motor coordination was strongly impaired, as assessed using a beam-walking test (Figure 7C and D; Supplementary Figure S7). We also analysed the muscle spindles in coTxNrg1 mutants: coTxNrg1 mice possessed less than half as many muscle spindles as control littermates (Figure 7E and F), and remaining spindles were shortened (Figure 7G; spindle length: 1805 ± 106 and 1593 ± 56 μm in control and coTxNrg1 mice, respectively; *P* = 0.048). In addition, the morphology of residual muscle spindles was profoundly altered and displayed thinner and discontinuous collagen IV staining (Figure 7H), which are signs of outer capsule degeneration (*cf.* Maier, 1997; Elsohemy *et al*, 2009). Further, intrafusal calbindin staining was weak in control mice of this age, and no longer observable by immunohistology in coTxNrg1 animals (Figure 7H). NF200<sup>+</sup> terminals were however still present, indicating that sensory innervation was preserved in remaining spindles (arrowheads in Figure 7H). The analysis of inducible mutant mice thus demonstrated that continuous Nrg1 signalling is essential to sustain muscle spindles and maintain motor coordination in the adult.

### Discussion

Bace1 is well known for its role in Alzheimer's disease, and is required for APP cleavage and production of Aβ peptides (Luo *et al*, 2001). In this study, we show that Bace1 controls muscle spindle ontogenesis and maintenance in the adult. We combined complex genetic analyses, biochemical studies and pharmacological interference to provide evidence that these functions depend on cleavage of membrane-tethered IgNrg1 by Bace1. Our findings identify Bace1 as an important molecule in proprioception.





**Figure 5** Bace1 and IgNrg1 control the ontogeny of muscle spindles. (A) Quantification of muscle spindles in hindlimbs of newborn mice of indicated genotypes. (B) Immunohistological analyses of DRG from control, Bace1<sup>-/-</sup> and IgNrg1β1<sup>OV</sup> animals at E15 and P0 using antibodies directed against TrkC, NF200 and Islet1/2. (C, D) Quantification of TrkC<sup>+</sup> proprioceptive neurons (C) and Islet<sup>+</sup> sensory neurons (D) in DRG of E15 and P0 animals. TrkC<sup>+</sup> proprioceptive neurons persisted in increased numbers in newborn IgNrg1β1<sup>OV</sup>, while the overall quantity of Islet1/2<sup>+</sup> sensory neurons was comparable. Bace1, co-IgNrg1 and wild-type mice display similar numbers of sensory or proprioceptive neurons. Scale bar, 25 μm (B).

### Bace1 and Nrg1 control ontogenesis and maintenance of muscle spindles

Our genetic study provides direct evidence that an Ig-containing isoform of Nrg1 supplies the muscle spindle-inducing activity. This is in accordance with the previous work that demonstrated a role of Nrg1 but not of CRD-containing isoforms in spindle induction (Hippenmeyer *et al*, 2002). IgNrg1β1 is the dominantly expressed Ig-containing isoform of Nrg1 in DRG and thus expected to participate in muscle spindle induction. Our analysis of mice overexpressing membrane-tethered IgNrg1β1 in neurons indicates that this isoform induces supernumerary muscle spindles, but this depends on the presence of Bace1.


A direct contact exists between proprioceptive axons and muscle during muscle spindle differentiation (Ernfors *et al*, 1994; Farinas *et al*, 1994; Klein *et al*, 1994; Tessarollo *et al*, 1994; Walro and Kucera, 1999; Chen *et al*, 2003). Membrane of sensory axons and muscle fibres are thus in close apposition. Nevertheless, our data show that membrane-tethered IgNrg1β1 isoforms are not able to elicit a response without being shed. Further, we find that Nrg1 ablation or Bace1 inhibition in the adult result in a regression of the muscle spindle pool, indicating that Bace1-dependent Nrg1 activity is required to sustain muscle spindles even in the

presence of an intact nerve. Previous studies had demonstrated that a continuous contact between the intrafusal fibres and sensory axons is a prerequisite for the maintenance of muscle spindles in adults. For instance, adult nerve transection in rats results in acute morphological changes followed by a massive loss of muscle spindles (Coprav *et al*, 1999; Elsohemy *et al*, 2009). However, the molecular identity of the signal(s) provided by sensory axons had not been defined, and our data indicate that Nrg1 provides a key signal for spindle maintenance.

### Functions of neuronally produced Nrg1 isoforms during development and adulthood

We show here that IgNrg1β1 requires Bace1 for its activity as spindle-promoting factor. IgNrg1β1<sup>OV</sup> mice ectopically express this particular isoform and display supernumerary muscle spindles, a phenotype that is completely reversed when Bace1 is ablated. This is accompanied by increased survival of proprioceptive neurons in IgNrg1β1<sup>OV</sup>; it is interesting to note that supernumerary muscle spindles in IgNrg1β1<sup>OV</sup> mice express NT3, a known rate-limiting factor for survival of proprioceptive neurons (Ernfors *et al*, 1994; Klein *et al*, 1994; Tessarollo *et al*, 1994). The expression of NT3 in such supernumerary muscle spindles might account






**Figure 6** The maturation of muscle spindles depends on Bace1 and IgNrg1. (A) Immunohistological staining of adult (P30) muscle spindles from the hindlimbs of mice of indicated genotypes. The capsule (collagen IV<sup>+</sup>) and sensory innervation (NF200<sup>+</sup>) are observable in all genotypes. Note that the intensity of calbindin staining of intrafusal fibres depends on the dose of IgNrg1 and on Bace1 (quantified in Supplementary Figure S6A). (B–E) Distribution of the diameter (B, C) and number of intrafusal fibres (D, E) in muscle spindles of Bace1 and IgNrg1 mutants. Scale bar, 20 μm (A).

for the enhanced survival of proprioceptive neurons during the selection phase. Several findings indicate that not all Nrg1 activity present at sensory–muscular contact sites is provided by IgNrg1β1 and depends on Bace1 for signalling. Previous data had indicated that the induction of muscle spindles strictly depends on neuronal Nrg1, i.e., no spindles form in the absence of Nrg1 (Hippenmeyer *et al*, 2002), but we observed remaining muscle spindles in Bace1 mutants, although their number and size were reduced. Thus, Bace1-independent Nrg1 isoforms appear to participate in muscle spindle induction, for instance the secreted IgNrg1β3 (GGF) isoform that is also produced at low levels in sensory neurons. Alternately, metalloproteinases that are known to process Nrg1 might contribute to IgNrg1 shedding *in vivo* (Montero *et al*, 2000; Shirakabe *et al*, 2001; Horiuchi *et al*, 2005; La Marca *et al*, 2011; Luo *et al*, 2011). Sensory neurons provide the IgNrg1 signal for spindle induction. Because of their limited availability, the relative overabundance of CRD-Nrg1, and the lack of IgNrg1-specific antibodies, we were unable to directly assess IgNrg1 processing in sensory neurons. However, our experiments indicate that the endogenous Bace1 and metalloproteinases cooperatively cleave IgNrg1β1 in hippocampal neurons, similar to the shedding of CRD-Nrg1 described recently (Fleck *et al*, 2013). Further work is required, for instance by the use of mouse genetics, to assess the contribution of metalloproteinases to IgNrg1β1 cleavage in sensory neurons.

We observed subtle differences in maturation of muscle spindles in Bace1 and IgNrg1 mutant mice; for instance, IgNrg1 but not Bace1 mutants frequently display spindles with unusually few intrafusal fibres. Thus, Bace1-dependent and -independent IgNrg1 isoforms might assume slightly different functions during muscle spindle maturation. Alternatively, these differences in muscle spindle morphology might represent a response to graded differences in Nrg1 signals. In particular, co-IgNrg1 animals display a very severe

reduction in spindle numbers (84% reduction), significantly smaller spindle diameter and fewer intrafusal fibres in remaining spindles; compound Bace1<sup>-/-</sup> IgNrg1<sup>Δ/+</sup> mutants spindles are reduced in numbers (71% reduction) and display milder changes in spindle size and intrafusal fibre numbers; finally, in Bace1 mutant and inhibitor-treated mice, spindles are reduced in numbers (~50%), but numbers of intrafusal fibres are comparable to those in wild-type mice. Thus, a graded lowering of Nrg1 signals in this series of mutants appears to be translated into more and more pronounced reduction in numbers of muscle spindles and into increasingly pronounced severities of morphological deficits in remaining spindles.

Nrg1 generated by sensory neurons exerts two unrelated functions, muscle spindle induction and myelination, and both are disrupted in Bace1 mutant mice. This argues strongly for a Bace1-dependent shedding of Nrg1 in the control of these events. In particular, our biochemical data show that Bace1 processes CRD-Nrg1β1 sequences in cultured HEK293 cells. Unprocessed CRD-Nrg1 has an apparent MW of 130–140 kD when expressed in cultured cells, and a Nrg1 isoform of 130 kD accumulates in the brain of Bace1<sup>-/-</sup> or Ly2811376-treated mice (Willem *et al*, 2006 and this study). Furthermore, hypomyelination is observed in Bace1<sup>-/-</sup> and CRD-Nrg1<sup>+/-</sup> mice (Michailov *et al*, 2004; Willem *et al*, 2006). Together, these experiments indicated that CRD-Nrg1 requires Bace1-dependent processing to control myelination. A recent report demonstrated that hypermyelination associated with overexpression of CRD-Nrg1 is not completely blocked by a Bace1 mutation, although a trend that did not reach a statistical significance was observed (Velanac *et al*, 2012). Thus, other substrates but CRD-Nrg1 might cause or participate in the impaired myelination observed in Bace1 mutants. Alternately, the transgenic overexpression of CRD-Nrg1 might bypass the processing by Bace1. In this context, it is noteworthy that Bace1-dependent processing only occurs during transit of substrates through endosomes, whereas



**Figure 7** Continuous Nrg1 signalling is required to sustain muscle spindles in the adult. (A) Schematic outline of Nrg1 ablation and spindle analysis in adult coTxNrg1 mice (Tx, tamoxifen). (B) qPCR analysis of Nrg1 transcripts in DRG of control and coTxNrg1 mice at P100. (C, D) Performance of coTxNrg1 mice (P120) on the beam walk test. coTxNrg1 mice display frequent missteps and distinctive hopping movements; the latter is also a feature of co-IgNrg1 mice. (E) 3D reconstruction of muscle spindles (white lines) in the tibialis anterior muscle of control and coTxNrg1 mice (P180). (F) Quantification of numbers of muscle spindles in tibialis anterior muscle of control (black bars) and coTxNrg1 (red bars) animals. (G) Distribution of muscle spindle length in coTxNrg1 and control mice. (H) Immunohistological analysis of muscle spindles from control and coTxNrg1 animals. The presence of NF200<sup>+</sup> endings indicates that sensory projections contacted spindles in both genotypes (arrowheads), but calbindin was not detectable in intrafusal fibres of coTxNrg1 mice. In addition, collagen IV staining in outer capsules was weak and interrupted, indicating that capsules degenerated in remaining spindles of coTxNrg1 mice. (I) Schematic summary of gait impairment in Ly2811376-treated, *Bace1*<sup>-/-</sup>, co-IgNrg1 and coTxNrg1 animals. Each panel displays the directions of paw movements (arrows) during a series of two steps. Both *Bace1*<sup>-/-</sup> and *Bace1*-inhibited mice display defective anterior/posterior coordination resulting in a swaying walk. In addition, co-IgNrg1 and coTxNrg1 mice show more pronounced deficits of posterior homologue coupling resulting in frequent hopping. Scale bars, 1.5 mm (E); 15 μm (H).

other proteases like those of the metalloproteinase family are active at the plasma membrane (Edwards *et al*, 2008; Willem *et al*, 2009; Weber and Saftig, 2012). In such case, metalloproteinases could contribute to enhanced and Bace1-independent Nrg1 signalling in mice overexpressing CRD-Nrg1. Alternatively, transgenic CRD-Nrg1 overexpression might saturate ErbB signalling; hence Schwann cells would no longer be affected by a reduction of bioactive Nrg1 caused by Bace1 mutation.

Two isoforms, Ig- and CRD-Nrg1, are expressed in sensory neurons (Meyer *et al*, 1997). We show here that IgNrg1 induces formation of muscle spindles, but is dispensable for myelination. In contrast, CRD-Nrg1 regulates the development of Schwann cells and myelination, but is not required for muscle spindle formation (Meyer *et al*, 1997; Wolpowitz *et al*, 2000; Hippenmeyer *et al*, 2002). The question arises of how Ig- and CRD-Nrg1 isoforms produced in the same sensory neurons exert their distinct roles. Both Nrg1 isoforms are cooperatively cleaved by Bace1 and metalloproteinases (this work, Willem *et al*, 2006; Fleck *et al*, 2013). Further, similar downstream signalling molecules such as ErbB2 and CREB participate in myelination and in muscle spindle induction (Garratt *et al*, 2000; Andrechek *et al*, 2002; Leu *et al*, 2003; Arthur-Farraj *et al*, 2011; Herndon *et al*, 2013). Divergent intra- or intercellular trafficking mechanisms might exist that result in separate subcellular localization and function of Ig- and CRD-Nrg1 isoforms.

### Functions of Bace1 in the mature nervous system

We demonstrate here that continuous Bace1 activity is required in the adult to sustain muscle spindles and therefore maintain accurate proprioception and coordinated movement. Bace1 was assigned various developmental functions, for instance in myelination and vascularization of the eye (Hu *et al*, 2006; Willem *et al*, 2006; Cai *et al*, 2012). Further behavioural deficits (prepulse inhibition, novelty-induced hyperactivity, social recognition) were observed, which might originate from disrupted synapse formation and maturation (Stefansson *et al*, 2002; Laird *et al*, 2005; Savonenko *et al*, 2008; Wang *et al*, 2008). Bace1 inhibition is considered a promising route for the treatment of Alzheimer's disease, and is expected to interfere with A $\beta$  generation and the formation of pathogenic A $\beta$  aggregates (Haass, 2004; Luo *et al*, 2011). To this date, several Bace1 inhibitors are tested in clinical phase II and planned for phase III. Adverse effects due to the inhibition of physiological functions of the protease must be considered. The data presented here indicate that one unwanted side effect of a long-term inhibition of Bace1 in adults might be disrupted muscle spindle functions, which are expected to impair coordinated movement.

## Materials and methods

### Animal strains and generation of IgNrg1<sup>flox</sup> and IgNrg1<sup>A</sup> alleles

Bace1<sup>-/-</sup> and IgNrg1 $\beta$ 1<sup>Ov</sup> (Nrg1 type I  $\beta$ 1 overexpression under the control of the Thy 1.2 promoter) strains have been described (Michailov *et al*, 2004; Dominguez *et al*, 2005). Exons 3 and 4 of Nrg1 encode the Ig-like domain (IgNrg1). IgNrg1<sup>flox</sup> mutants contain LoxP sites 5' and 3' of exon 4, which were inserted by homologous recombination into embryonic stem (ES) cells. Targeted ES cells were used to generate a mutant strain using standard techniques. The IgNrg1<sup>A</sup> strain was generated by crossing IgNrg1<sup>flox</sup> with Deleter<sup>cre</sup> animals (Schwenk *et al*, 1995),

in which cre recombinase is expressed in all tissues including germ cells, thus inducing complete recombination. We obtained IgNrg1<sup>A</sup> mice upon backcrossing Deleter<sup>cre</sup>IgNrg1<sup>flox</sup> animals with wild-type animals. The Wnt1-cre stain expresses cre in neural crest cells, causing recombination in the progenitors that give rise to sensory neurons (Danielian *et al*, 1998; Le Douarin and Kalcheim, 1999). Wnt1<sup>cre</sup>IgNrg1<sup>flox/A</sup> (co-IgNrg1) mice were used for further analysis. Krox20<sup>cre/+</sup>ErbB2<sup>flox/flox</sup> animals (here also called coErbB2) present severe hypomyelination similar to Bace1<sup>-/-</sup> mice, and were previously described (Garratt *et al*, 2000). Phenotypes were analysed on a mixed C57Bl/6;129Ola background and mutants always compared with littermates.

### Bace1 inhibition in adult mice

*In vivo* pharmacology studies were conducted with wild type, heterozygous and homozygous Bace1 mutant mice (EPO Berlin-Buch GmbH). Briefly, the Bace1 inhibitor Ly2811376 (Eli Lilly) was prepared as a 160  $\mu$ g/ $\mu$ l stock solution in pharماسolve (ISP technologies Inc), and aliquots were diluted 1/16 in corn oil (Ly2811376 concentration: 10  $\mu$ g/ $\mu$ l). The animals were treated daily with Ly2811376 at a dose of 100  $\mu$ g/g of body weight/day for 29 consecutive days, or a corresponding volume of vehicle solution, a treatment that did not change body weight. Animals were sacrificed within 4 h after the last treatment. Brain tissue was snap frozen and kept at -80°C until further processing.

### Ablation of Nrg1 expression during adulthood

The cre-ER<sup>TM</sup> and Nrg1<sup>flox/flox</sup> strains have been described (Yang *et al*, 2001; Hayashi and McMahon, 2002). Briefly, in cre-ER<sup>TM</sup> mice the cre-ER<sup>TM</sup> fusion protein is ubiquitously expressed under the control of CAG promoter (CMV enhancer/chicken beta-actin promoter, *cf.* Hayashi and McMahon, 2002). We used a previously established colony of cre-ER<sup>TM</sup>Nrg1<sup>flox/flox</sup> mice, in which cre-mediated recombination of Nrg1<sup>flox</sup> alleles only occurs after tamoxifen binds to the oestrogen receptor domain (ER<sup>TM</sup>) of cre-ER<sup>TM</sup> (Fricker *et al*, 2013). Recombined Nrg1<sup>flox</sup> alleles generate truncated Nrg1 mRNAs lacking a functional EGF domain (Yang *et al*, 2001). Ten-week-old cre-ER<sup>TM</sup>Nrg1<sup>flox/flox</sup> mice were treated with tamoxifen (Fricker *et al*, 2013). Animals were kept for further 16 weeks before dissection. Cre-ER<sup>TM</sup>Nrg1<sup>flox/flox</sup> animals treated with tamoxifen are referred as coTxNrg1, and were compared with three control groups (cre-ER<sup>TM</sup>Nrg1<sup>flox/flox</sup> vehicle treated; cre-ER<sup>TM</sup>Nrg1<sup>+/+</sup> tamoxifen treated; Nrg1<sup>flox/flox</sup> tamoxifen treated). All control groups displayed similar characteristics and were therefore grouped for the spindle length analysis.

### Animal handling and assessment of motor coordination

The motor coordination of 5–7 months' old Ly2811376-treated, Bace1 and co-IgNrg1 mutant mice were assessed using gait analysis (Treadscan, Clever Sys Inc. Beare *et al*, 2009) and inverted grid tests (Coughenour *et al*, 1977; Landauer *et al*, 2003). We assessed the motor coordination of the cohort of 4-month-old coTxNrg1 using beam walking (as described in Carter *et al*, 2001; Baskin *et al*, 2003; Taylor *et al*, 2005). All experiments were conducted in accordance with institutional German regulations and home office guidelines in UK, respectively.

### Double in situ hybridization and immunohistochemistry

Fluorescent double *in situ* hybridization and immunohistological analyses were performed as described (Wende *et al*, 2012) and acquired using Zeiss LSM700 confocal microscope. The sequences of IgNrg1 and Bace1 ISH probes correspond to the 5'UTR and type I-specific coding sequence of Nrg1, or the coding sequence and 3'UTR of Bace1, respectively. Antibodies used are Egr3 (rabbit, 1:300, Santa Cruz), Neurofilament 200 (chicken, 1:20 000, Millipore), calbindin (rabbit, 1:500, Swant), collagen IV (goat, 1:1000, Millipore), S100 (rabbit, 1:1000, Dakocytomation), TrkC (rat, 1:500, R&D Systems), Islet1/2 (mouse, 1:200, DSHB). DAPI (250 nM, Invitrogen) was used as nuclear counter stain.

Colocalization of Bace1 and IgNrg1 transcripts in DRG neurons was performed on eight DRG sections from four animals. TrkC<sup>+</sup> or Islet<sup>+</sup> sensory neurons were quantified from serial sections (12  $\mu$ m) of L5 DRG, in the five centremost sections of 3–6 animals/genotype/age. Only TrkC<sup>+</sup> cells displaying nuclei in the focal plane were counted.



### Muscle spindle identification and quantification

The number of muscle spindles per section were counted in lower hindlimbs or the tibialis anterior of 4–6 animals per genotype/treatment. Wild-type littermates were used as controls. The lower hindlimb (P0/P30) or the tibialis anterior (P30/P180) were sectioned transversally (60 µm), every fourth section was stained, and five (P0 animals) or 10 (adult animals) sections of the center of the muscle were evaluated per animal. Muscle spindles were defined as encapsulated structures displaying specific morphological hallmarks (nuclear clustering, periaxial space of Sherrington) and containing at least two intrafusal fibres expressing Egr3 (P0) or calbindin (adult). Morphological criteria and immunohistological signals were used to prevent an underestimation of muscle spindle numbers caused by loss of innervation or reduction of Egr3/calbindin expression in mutant strains. Expression of calbindin in intrafusal fibres was particularly weak in P180 control mice, and absent in age-matched coTxNrg1 mutants. We therefore used morphological criteria (nuclear aggregation, space of Sherrington) to identify muscle spindles in P180 animals.

Muscle spindle diameter and intrafusal fibre counts were obtained from immunohistological sections and were determined at the level of the annulospiral endings (equatorial plane), identified by NF200 staining; 25 muscle spindles/animal were analysed. The area enclosed by the outer capsule of the spindle was measured using ImageJ software 1.46j. Intrafusal fibres were identified using Egr3/calbindin staining and cell morphology. Each point represents mean ± s.e.m. of data obtained from 5–6 animals per genotype.

For 3D reconstructions of the tibialis anterior muscle, pictures of sections were assembled as a stack. On individual pictures, the surface of the muscle was outlined and the position of muscle spindles was indicated. Muscle spindles were then aligned for reconstructions using ImageJ 1.46j. Muscle spindle length was estimated on reconstructed muscle, data points represent mean ± s.e.m. of 3–6 animals per condition.

### Cell culture and biochemical analyses

The Bace1 plasmid was described in Westmeyer *et al* (2004). For HEK293 cell culture experiments, IgNrg1β1/2 (type I isoforms) and CRD-Nrg1β1 cDNAs were inserted into pSecTag and pcDNA3.1 expression plasmids, respectively. IgNrg1β1 and β2 carry an HA tag close to the N-terminus, while CRD-Nrg1β1 bears its HA tag between EGF and transmembrane domains (Fleck *et al*, 2013). In constructs used for SEAP assay, the alkaline phosphatase sequences substituted sequences of the type I-specific and Ig domains of Nrg1 (Willem *et al*, 2006). The sequences of all constructs were verified by sequencing.

HEK293 cells were cultured in DMEM + 10% FCS + penicillin/streptomycin (Gibco-BRL Invitrogen) and transfected (5 µg DNA) using Lipofectamine 2000 (Invitrogen) following the manufacturer's instructions. The hippocampal neurons were isolated from embryonic day 18 rats as described previously (Kaech and Banker, 2006). Dissociated neurons were plated at 17 700 cells per cm<sup>2</sup> onto 6 cm dishes coated with poly-L-lysine (1 mg/ml; Sigma) and cultured in Neurobasal medium supplemented with 2% B27 and 0.5 mM L-glutamine (all from Invitrogen). We transfected primary neurons after 4 days of culture with pSecTag-HA-IgNrg1β1 plasmid (5 µg DNA) using Lipofectamine 2000. After 9 days in culture, cells were incubated for 24 h in the presence/absence of Bace1 (C3; 0.2 µM) or metalloproteinase (GM6001; 20 µM) inhibitors (EMD Biosciences, Merck). Supernatants and cells were independently collected either 24 h post-transfection (HEK293) or following the inhibitor treatment (primary neurons) for western blot analysis of Nrg1 cleavage. Supernatants were cleared by centrifugation and cells were lysed

in RIPA buffer containing protease inhibitor mix (Sigma P8340). Immunoprecipitation from neuronal supernatants was performed with anti-HA Agarose (Sigma A2095; 25 mg/ml) and washed according to standard protocols (Fleck *et al*, 2013). Western blots were performed as described (Fleck *et al*, 2013) using the 4F10 antibody, which recognizes an epitope that is exclusively presented by Bace1-cleaved N-terminal Nrg1 fragments (1:40) and commercial antibodies directed against C-terminal Nrg1 (sc348, Santa Cruz, 1:2000), Bace1 (D10E5, Cell signalling, 1:1000), calnexin (AF18, Enzo, 1:5000) and HA (3F10-HRP coupled, Roche, 1:10 000). Secondary anti-IgG-HRP antibodies were from Santa Cruz (1:2000) or Promega (1:10 000).

For SEAP analysis, the medium was changed 24 h after transfection and cells were incubated for 16 h in the presence/absence of 5 µM C3 and 20 µM GM6001 (EMD Biosciences, Merck). The SEAP assay was performed as described (Willem *et al*, 2006).

### Semi-quantitative PCR

qPCR was essentially performed as described (Wende *et al*, 2012). The primers used are shown in Supplementary Table I. In Nrg1 isotype analyses, the amount of CRD-Nrg1 transcripts in control DRG was set to 100%, and amounts of other isoforms in control and mutant DRG were displayed as proportion of this value.

### Statistics

Data are presented as mean ± s.e.m. The differences observed were assessed by ANOVA followed by 2-tailed unpaired Student's modified *t*-test. Significance is displayed as follows: <sup>NS</sup>*P* > 0.05; \**P* < 0.05; \*\**P* < 0.01; \*\*\**P* < 0.001. Supplementary Table II details the statistical analysis for the quantification of muscle spindles in newborn Bace1 and IgNrg1 mutants presented in Figure 5A.

### Supplementary data

Supplementary data are available at *The EMBO Journal* Online (<http://www.embojournal.org>).

## Acknowledgements

We thank Bettina Brandt, Carola Griffel, Nadine Baseler, Ana Antunes-Martins, Heike Hampel and Andrea Seibel for technical support; Petra Stallerow and Claudia Päseler for animal husbandry; Walter Birchmeier and Dominique Bröhl for critical reading of the manuscript; and Christiane Volbracht (H. Lundbeck A/S (Copenhagen)) for the gift of the Ly2811376 Bace1 inhibitor. DLB is a Wellcome Trust Senior Clinical Scientist (095698/z/11/z). This work was supported by HelMA (Helmholtz Association) and BMBF (NGFNplus IG Alzheimer).

**Author contributions:** CC performed histological and behavioural analysis. MW, ST and CH performed and analysed biochemical experiments; FRF and DLB provided tissue from and performed behavioural testing of coTxNrg1 mice; HW generated the IgNrg1 strain and performed ISH experiments. AW-G supervised Ly2811376 treatment; KAN and PS provided IgNrg1β1<sup>0v</sup> and Bace1<sup>-/-</sup> strains, respectively; and ANG analysed myelination. CB and CC supervised the project, designed experiments and wrote the article.

## Conflict of interest

The authors declare that they have no conflict of interest.

## References

- Andrechek ER, Hardy WR, Gargis-Gabardo AA, Perry RL, Butler R, Graham FL, Kahn RC, Rudnicki MA, Muller WJ (2002) ErbB2 is required for muscle spindle and myoblast cell survival. *Mol Cell Biol* **22**: 4714–4722
- Arthur-Farraj P, Wanek K, Hantke J, Davis CM, Jayakar A, Parkinson DB, Mirsky R, Jessen KR (2011) Mouse schwann cells need both NRG1 and cyclic AMP to myelinate. *Glia* **59**: 720–733
- Baskin YK, Dietrich WD, Green EJ (2003) Two effective behavioral tasks for evaluating sensorimotor dysfunction following traumatic brain injury in mice. *J Neurosci Methods* **129**: 87–93
- Beare JE, Morehouse JR, DeVries WH, Enzmann GU, Burke DA, Magnuson DS, Whittemore SR (2009) Gait analysis in normal and spinal contused mice using the TreadScan system. *J Neurotrauma* **26**: 2045–2056

- Birchmeier C (2009) ErbB receptors and the development of the nervous system. *Exp Cell Res* **315**: 611–618
- Cai J, Qi X, Kociok N, Skosyrski S, Emilio A, Ruan Q, Han S, Liu L, Chen Z, Bowes Rickman C, Golde T, Grant MB, Saftig P, Serneels L, de Strooper B, Jousen AM, Boulton ME (2012) beta-Secretase (BACE1) inhibition causes retinal pathology by vascular dysregulation and accumulation of age pigment. *EMBO Mol Med* **4**: 980–991
- Carter RJ, Morton J, Dunnett SB (2001) Motor coordination and balance in rodents. *Curr Protoc Neurosci* Chapter 8, Unit 8, 12
- Chen HH, Hippenmeyer S, Arber S, Frank E (2003) Development of the monosynaptic stretch reflex circuit. *Curr Opin Neurobiol* **13**: 96–102
- Chen HH, Tourtellotte WG, Frank E (2002) Muscle spindle-derived neurotrophin 3 regulates synaptic connectivity between muscle sensory and motor neurons. *J Neurosci* **22**: 3512–3519
- Copray JC, Bastiaansen M, Gibbons H, van Roon WM, Comer AM, Lipski J (1999) Neurotrophic requirements of rat embryonic catecholaminergic neurons from the rostral ventrolateral medulla. *Brain Res* **116**: 217–222
- Coughenour LL, McLean JR, Parker RB (1977) A new device for the rapid measurement of impaired motor function in mice. *Pharmacol Biochem Behav* **6**: 351–353
- Danielian PS, Muccino D, Rowitch DH, Michael SK, McMahon AP (1998) Modification of gene activity in mouse embryos *in utero* by a tamoxifen-inducible form of Cre recombinase. *Curr Biol* **8**: 1323–1326
- Dominguez D, Tournoy J, Hartmann D, Huth T, Cryns K, Deforce S, Serneels L, Camacho IE, Marjaux E, Craessaerts K, Roebroek AJ, Schwake M, D'Hooge R, Bach P, Kalinke U, Moechars D, Alzheimer C, Reiss K, Saftig P, De Strooper B (2005) Phenotypic and biochemical analyses of BACE1- and BACE2-deficient mice. *J Biol Chem* **280**: 30797–30806
- Edwards DR, Handsley MM, Pennington CJ (2008) The ADAM metalloproteinases. *Mol Aspects Med* **29**: 258–289
- Elsohemy A, Butler R, Bain JR, Fahnestock M (2009) Sensory protection of rat muscle spindles following peripheral nerve injury and reinnervation. *Plast Reconstr Surg* **124**: 1860–1868
- Ernfors P, Lee KF, Kucera J, Jaenisch R (1994) Lack of neurotrophin-3 leads to deficiencies in the peripheral nervous system and loss of limb proprioceptive afferents. *Cell* **77**: 503–512
- Falls DL (2003) Neuregulins: functions, forms, and signaling strategies. *Exp Cell Res* **284**: 14–30
- Farinas I, Jones KR, Backus C, Wang XY, Reichardt LF (1994) Severe sensory and sympathetic deficits in mice lacking neurotrophin-3. *Nature* **369**: 658–661
- Fleck D, van Bebber F, Colombo A, Galante C, Schwenk BM, Rabe L, Hampel H, Novak B, Kremmer E, Tahirovic S, Edbauer D, Lichtenthaler SF, Schmid B, Willem M, Haass C (2013) Dual Cleavage of Neuregulin 1 Type III by BACE1 and ADAM17 Liberates Its EGF-Like Domain and Allows Paracrine Signaling. *J Neurosci* **33**: 7856–7869
- Freese C, Garratt AN, Fahrenholz F, Endres K (2009) The effects of alpha-secretase ADAM10 on the proteolysis of neuregulin-1. *FEBS J* **276**: 1568–1580
- Fricker FR, Antunes-Martins A, Galino J, Paramsothy R, La Russa F, Perkins J, Goldberg R, Brelstaff J, Zhu N, McMahon S, Orengo C, Garratt A, Birchmeier C, Bennett BD (2013) Axonal Neuregulin-1 is a rate limiting but not essential factor for nerve remyelination. *Brain* (In press)
- Fricker FR, Lago N, Balarajah S, Tsantoulas C, Tanna S, Zhu N, Fageiry SK, Jenkins M, Garratt AN, Birchmeier C, Bennett DL (2011) Axonally derived neuregulin-1 is required for remyelination and regeneration after nerve injury in adulthood. *J Neurosci* **31**: 3225–3233
- Garratt AN, Voiculescu O, Topilko P, Charnay P, Birchmeier C (2000) A dual role of erbB2 in myelination and in expansion of the schwann cell precursor pool. *J Cell Biol* **148**: 1035–1046
- Gould TW, Yonemura S, Oppenheim RW, Ohmori S, Enomoto H (2008) The neurotrophic effects of glial cell line-derived neurotrophic factor on spinal motoneurons are restricted to fusimotor subtypes. *J Neurosci* **28**: 2131–2146
- Grossmann KS, Wende H, Paul FE, Cheret C, Garratt AN, Zurborg S, Feinberg K, Besser D, Schulz H, Peles E, Selbach M, Birchmeier W, Birchmeier C (2009) The tyrosine phosphatase Shp2 (PTPN11) directs Neuregulin-1/ErbB signaling throughout Schwann cell development. *Proc Natl Acad Sci USA* **106**: 16704–16709
- Haass C (2004) Take five–BACE and the gamma-secretase quartet conduct Alzheimer's amyloid beta-peptide generation. *EMBO J* **23**: 483–488
- Hayashi S, McMahon AP (2002) Efficient recombination in diverse tissues by a tamoxifen-inducible form of Cre: a tool for temporally regulated gene activation/inactivation in the mouse. *Dev Biol* **244**: 305–318
- Herndon CA, Ankenbruck N, Lester B, Bailey J, Fromm L (2013) Neuregulin1 signaling targets SRF and CREB and activates the muscle spindle-specific gene Egr3 through a composite SRF-CREB-binding site. *Exp Cell Res* **319**: 718–730
- Hippenmeyer S, Shneider NA, Birchmeier C, Burden SJ, Jessell TM, Arber S (2002) A role for neuregulin1 signaling in muscle spindle differentiation. *Neuron* **36**: 1035–1049
- Horiuchi K, Zhou HM, Kelly K, Manova K, Blobel CP (2005) Evaluation of the contributions of ADAMs 9, 12, 15, 17, and 19 to heart development and ectodomain shedding of neuregulins beta1 and beta2. *Dev Biol* **283**: 459–471
- Hu X, He W, Diaconu C, Tang X, Kidd GJ, Macklin WB, Trapp BD, Yan R (2008) Genetic deletion of BACE1 in mice affects remyelination of sciatic nerves. *FASEB J* **22**: 2970–2980
- Hu X, Hicks CW, He W, Wong P, Macklin WB, Trapp BD, Yan R (2006) Bace1 modulates myelination in the central and peripheral nervous system. *Nat Neurosci* **9**: 1520–1525
- Hunt CC (1990) Mammalian muscle spindle: peripheral mechanisms. *Physiol Rev* **70**: 643–663
- Kaech S, Banker G (2006) Culturing hippocampal neurons. *Nat Protoc* **1**: 2406–2415
- Kandalepas PC, Vassar R (2012) Identification and biology of beta-secretase. *J Neurochem* **120**(Suppl 1): 55–61
- Klein R, Silos-Santiago I, Smeyne RJ, Lira SA, Brambilla R, Bryant S, Zhang L, Snider WD, Barbacid M (1994) Disruption of the neurotrophin-3 receptor gene trkC eliminates the muscle afferents and results in abnormal movements. *Nature* **368**: 249–251
- Kuhn PH, Koroniak K, Hogl S, Colombo A, Zeitschel U, Willem M, Volbracht C, Schepers U, Imhof A, Hoffmeister A, Haass C, Rossner S, Brase S, Lichtenthaler SF (2012) Secretome protein enrichment identifies physiological BACE1 protease substrates in neurons. *EMBO J* **31**: 3157–3168
- La Marca R, Cerri F, Horiuchi K, Bachi A, Feltri ML, Wrabetz L, Blobel CP, Quattrini A, Salzer JL, Taveggia C (2011) TACE (ADAM17) inhibits Schwann cell myelination. *Nat Neurosci* **14**: 857–865
- Laird FM, Cai H, Savonenko AV, Farah MH, He K, Melnikova T, Wen H, Chiang HC, Xu G, Koliatsos VE, Borchelt DR, Price DL, Lee HK, Wong PC (2005) BACE1, a major determinant of selective vulnerability of the brain to amyloid-beta amyloidogenesis, is essential for cognitive, emotional, and synaptic functions. *J Neurosci* **25**: 11693–11709
- Landauer MR, Srinivasan V, Seed TM (2003) Genistein treatment protects mice from ionizing radiation injury. *J Appl Toxicol* **23**: 379–385
- Le Douarin N, Kalcheim C (1999) *The neural crest*. Cambridge University Press
- Leu M, Bellmunt E, Schwander M, Farinas I, Brenner HR, Muller U (2003) ErbB2 regulates neuromuscular synapse formation and is essential for muscle spindle development. *Development* **130**: 2291–2301
- Luo X, Prior M, He W, Hu X, Tang X, Shen W, Yadav S, Kiryu-Seo S, Miller R, Trapp BD, Yan R (2011) Cleavage of neuregulin-1 by BACE1 or ADAM10 protein produces differential effects on myelination. *J Biol Chem* **286**: 23967–23974
- Luo Y, Bolon B, Kahn S, Bennett BD, Babu-Khan S, Denis P, Fan W, Kha H, Zhang J, Gong Y, Martin L, Louis JC, Yan Q, Richards WG, Citron M, Vassar R (2001) Mice deficient in BACE1, the Alzheimer's beta-secretase, have normal phenotype and abolished beta-amyloid generation. *Nat Neurosci* **4**: 231–232
- Maier A (1997) Development and regeneration of muscle spindles in mammals and birds. *Int J Dev Biol* **41**: 1–17
- May PC, Dean RA, Lowe SL, Martenyi F, Sheehan SM, Boggs LN, Monk SA, Mathes BM, Morgott DJ, Watson BM, Stout SL, Timm DE, Smith Labell E, Gonzales CR, Nakano M, Jhee SS, Yen M, Ereshefsky L, Lindstrom TD, Calligaro DO *et al* (2011) Robust central reduction of amyloid-beta in humans with an orally

- available, non-peptidic beta-secretase inhibitor. *J Neurosci* **31**: 16507–16516
- Meyer D, Yamaai T, Garratt A, Riethmacher-Sonnenberg E, Kane D, Theill LE, Birchmeier C (1997) Isoform-specific expression and function of neuregulin. *Development* **124**: 3575–3586
- Michailov GV, Sereda MW, Brinkmann BG, Fischer TM, Haug B, Birchmeier C, Role L, Lai C, Schwab MH, Nave KA (2004) Axonal neuregulin-1 regulates myelin sheath thickness. *Science* **304**: 700–703
- Montero JC, Yuste L, Diaz-Rodriguez E, Esparis-Ogando A, Pandiella A (2000) Differential shedding of transmembrane neuregulin isoforms by the tumor necrosis factor-alpha-converting enzyme. *Mol Cell Neurosci* **16**: 631–648
- Savonenko AV, Melnikova T, Laird FM, Stewart KA, Price DL, Wong PC (2008) Alteration of BACE1-dependent NRG1/ErbB4 signaling and schizophrenia-like phenotypes in BACE1-null mice. *Proc Natl Acad Sci USA* **105**: 5585–5590
- Schwenk F, Baron U, Rajewsky K (1995) A cre-transgenic mouse strain for the ubiquitous deletion of loxP-flanked gene segments including deletion in germ cells. *Nucleic Acids Res* **23**: 5080–5081
- Shirakabe K, Wakatsuki S, Kurisaki T, Fujisawa-Sehara A (2001) Roles of Meltrin beta/ADAM19 in the processing of neuregulin. *J Biol Chem* **276**: 9352–9358
- Stefansson H, Sigurdsson E, Steinthorsdottir V, Bjornsdottir S, Sigmundsson T, Ghosh S, Brynjolfsson J, Gunnarsdottir S, Ivarsson O, Chou TT, Hjaltason O, Birgisdottir B, Jonsson H, Gudnadottir VG, Gudmundsdottir E, Bjornsson A, Ingvarsson B, Ingason A, Sigfusson S, Hardardottir H *et al* (2002) Neuregulin 1 and susceptibility to schizophrenia. *Am J Hum Genet* **71**: 877–892
- Taylor MD, Holdeman AS, Weltmer SG, Ryals JM, Wright DE (2005) Modulation of muscle spindle innervation by neurotrophin-3 following nerve injury. *Exp Neurol* **191**: 211–222
- Tessarollo L, Vogel KS, Palko ME, Reid SW, Parada LF (1994) Targeted mutation in the neurotrophin-3 gene results in loss of muscle sensory neurons. *Proc Natl Acad Sci USA* **91**: 11844–11848
- Tourtellotte WG, Keller-Peck C, Milbrandt J, Kucera J (2001) The transcription factor Egr3 modulates sensory axon-myotube interactions during muscle spindle morphogenesis. *Dev Biol* **232**: 388–399
- Tourtellotte WG, Milbrandt J (1998) Sensory ataxia and muscle spindle agenesis in mice lacking the transcription factor Egr3. *Nat Genet* **20**: 87–91
- Vassar R, Kovacs DM, Yan R, Wong PC (2009) The beta-secretase enzyme BACE in health and Alzheimer's disease: regulation, cell biology, function, and therapeutic potential. *J Neurosci* **29**: 12787–12794
- Velanac V, Unterbarnscheidt T, Hinrichs W, Gummert MN, Fischer TM, Rossner MJ, Trimarco A, Brivio V, Taveggia C, Willem M, Haass C, Mobius W, Nave KA, Schwab MH (2012) Bace1 processing of NRG1 type III produces a myelin-inducing signal but is not essential for the stimulation of myelination. *Glia* **60**: 203–217
- Walro JM, Kucera J (1999) Why adult mammalian intrafusal and extrafusal fibers contain different myosin heavy-chain isoforms. *Trends Neurosci* **22**: 180–184
- Wang H, Song L, Laird F, Wong PC, Lee HK (2008) BACE1 knock-outs display deficits in activity-dependent potentiation of synaptic transmission at mossy fiber to CA3 synapses in the hippocampus. *J Neurosci* **28**: 8677–8681
- Weber S, Saftig P (2012) Ectodomain shedding and ADAMs in development. *Dev* **139**: 3693–3709
- Wende H, Lechner SG, Cheret C, Bourane S, Kolanczyk ME, Pattyn A, Reuter K, Munier FL, Carroll P, Lewin GR, Birchmeier C (2012) The transcription factor c-Maf controls touch receptor development and function. *Sci* **335**: 1373–1376
- Westmeyer GG, Willem M, Lichtenthaler SF, Lurman G, Multhaup G, Assfalg-Machleidt I, Reiss K, Saftig P, Haass C (2004) Dimerization of beta-site beta-amyloid precursor protein-cleaving enzyme. *J Biol Chem* **279**: 53205–53212
- Willem M, Garratt AN, Novak B, Citron M, Kaufmann S, Rittger A, De Strooper B, Saftig P, Birchmeier C, Haass C (2006) Control of peripheral nerve myelination by the beta-secretase BACE1. *Science* **314**: 664–666
- Willem M, Lammich S, Haass C (2009) Function, regulation and therapeutic properties of beta-secretase (BACE1). *Semin Cell Dev Biol* **20**: 175–182
- Wolpowitz D, Mason TB, Dietrich P, Mendelsohn M, Talmage DA, Role LW (2000) Cysteine-rich domain isoforms of the neuregulin-1 gene are required for maintenance of peripheral synapses. *Neuron* **25**: 79–91
- Yang X, Arber S, William C, Li L, Tanabe Y, Jessell TM, Birchmeier C, Burden SJ (2001) Patterning of muscle acetylcholine receptor gene expression in the absence of motor innervation. *Neuron* **30**: 399–410
- Zelena J, Soukup T (1993) Increase in the number of intrafusal muscle fibres in rat muscles after neonatal motor denervation. *Neuroscience* **52**: 207–218
- Zhou L, Barao S, Laga M, Bockstael K, Borgers M, Gijzen H, Annaert W, Moechars D, Mercken M, Gevaert K, De Strooper B (2012) The neural cell adhesion molecules L1 and CHL1 are cleaved by BACE1 protease *in vivo*. *J Biol Chem* **287**: 25927–25940



The EMBO Journal is published by Nature Publishing Group on behalf of the European Molecular Biology Organization. This article is licensed under a Creative Commons Attribution-NonCommercial-Share Alike 3.0 Unported Licence. To view a copy of this licence visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>.