

OPEN ACCESS

**Repository of the Max Delbrück Center for Molecular Medicine (MDC)
in the Helmholtz Association**

<http://edoc.mdc-berlin.de/14987>

Intake of meat mutagens and risk of prostate cancer in a cohort of U.S. health professionals

Rohrmann, S., Nimptsch, K., Sinha, R., Willett, W.C., Giovannucci, E.L., Platz, E.A., Wu, K.

This is the final version of the manuscript. The original article has been published in final edited form in:

Cancer Epidemiology, Biomarkers & Prevention

2015 OCT ; 24(10): 1557-1563

Published first July 29, 2015

doi: [10.1158/1055-9965.EPI-15-0068-T](https://doi.org/10.1158/1055-9965.EPI-15-0068-T)

Publisher: copyright ©2015 [American Association for Cancer Research](#)

Intake of meat mutagens and risk of prostate cancer in a cohort of U.S. health professionals

Sabine Rohrmann^{1*}, Katharina Nimptsch^{2,4}, Rashmi Sinha³, Walter C Willett^{4,5,6},
Edward L. Giovannucci^{4,5,6}, Elizabeth A Platz^{7,8,9**}, and Kana Wu^{4**}

*Corresponding author

**Co-Senior Authors

¹ Division of Chronic Disease Epidemiology, Epidemiology, Biostatistics and Prevention Institute (EBPI), University of Zurich, Zurich, Switzerland

² Molecular Epidemiology Group, Max Delbrueck Center for Molecular Medicine, Molecular Epidemiology Group, Berlin, Germany

³ Nutritional Epidemiology Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, 6120 Executive Blvd, Rockville, MD

⁴ Department of Nutrition, Harvard School of Public Health, Boston, MA

⁵ Department of Epidemiology, Harvard School of Public Health, Boston, MA

⁶ Channing Division of Network Medicine, Department of Medicine, Brigham & Women's Hospital and Harvard Medical School, Boston, MA

⁷ Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD

⁸ Department of Urology and the James Buchanan Brady Urological Institute, Johns Hopkins School of Medicine, Baltimore, MD

⁹Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins, Baltimore, MD

Running title: Heterocyclic amines and risk of prostate cancer

Financial support: This study was supported by National Institutes of Health grants UM1 CA167552 (to WC Willett), P01 CA055075 (to WC Willett), and R01 CA133891 (to EL Giovannucci).

Corresponding author:

Sabine Rohrmann

Division of Chronic Disease Epidemiology

Epidemiology, Biostatistics and Prevention Institute (EBPI)

University of Zurich

Hirschengraben 84

8001 Zurich

Switzerland

e-mail: sabine.rohrmann@ifpsm.uzh.ch

phone: +41 44 634 5256

fax: +41 44 634 5444

Conflicts of interest: None.

word count (excluding references): 3312

number of figures: 0

number of tables: 3

Abstract

Background: Evidence relating heterocyclic aromatic amines (HCA), associated with high-temperature cooking methods, to prostate cancer risk is inconsistent

Methods: In a large US cohort study, intakes of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP), 2-amino-3,8-dimethylimidazo[4,5-f]quinoxaline (MeIQx) and 2-amino-3,4,8-trimethylimidazo[4,5-f]quinoxaline (DiMeIQx) and a meat-derived mutagenicity index (MDM) were assessed using a cooking method questionnaire administered in 1996. Until 2010, 2,770 prostate cancer cases were observed among 26,030 participants.

Results: Intake of PhIP from red meat was statistically significantly associated with total prostate cancer risk (top vs. bottom quintile HR=1.18, 95% CI 1.03-1.35), but not other HCAs (MeIQx, 1.12, 0.98-1.27, PhIP from white meat, 1.07, 0.94-1.21, DiMeIQx, 1.09, 0.97-1.21) or MDM (1.13, 1.00-1.28). For high grade (Gleason sum 7 with pattern 4+3 and Gleason sum 8-10, n=483 cases) and advanced cancers (n=281), we only observed positive associations for PhIP from red meat (top vs. bottom quintile: high grade: HR=1.44, 95% CI 1.04-1.98, p-trend=0.03; advanced: HR=1.50, 95% 0.99-2.26; p-trend=0.12), but associations for advanced cancers did not reach statistical significance. Observed associations remained similar after adjustment for total, unprocessed or processed red meat intake.

Conclusion. Observed positive associations between PhIP intake from red meat and prostate cancer, particularly high-grade and possibly also advanced prostate cancer need to be confirmed in other studies.

Impact. Results do not provide strong evidence that HCAs increase risk of prostate cancers.

Introduction

Higher meat, especially red meat consumption has been suggested to be associated with higher risk of prostate cancer, but the evidence is not consistent (reviewed in (1)).

One possible mechanism through which red meat may be involved in prostate carcinogenesis is the formation of mutagenic heterocyclic amines (HCA) in muscle meat when meats are cooked at high temperature and for long duration (e.g. grilling, barbecuing, frying, and broiling) (2).

HCAs were first detected in the 1970s in smoke condensate of grilled fish and were shown to be mutagenic (3). HCAs are formed from precursors (creatine/creatinine, sugar, amino acids) found in the muscle of meat and fish cooked at temperatures exceeding 130°C (4, 5). HCA production and amount depend mainly on cooking method, temperature, and the type of meat or fish (4). Meat drippings and gravy made from these drippings contain considerable amounts of HCA (6). The most abundant HCAs in the human diet are 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP), 2-amino-3,8-dimethylimidazo[4,5-f]quinoxaline (MeIQx) and 2-amino-3,4,8-trimethylimidazo[4,5-f]quinoxaline (DiMeIQx) (7).

Thus far, five cohort studies have examined the association between HCA intake and prostate cancer risk. Some reported positive associations for red meat that was consumed mostly well-done (8-10) or for HCA intake (10), in particular with advanced disease. Two found no associations (11, 12). In addition, three (13-15) of five (16, 17) case-control studies reported no clear associations between HCA intake and risk of prostate cancer.

The aim of this study was to examine in the PSA era the association of HCA intake and a meat-derived mutagenicity (MDM) index with risk of prostate cancer in a large prospective US cohort study.

Materials and Methods

Study population

The Health Professionals Follow-up Study includes 51,529 dentists, veterinarians, pharmacists, optometrists, osteopathic physicians, and podiatrists who were between 40 and 75 years of age at enrollment in 1986. At baseline, all participants completed a 131-item semi-quantitative food-frequency questionnaire (FFQ) and provided information on age, race or ethnicity, weight, height, physical activity, cigarette smoking, alcohol consumption, and medical history. Every two years questionnaires were mailed to collect updated information on exposure and new diagnoses; follow-up FFQs were mailed every four years. The validity and reproducibility of our FFQs have been documented previously (18). Deaths were reported by next-of-kin, the postal service, and searches of the National Death Index. The study was approved by the Human Subjects Committee at the Harvard School of Public Health.

In 1996, questions on meat preparation (cooking method and degree of doneness) were included on the biennial questionnaire. This analysis is based on the cohort of men who provided information on meat preparation in 1996 and who responded to the 1994 FFQ. Of these participants, we excluded all men with a cancer diagnosis except for non-melanoma skin cancer prior to 1996 and men with missing information on meat

preparation (no answer to any of the cooking method questions, missing frequency of meat intake of at least one cooked meat item, or no information on bacon consumption, which is needed to estimate HCA intake). The final cohort for this analysis included 23,030 men.

Case ascertainment

On the biennial follow-up questionnaires, men were asked to report whether they were diagnosed with prostate cancer, in the previous two years. Diagnosis was confirmed by review of medical records (>90% of the cases). For non-responders, we used information from the National Death Index, postal service and next-of-kin to determine whether a participant had died. After the review of death certificates, informed consent was obtained from next-of-kin of participants who died of prostate cancer to obtain medical records. Study investigators reviewed the medical records to confirm a prostate cancer diagnosis and prostate cancer deaths.

This analysis was based on 2770 non-T1a prostate cancer cases ascertained after the date of return of the 1996 questionnaire through January 31st, 2010. Of these cancer cases, 281 were classified as advanced or lethal (diagnosed at an advanced stage [T3b, T4, N1, and M1], developed metastases during follow-up, or died due to prostate cancer). We confirmed 2285 cases as organ-confined or having limited extraprostatic extension (T1b, T1c, T2, T3a and N0 or Nx and M0), and the remainder could not be assigned a stage. We defined low-grade prostate cancer as patients having Gleason sum of 2-6 or 7 with a pattern of 3 plus 4 (n=1859) and high-grade cases as those with Gleason sum 7 and pattern of 4 plus 3 and those with Gleason sum 8-10 (n=483).

Assessment of meat consumption and cooking methods

Total, red (processed and unprocessed), and white meat intake was estimated from the 1994 FFQ. For each food item, a commonly used unit or serving size was specified.

The participants were asked to indicate how often, on average, they consumed each food item, with nine possible response categories ranging from 'never' to 'six or more times per day'. In 1996, participants completed a cooking methods questionnaire based on the results of a pilot study, which ascertained a group of cooking method questions that best predicted HCA intake in the HPFS (7). Information on cooking methods and typical outside appearance of pan-fried, broiled, and grilled or barbecued chicken, hamburger, and steak; fried, microwaved, and broiled bacon; fried sausage; roast beef; and homemade gravy were assessed (7). Based on results of the pilot study, variation in doneness level of fried bacon was small. Therefore, fried bacon intake was based on the 1994 FFQ and assumed to be medium browned (7). The Charred Database (19) contains data on HCA and MDM concentration in various meat items depending on cooking method and outside appearance (20). The mutagenic activity of meat samples was assessed by the Ames/Salmonella test. The resulting MDM is a measure of the total mutagenic activity found in cooked meats, which integrates mutagenic activity from all classes of mutagens, including heterocyclic amines, but also benzo[a]pyrene or yet unidentified compounds, found in cooked meats (21).

A participant's HCA intake or MDM was calculated by multiplying the given consumption frequency of each food by its HCA content or MDM value for the specified cooking method and degree of browning derived from the Charred Database (20).

Statistical analysis

Intake of PhIP, MeIQx, DiMeIQx, and MDM index was categorized into quintiles based on the distribution in the analytic cohort. We calculated hazard ratios (HR) and corresponding 95% confidence intervals (CI) using Cox proportional hazards regression to evaluate the association of HCA intake and MDM index with total and advanced or lethal prostate cancer. We observed no violation of the proportional hazards assumption. The models were adjusted for age, race/ethnicity, pack-years smoked in the past ten years, family history of prostate cancer, updated history of diabetes mellitus, height, body mass index at age 21, updated vigorous leisure-time physical activity, updated current aspirin use, cumulative average updated intake of tomato products and fish, and updated intake of energy, alcohol, vitamin E, α -linolenic acid, and total calcium. We decided on this set of confounders a priori based on the overall literature. Cumulative updated average intake represents the average intake of a food or nutrient from all available FFQs up to the start of each follow-up interval (22). To test whether associations between HCA intake or MDM index might be explained by meat intake *per se*, we ran models that additionally adjusted for (a) total red meat, (b) unprocessed red meat (which includes regular hamburger; lean or extra-lean hamburger; beef, pork, or lamb as a sandwich or mixed dish (e.g., stew, casserole, lasagna, etc.); beef or lamb as a main dish (e.g., steak, roast, ham, etc.); pork as a main dish (e.g., ham or chops)), (c) processed red meat salami, bologna, or other processed meat sandwiches; other processed meats (e.g., sausage, kielbasa, etc.); bacon; hot dogs), and (d) white meat (chicken and turkey). We also examined the association of total red meat, unprocessed red meat, processed red meat and white

meat consumption in 1994 (closest date to HCA assessment in 1996) and the risk of prostate cancer. To test for trend, we entered the median of each quintile of HCA intake or MDM index as a single continuous variable into the model and used the Wald test to assess statistical significance of the coefficient. We ran stratified models to determine if the associations of HCA intake or MDM index with prostate cancer varied by age (<60 vs. \geq 60 years in 1996), current body mass index, aspirin use (non-users in 1996 vs. user; infrequent vs. frequent user; never user vs. user), consumption of tomato products (sum of tomato sauce and pizza; < vs. \geq 2.0 servings/week), intake of vitamin E (< vs. \geq median intake), and history of prostatitis. In a sensitivity analysis, we restricted the analysis to men who have had at least one PSA test until the end of the observation period in 2010. The presence of multiplicative interaction was assessed by including a cross-product term for these factors and HCA intake or MDM index in the regression model along with the main effect terms. The statistical significance of the coefficient for the cross-product term was evaluated by the Wald test. A p-value of <0.05 was considered statistically significant.

Results

The analysis included 2770 prostate cancer cases in 314,746 person-years of follow-up. Selected baseline characteristics of the study participants by highest and lowest category of PhIP, MeIQx, and DiMeIQx intake are shown in Table 1. Participants in the highest categories engaged less often in vigorous physical activity and were more likely to have smoked in the past 10 years. These participants also had a higher intake of red meat, but lower intake of vitamin E than participants in the lowest quintiles.

Intake of PhIP from red meat was statistically significantly associated with total prostate cancer risk (multivariable HR=1.18, 95% CI 1.03-1.35, top vs. bottom quintile; Table 2). However, intake of PhIP from white meat or intake of MeIQx, DiMeIQx or the MDM index were not associated with total prostate cancer risk (Table 2). For high grade cancers (Gleason sum 7 with pattern 4+3 and Gleason sum 8-10, n=483 cases), we also observed an increased risk among participants with high intake of PhIP from red meat (HR=1.44, 95% CI 1.04-1.98, p-trend=0.03), but we did not observe any significant associations with MDM index (HR=1.05, 95% CI 0.78-1.41, p-trend=0.54), MeIQx, PhIP from white meat or DiMeIQx.

For advanced cancers (n=281) we only observed positive associations for PhIP from red meat (top vs. bottom quintile: HR: 1.50 (95% 0.99-2.26; p-trend=0.12), but associations for advanced cancers did not reach statistical significance (Table 2). HCA intake or MDM index were not statistically significantly associated with the risk of organ-confined or low-grade prostate cancer (data not shown).

Adding total red meat intake to the multivariable model did not materially change the associations between HCA intake or MDM index with total, advanced and high-grade prostate (data not shown); results were also similar after adding unprocessed red meat or processed meat separately to the models (data not shown). Similarly, no major changes were observed when adding white meat to the model. In order to evaluate whether the effect on high-grade and advanced tumors was due to HCA or other carcinogens in cooked meat, we included both MDM and PhIP from red meat into the same model; but the risk estimates for PhIP from red meat were similar (top vs. bottom

quintile, HR, 95% CI; high grade: PhIP from red meat 1.50; 1.06-2.14; MDM 0.90, 0.65-1.23; advanced: PhIP from red meat 1.45, 0.92-2.28; MDM 1.17, 0.76-1.81).

Generally, the associations of HCA intake or MDM index with total prostate cancer were not modified by age, BMI, aspirin use, vitamin E intake, or history of prostatitis (all p-interaction > 0.05; data not shown). However, we observed an increased risk of total prostate cancer associated with high HCA intake or MDM index among those with lower (but not higher) consumption of tomato products; however interactions were not statistically significant, except for MDM (p-interaction=0.03; Table 3). Excluding men who had never had a PSA test did not change our results appreciably (data not shown). In joint analysis we observed that men with both low tomato consumption and high MDM had a 22% (95% CI 2%-47%) higher risk of developing prostate cancer when compared to those with high tomato consumption and low MDM (data not shown).

Discussion

In this large prospective cohort study conducted in the PSA era, we did not observe statistically significant associations between HCA intake and risk of prostate cancer. However, we observed positive associations between PhIP intake from red meat and risk of total, high grade and advanced prostate cancer. Also, we noted that men with low tomato product consumption had an increased risk of total prostate cancer associated with intake of some HCAs or the MDM index, whereas no increased risks were observed among men with high tomato product consumption.

Heterocyclic amines have consistently been shown to be mutagenic in mutagenicity assays and carcinogenic in animal models (23, 24). PhIP contributes most to HCA intake in the US, whereas DiMeIQx intake is comparably low, but is the most mutagenic HCA of the three examined HCAs (25). Epidemiologic evidence for an association between HCA intake and prostate cancer, however, is limited. In the most recent study, conducted among participants of the California Collaborative Prostate Cancer Study (16), high consumption of red meat cooked at high temperature and well done red meat were positively associated with risk of advanced, but not overall prostate cancer. Intake of PhIP was also positively, although not statistically significantly related with advanced prostate cancer. This result is similar to what has been reported in the NIH-AARP cohort (8). High intake of very well done meat and of PhIP were associated with an increased risk of prostate cancer in the Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial (10). In the Agricultural Health Study (9), consumption of well or very well done meat was related to an increased risk of total and in particular advanced prostate cancer, and there were non-statistically significant positive associations with DiMeIQx and MeIQx intakes. In the Multiethnic Cohort (11) and the EPIC-Heidelberg cohort (12), no associations of HCA intake or type of meat cooking or degree of doneness with risk of prostate cancer overall or with advanced disease were observed. In two case-control studies HCA intake was not clearly associated with prostate cancer risk (13-15), although another study by John et al. (13) reported a positive association of well done and grilled red meat consumption with risk of advanced prostate cancer. A fourth case-control study including 470 cases of aggressive prostate cancer related high intake of well done meat and intake of MeIQx and DiMeIQx to an increased risk (17).

It is of interest to note that we observed associations between PhIP intake from red but not from white meat with risk of high-grade prostate cancer, although PhIP intake from white meat is twice as high as the intake from red meat. In addition, associations for PhIP from red meat remained similar after we adjusted for total, unprocessed and processed meat intake separately. This argues against a causal association between PhIP intake (or HCA intake in general) and prostate cancer risk, and suggest the possibility that not HCA themselves but other mutagenic compounds that arise from cooking of red meat may also be a risk factor for prostate cancer. MDM integrates mutagenic activity of different compounds in cooked meats such as heterocyclic amines or benzo[a]pyrene, but also yet unidentified compounds (21). However, when we examined associations between PhIP from red meat and MDM and high grade or advanced cancers in the same model estimates for PhIP from red meat were similar to estimates without MDM in the model.

In the rat, PhIP induced mutations in the ventral prostate, which was associated with an infiltrate of inflammatory cells (26). Similarly, PhIP induced prostate cancer in a CYP1A-humanized mice model, which is preceded by inflammatory proliferative epithelial lesions (27). In our analysis, we did not observe evidence for effect modification of the association between HCA intake or MDM index and prostate cancer risk by use of aspirin, intake of dietary vitamin E, or BMI, all of which are factors that influence inflammatory processes. We did, however, observe that associations were modified by consumption of tomato products. Lycopene is a potent antioxidant found in tomatoes (28), but the mechanism by which lycopene may modify the association of HCA intake and prostate cancer is currently unclear. Animal studies did not provide clear evidence

that administration of lycopene reduced formation of HCA-DNA adducts, reduced oxidative stress induced by HCA administration, or influenced levels of phase-II enzymes (29, 30). However, lycopene may affect cell cycle, apoptosis, inflammation, and angiogenesis by a variety of pathways (31, 32). To the best of our knowledge, this is the first study to observe a modifying effect of tomato intake on risk of prostate cancer and, thus, our results need to be confirmed by other studies. However, we cannot exclude the possibility that our results are due to chance and, thus, warrant confirmation from other studies.

We observed positive associations of PhIP intake from red meat with total, high grade and advanced prostate cancer; even though associations for advanced cancers did not reach statistical significance. However, the number of advanced or lethal cases in our analytical cohort was small because we began follow-up in 1996 after the completion of the questionnaire, i.e., in the PSA era; about 95% of men included in our analysis had reported at least one PSA between 1996 until the end of the follow-up period. Other studies that had reported associations of HCA intake with advanced prostate cancer (9, 13, 16, 17) did not separately evaluate high-grade disease.

We assessed meat cooking habits in 1996 and used that information throughout follow-up. It is possible that participants might have changed their meat preparation habits or overall meat intake during follow-up, which may result in another source of measurement error. For example, red meat consumption decreased in the HPFS over time. In 2004, we assessed meat cooking methods a second time in the HPFS cohort and correlations between calculated HCA intake in 1996 and 2004 were $r=0.31$, 0.40 ,

and 0.23 (all p-values < 0.0001) for PhIP, MeIQx, and DiMeIQx, respectively. HCA intake estimations suffer from imprecision. Assessing a person's dietary intake by means of a FFQ is prone to recall bias due to over- or underestimation of dietary intake by the study participant leading to misclassification with respect to dietary intake. We estimated HCA intake and MDM index based on participants' responses to questions on the cooking method of specific meat items. Doneness levels specified by participants are proxies of the actual intake of heterocyclic amines, which varies by duration of cooking, temperatures used (33), type of oil used (34, 35), handling of the meat before and during the cooking process (36) as well as of the cooking method itself (37) such that cooking short time by high temperature and long time by lower temperature can result in the same degree of browning but not in the same degree of doneness. In addition, the use of limited data on the HCA content in differently prepared meats for the computation of HCA intake is another major shortcoming of this approach to quantify intake (38). However, the cooking questions used in this study are based on a pilot study (7) that determined which set of questions best predicted HCA intake in our cohorts. Also, because data on cooking methods were obtained prior to cancer diagnosis, possible measurement errors due to the aforementioned factors should be non-differential, which tends to attenuate associations.

In conclusion, our results do not provide strong evidence that HCAs in general increase risk of prostate cancers. Observed associations between PhIP intake from red meat and prostate cancer, particularly high-grade and possibly also advanced prostate cancer need to be confirmed in other studies.

Acknowledgements

We would like to thank the participants and staff of the Health Professionals Follow-up Study for their valuable contributions as well as the following state cancer registries for their help: AL, AZ, AR, CA, CO, CT, DE, FL, GA, ID, IL, IN, IA, KY, LA, ME, MD, MA, MI, NE, NH, NJ, NY, NC, ND, OH, OK, OR, PA, RI, SC, TN, TX, VA, WA, WY. The authors assume full responsibility for analyses and interpretation of these data.

References

1. Alexander DD, Mink PJ, Cushing CA, Scurman B. A review and meta-analysis of prospective studies of red and processed meat intake and prostate cancer. *Nutr J* 2010; 9:50.
2. Michaud DS, Augustsson K, Rimm EB, Stampfer MJ, Willet WC, Giovannucci E. A prospective study on intake of animal products and risk of prostate cancer. *Cancer Causes Control* 2001; 12:557-67.
3. Sugimura T. Overview of carcinogenic heterocyclic amines. *Mutat Res* 1997; 376:211-9.
4. Skog KI, Johansson MA, Jagerstad MI. Carcinogenic heterocyclic amines in model systems and cooked foods: a review on formation, occurrence and intake. *Food Chem Toxicol* 1998; 36:879-96.
5. Jagerstad M, Skog K, Grivas S, Olsson K. Formation of heterocyclic amines using model systems. *Mutat Res* 1991; 259:219-33.
6. Gross GA, Gruter A. Quantitation of mutagenic/carcinogenic heterocyclic aromatic amines in food products. *J Chromatogr* 1992; 592:271-8.
7. Byrne C, Sinha R, Platz EA, Giovannucci E, Colditz GA, Hunter DJ, et al. Predictors of dietary heterocyclic amine intake in three prospective cohorts. *Cancer Epidemiol Biomarkers Prev* 1998; 7:523-9.
8. Sinha R, Park Y, Graubard BI, Leitzmann MF, Hollenbeck A, Schatzkin A, et al. Meat and meat-related compounds and risk of prostate cancer in a large prospective cohort study in the United States. *Am J Epidemiol* 2009; 170:1165-77.
9. Koutros S, Cross AJ, Sandler DP, Hoppin JA, Ma X, Zheng T, et al. Meat and meat mutagens and risk of prostate cancer in the Agricultural Health Study. *Cancer Epidemiol Biomarkers Prev* 2008; 17:80-7.
10. Cross AJ, Peters U, Kirsh VA, Andriole GL, Reding D, Hayes RB, et al. A Prospective Study of Meat and Meat Mutagens and Prostate Cancer Risk. *Cancer Res* 2005; 65:11779-84.
11. Sharma S, Cao X, Wilkens LR, Yamamoto J, Lum-Jones A, Henderson BE, et al. Well-done meat consumption, NAT1 and NAT2 acetylase genotypes and prostate cancer risk: the multiethnic cohort study. *Cancer Epidemiol Biomarkers Prev* 2010; 19:1866-70.
12. Sander A, Linseisen J, Rohrmann S. Intake of heterocyclic aromatic amines and the risk of prostate cancer in the EPIC-Heidelberg cohort. *Cancer Causes Control* 2010; 22:109-14.
13. John EM, Stern MC, Sinha R, Koo J. Meat consumption, cooking practices, meat mutagens, and risk of prostate cancer. *Nutr Cancer* 2011; 63:525-37.
14. Rovito PM, Jr, Morse PD, Spinek K, Newman N, Jones RF, Wang CY, et al. Heterocyclic amines and genotype of N-acetyltransferases as risk factors for prostate cancer. 2005; 8:69-74.
15. Norrish AE, Ferguson LR, Knize MG, Felton JS, Sharpe SJ, Jackson RT. Heterocyclic amine content of cooked meat and risk of prostate cancer. *J Natl Cancer Inst* 1999; 91:2038-44.
16. Joshi AD, Corral R, Catsburg C, Lewinger JP, Koo J, John EM, et al. Red meat and poultry, cooking practices, genetic susceptibility and risk of prostate cancer: results from a multiethnic case-control study. *Carcinogenesis* 2012; 33:2108-18.
17. Punnen S, Hardin J, Cheng I, Klein EA, Witte JS. Impact of meat consumption, preparation, and mutagens on aggressive prostate cancer. *PLoS One* 2011; 6:e27711.
18. Rimm EB, Giovannucci EL, Stampfer MJ, Colditz GA, Litin LB, Willett WC. Reproducibility and validity of an expanded self-administered semiquantitative food frequency questionnaire among male health professionals. *Am J Epidemiol* 1992; 135:1114-26; discussion 27-36.
19. National Cancer Institute. CHARRED: computerized heterocyclic amines database resource for research in the epidemiologic of disease. 2006 [cited 2015 May 7]; Available from: <http://dceg.cancer.gov/tools/design/charred>

20. Sinha R, Cross A, Curtin J, Zimmerman T, McNutt S, Risch A, et al. Development of a food frequency questionnaire module and databases for compounds in cooked and processed meats. *Molecular Nutrition & Food Research* 2005; 49:648-55.
21. Sinha R, Kulldorff M, Chow WH, Denobile J, Rothman N. Dietary intake of heterocyclic amines, meat-derived mutagenic activity, and risk of colorectal adenomas. *Cancer Epidemiol Biomarkers Prev* 2001; 10:559-62.
22. Willett WC. *Nutritional Epidemiology*. Second ed. New York: Oxford University Press; 1998.
23. Shirai T, Tamano S, Sano M, Masui T, Hasegawa R, Ito N. Carcinogenicity of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) in rats: dose-response studies. *Princess Takamatsu Symp* 1995; 23:232-9.
24. Snyderwine EG, Turesky RJ, Turteltaub KW, Davis CD, Sadrieh N, Schut HA, et al. Metabolism of food-derived heterocyclic amines in nonhuman primates. *Mutat Res* 1997; 376:203-10.
25. Wang CY, Debiec-Rychter M, Schut HAJ, Morse P, Jones RF, Archer C, et al. N-Acetyltransferase expression and DNA binding of N-hydroxyheterocyclic amines in human prostate epithelium. *Carcinogenesis* 1999; 20:1591-6.
26. Nakai Y, Nelson WG, De Marzo AM. The dietary charred meat carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine acts as both a tumor Initiator and promoter in the rat ventral prostate. *Cancer Res* 2007; 67:1378-84.
27. Li G, Wang H, Liu AB, Cheung C, Reuhl KR, Bosland MC, et al. Dietary carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine-induced prostate carcinogenesis in CYP1A-humanized mice. *Cancer Prev Res (Phila)* 2012; 5:963-72.
28. Stahl W, Sies H. Lycopene: a biologically important carotenoid for humans? *Arch Biochem Biophys* 1996; 336:1-9.
29. Breinholt V, Lauridsen ST, Daneshvar B, Jakobsen J. Dose-response effects of lycopene on selected drug-metabolizing and antioxidant enzymes in the rat. *Cancer Lett* 2000; 154:201-10.
30. Dingley KH, Ubick EA, Chiarappa-Zucca ML, Nowell S, Abel S, Ebeler SE, et al. Effect of dietary constituents with chemopreventive potential on adduct formation of a low dose of the heterocyclic amines PhIP and IQ and phase II hepatic enzymes. *Nutr Cancer* 2003; 46:212-21.
31. Trejo-Solis C, Pedraza-Chaverri J, Torres-Ramos M, Jimenez-Farfan D, Cruz Salgado A, Serrano-Garcia N, et al. Multiple molecular and cellular mechanisms of action of lycopene in cancer inhibition. *Evid Based Complement Alternat Med* 2013; 2013:705121.
32. Holzapfel NP, Holzapfel BM, Champ S, Feldthusen J, Clements J, Huttmacher DW. The potential role of lycopene for the prevention and therapy of prostate cancer: from molecular mechanisms to clinical evidence. *Int J Mol Sci* 2013; 14:14620-46.
33. Skog K, Steineck G, Augustsson K, Jagerstad M. Effect of cooking temperature on the formation of heterocyclic amines in fried meat products and pan residues. *Carcinogenesis* 1995; 16:861-7.
34. Johansson M, Skog K, Jagerstad M. Effects of edible oils and fatty acids on the formation of mutagenic heterocyclic amines in a model system. *Carcinogenesis* 1993; 14:89-94.
35. Persson E, Graziani G, Ferracane R, Fogliano V, Skog K. Influence of antioxidants in virgin olive oil on the formation of heterocyclic amines in fried beefburgers. *Food Chem Toxicol* 2003; 41:1587-97.
36. Salmon CP, Knize MG, Felton JS. Effects of marinating on heterocyclic amine carcinogen formation in grilled chicken. *Food Chem Toxicol* 1997; 35:433-41.
37. Skog K. Problems associated with the determination of heterocyclic amines in cooked foods and human exposure. *Food and Chemical Toxicology* 2002; 40:1197-203.
38. Rohrmann S, Zoller D, Hermann S, Linseisen J. Intake of heterocyclic aromatic amines from meat in the EPIC-Heidelberg cohort. *Brit J Nutr* 2007; 98:1112-5.

Table 1. Baseline (1996) age-standardized characteristics by lowest and highest quintiles of HCA and MDM activity intake, Health Professionals

Follow-up Study

| Quintile (median intake) | Total MeIQx (ng/d) | | Total DiMeIQx (ng/d) | | Total PhIP (ng/d) | | MDM index (revertant colonies/d) | |
|---|--------------------|-------------|----------------------|-------------|-------------------|-------------|-------------------------------------|-------------|
| | Q1 (1.6) | Q5 (37.8) | Q1 (0) | Q5 (4.0) | Q1 (13.2) | Q5 (218.1) | Q1 (693) | Q5 (8233) |
| N | 531 | 558 | 869 | 562 | 523 | 568 | 520 | 577 |
| Age in 1996 (mean, SD) | 61.4 (8.3) | 59.7 (7.9) | 61.7 (8.3) | 58.5 (7.4) | 62.8 (8.5) | 58.2 (7.3) | 62.0 (8.4) | 58.8 (7.6) |
| Caucasian (%) | 90 | 93 | 92 | 93 | 91 | 93 | 91 | 93 |
| Smoked in past ten years (%) | 12 | 18 | 15 | 17 | 14 | 16 | 13 | 17 |
| Family history of prostate cancer (%) | 12 | 15 | 13 | 15 | 13 | 14 | 12 | 14 |
| History of diabetes (up to 1996; %) | 3 | 5 | 4 | 4 | 3 | 4 | 3 | 5 |
| Current use of aspirin in 1996 (%) | 70 | 76 | 72 | 76 | 70 | 76 | 70 | 76 |
| Screening PSA test by 2006 (%) | 96 | 93 | 95 | 95 | 94 | 95 | 95 | 95 |
| Body height (inches), mean (SD) | 70.0 (2.6) | 70.5 (2.5) | 70.2 (2.7) | 70.4 (2.5) | 70.2 (2.6) | 70.3 (2.5) | 70.1 (2.7) | 70.4 (2.6) |
| BMI at age 21 (kg/m ²), mean (SD) | 23.0 (2.6) | 23.0 (2.7) | 23.0 (2.6) | 23.1 (2.7) | 22.9 (2.6) | 23.1 (2.6) | 23.0 (2.7) | 23.0 (2.7) |
| BMI in 1996 (kg/m ²), mean (SD) | 25.0 (3.0) | 26.5 (3.5) | 25.6 (3.3) | 26.1 (3.4) | 25.1 (3.2) | 26.3 (3.3) | 25.1 (3.2) | 26.2 (3.5) |
| Vigorous MET-h/wk, mean (SD) | 20.7 (29.6) | 11.9 (23.3) | 16.8 (28.1) | 14.9 (24.3) | 17.8 (29.1) | 15.1 (25.7) | 18.5 (29.2) | 14.3 (24.5) |
| Daily intake in 1994, mean (SD) | | | | | | | | |
| Alcohol intake (drinks/w) | 4.57 (6.00) | 6.23 (7.71) | 5.25 (6.67) | 6.27 (7.27) | 4.50 (6.27) | 6.91 (7.78) | 4.77 (6.28) | 6.27 (7.52) |
| Tomato products (servings/w) | 1.55 (1.29) | 1.73 (1.29) | 1.59 (1.26) | 1.75 (1.26) | 1.55 (1.29) | 1.75 (1.27) | 1.53 (1.25) | 1.75 (1.31) |
| Red meat (servings/w) | 3.65 (3.19) | 10.9 (5.38) | 5.98 (4.52) | 8.11 (5.23) | 4.88 (4.13) | 8.47 (5.19) | 4.46 (3.63) | 8.98 (5.36) |
| Fish (servings/w) | 2.57 (1.91) | 1.78 (1.34) | 2.32 (1.80) | 2.21 (1.65) | 2.12 (1.71) | 2.32 (1.66) | 2.19 (1.73) | 2.22 (1.67) |
| White meat (servings/w) | 2.68 (1.68) | 2.66 (1.59) | 2.69 (1.66) | 2.96 (1.58) | 2.25 (1.53) | 3.21 (1.67) | 2.28 (1.52) | 3.13 (1.66) |
| Calcium (mg/d) | 981 (441) | 843 (343) | 943 (423) | 875 (361) | 990 (448) | 847 (355) | 972 (441) | 864 (358) |
| Vitamin E (mg/d) | 108 (129) | 64.5 (97.3) | 93.6 (122) | 77.1 (109) | 97.6 (126) | 78.9 (111) | 97.5 (126) | 76.2 (107) |
| Alpha-linolenic acid (mg/d) | 1129 (359) | 1155 (361) | 1118 (348) | 1133 (369) | 1139 (359) | 1137 (344) | 1134 (364) | 1148 (347) |

Table 2. Association of heterocyclic amine intake and MDM index with prostate cancer, Health Professionals
 Follow-up Study 1996-2010

| | Total prostate cancer (n=2770) | | | | High-grade prostate cancer (n=483) ¹ | | | | Advanced or lethal prostate cancer (n=281) ² | | | |
|------------------------------------|--------------------------------|-----------------|-----------------|------------|---|-----------------|-----------------|------------|---|-----------------|-----------------|------------|
| | N | HR ³ | HR ⁴ | 95% CI | N | HR ³ | HR ⁴ | 95% CI | N | HR ³ | HR ⁴ | 95% CI |
| MelQx (ng/d) | | | | | | | | | | | | |
| <3.6 | 531 | 1.00 | 1.00 | ref. | 92 | 1.00 | 1.00 | ref. | 60 | 1.00 | 1.00 | ref. |
| 3.7 - <8.0 | 546 | 1.03 | 1.04 | 0.92 -1.17 | 97 | 1.12 | 1.12 | 0.83 -1.49 | 45 | 0.85 | 0.83 | 0.56 -1.24 |
| 8.1 - <14.1 | 581 | 1.08 | 1.12 | 0.99 -1.27 | 102 | 1.19 | 1.17 | 0.87 -1.57 | 66 | 1.24 | 1.24 | 0.86 -1.79 |
| 14.1 - <25.0 | 554 | 1.10 | 1.08 | 0.95 -1.22 | 94 | 1.11 | 1.09 | 0.80 -1.48 | 54 | 1.03 | 1.00 | 0.67 -1.47 |
| ≥25.0 | 558 | 1.11 | 1.12 | 0.98 -1.27 | 98 | 1.17 | 1.15 | 0.84 -1.56 | 56 | 1.11 | 1.08 | 0.72 -1.60 |
| <i>p-trend</i> | | 0.11 | 0.07 | | | 0.35 | 0.49 | | | 0.38 | 0.52 | |
| PhIP (ng/d) | | | | | | | | | | | | |
| <25.6 | 523 | 1.00 | 1.00 | ref. | 104 | 1.00 | 1.00 | ref. | 66 | 1.00 | 1.00 | ref. |
| 25.6 - <51.4 | 571 | 1.08 | 1.08 | 0.96 -1.22 | 103 | 1.02 | 1.00 | 0.76 -1.33 | 51 | 0.84 | 0.84 | 0.58 -1.22 |
| 51.4 - <89.4 | 553 | 1.04 | 1.02 | 0.90 -1.15 | 80 | 0.80 | 0.77 | 0.57 -1.04 | 63 | 1.06 | 1.06 | 0.74 -1.51 |
| 89.4 - <152.9 | 555 | 1.06 | 1.06 | 0.94 -1.20 | 92 | 0.96 | 0.95 | 0.71 -1.27 | 52 | 0.97 | 0.98 | 0.67 -1.43 |
| ≥152.9 | 568 | 1.09 | 1.08 | 0.95 -1.22 | 104 | 1.13 | 1.08 | 0.81 -1.44 | 49 | 1.00 | 1.01 | 0.68 -1.49 |
| <i>p-trend</i> | | 0.29 | 0.35 | | | 0.60 | 0.79 | | | 0.77 | 0.73 | |
| PhIP from red meat (ng/d) | | | | | | | | | | | | |
| <0.7 | 441 | 1.00 | 1.00 | ref. | 50 | 1.00 | 1.00 | ref. | 51 | 1.00 | 1.00 | ref. |
| 0.7 - <9.5 | 597 | 1.05 | 1.05 | 0.92 -1.19 | 100 | 1.02 | 1.02 | 0.75 -1.37 | 63 | 1.05 | 1.05 | 0.71 -1.54 |
| 9.5 - <18.4 | 571 | 1.11 | 1.11 | 0.97 -1.26 | 90 | 1.04 | 1.05 | 0.77 -1.44 | 52 | 0.99 | 1.00 | 0.67 -1.50 |
| 18.4 - <45.0 | 638 | 1.05 | 1.04 | 0.91 -1.18 | 107 | 1.07 | 1.06 | 0.78 -1.45 | 56 | 0.98 | 0.98 | 0.65 -1.47 |
| ≥45.0 | 523 | 1.18 | 1.18 | 1.03 -1.35 | 106 | 1.47 | 1.44 | 1.04 -1.98 | 59 | 1.48 | 1.50 | 0.99 -2.26 |
| <i>p-trend</i> | | 0.03 | 0.05 | | | 0.02 | 0.03 | | | 0.12 | 0.12 | |
| PhIP from white meat (ng/d) | | | | | | | | | | | | |
| <14.4 | 510 | 1.00 | 1.00 | ref. | 94 | 1.00 | 1.00 | ref. | 60 | 1.00 | 1.00 | ref. |
| 14.4 - <29.9 | 578 | 1.09 | 1.08 | 0.96 -1.22 | 107 | 1.11 | 1.09 | 0.82 -1.45 | 65 | 1.16 | 1.18 | 0.82 -1.69 |
| 29.9 - <56.3 | 580 | 1.10 | 1.10 | 0.97 -1.24 | 96 | 1.04 | 1.03 | 0.77 -1.37 | 51 | 1.00 | 1.00 | 0.68 -1.47 |
| 56.3 - <110.4 | 547 | 1.05 | 1.05 | 0.92 -1.18 | 98 | 1.06 | 1.06 | 0.79 -1.42 | 59 | 1.10 | 1.12 | 0.77 -1.63 |
| ≥110.4 | 555 | 1.07 | 1.07 | 0.94 -1.21 | 88 | 1.02 | 0.99 | 0.73 -1.34 | 46 | 0.99 | 1.02 | 0.68 -1.53 |
| <i>p-trend</i> | | 0.49 | 0.51 | | | 0.98 | 0.87 | | | 0.92 | 0.99 | |
| DiMelQx (ng/d) | | | | | | | | | | | | |
| <0.10 | 869 | 1.00 | 1.00 | ref. | 164 | 1.00 | 1.00 | ref. | 86 | 1.00 | 1.00 | ref. |
| 0.10 - <0.20 | 272 | 1.15 | 1.14 | 0.99 -1.31 | 49 | 1.14 | 1.13 | 0.82 -1.56 | 34 | 1.64 | 1.67 | 1.11 -2.51 |
| 0.20 - <0.87 | 539 | 1.12 | 1.11 | 1.00 -1.24 | 85 | 0.95 | 0.94 | 0.72 -1.23 | 48 | 1.10 | 1.10 | 0.77 -1.58 |
| 0.87 - <1.90 | 528 | 1.00 | 1.00 | 0.89 -1.11 | 91 | 0.92 | 0.90 | 0.69 -1.16 | 60 | 1.28 | 1.29 | 0.92 -1.81 |
| ≥1.90 | 562 | 1.09 | 1.09 | 0.97 -1.21 | 94 | 1.02 | 1.00 | 0.77 -1.30 | 53 | 1.26 | 1.28 | 0.90 -1.83 |
| <i>p-trend</i> | | 0.32 | 0.37 | | | 0.74 | 0.61 | | | 0.23 | 0.21 | |
| MDM index (revertant colonies/day) | | | | | | | | | | | | |
| < 1222 | 520 | 1.00 | 1.00 | ref. | 98 | 1.00 | 1.00 | ref. | 54 | 1.00 | 1.00 | ref. |
| 1222- < 2256 | 575 | 1.09 | 1.09 | 0.96 -1.23 | 102 | 1.04 | 1.01 | 0.76 -1.34 | 64 | 1.32 | 1.31 | 0.90 -1.90 |
| 2256 - < 3551 | 531 | 1.02 | 1.03 | 0.91 -1.16 | 80 | 0.85 | 0.83 | 0.61 -1.13 | 50 | 1.04 | 1.03 | 0.69 -1.53 |
| 3551 - < 5663 | 567 | 1.10 | 1.11 | 0.98 -1.25 | 105 | 1.14 | 1.12 | 0.84 -1.49 | 57 | 1.25 | 1.25 | 0.85 -1.84 |
| ≥ 5663 | 577 | 1.12 | 1.13 | 1.00 -1.28 | 98 | 1.10 | 1.05 | 0.78 -1.41 | 56 | 1.32 | 1.30 | 0.88 -1.93 |
| <i>p-trend</i> | | 0.08 | 0.07 | | | 0.38 | 0.54 | | | 0.25 | 0.30 | |

¹Gleason 7 4+3 pattern and 8-10

²T3b, T4, N1, M1 at diagnosis or had metastases during follow-up or died due to prostate cancer

³adjusted for age

⁴adjusted for age, race/ethnicity, pack-years smoked in the past ten years, family history of prostate cancer, updated history of diabetes mellitus, height, body mass index at age 21, updated vigorous leisure-time physical activity, updated current aspirin use, cumulative average updated intake of tomato products and fish, and updated intake of energy, alcohol, vitamin E, α -linolenic acid, and total calcium

Table 3. Association of heterocyclic amine intake and MDM index with total prostate cancer by consumption of tomato products, Health Professionals Follow-up Study 1996-2010

| | Total prostate cancer | | | | | p ² |
|------------------------------------|--|------------|---|------------|----------------|----------------|
| | Lower consumption of tomato products (< 2 portions/week; 1607 cases) | | Higher consumption of tomato products (≥ 2 portions/week; 1163 cases) | | p ² | |
| | HR ¹ | 95% CI | HR ¹ | 95% CI | | |
| MelQx (ng/d) | | | | | | |
| <3.6 | 1.00 | ref. | 1.00 | ref. | | |
| 3.7 - <8.0 | 1.08 | 0.92 -1.26 | 0.98 | 0.80 -1.19 | | |
| 8.1 - <14.1 | 1.19 | 1.02 -1.40 | 1.01 | 0.83 -1.23 | | |
| 14.1 - <25.0 | 1.14 | 0.97 -1.34 | 0.98 | 0.80 -1.20 | | |
| ≥25.0 | 1.15 | 0.97 -1.37 | 1.07 | 0.87 -1.31 | | |
| <i>p-trend</i> | 0.08 | | 0.55 | | | 0.27 |
| PhIP (ng/d) | | | | | | |
| <25.6 | 1.00 | ref. | 1.00 | ref. | | |
| 25.6 - <51.4 | 1.04 | 0.89 -1.21 | 1.15 | 0.94 -1.41 | | |
| 51.4 - <89.4 | 1.07 | 0.92 -1.26 | 0.95 | 0.77 -1.17 | | |
| 89.4 - <152.9 | 1.09 | 0.93 -1.28 | 1.01 | 0.82 -1.23 | | |
| ≥152.9 | 1.14 | 0.96 -1.34 | 1.02 | 0.84 -1.25 | | |
| <i>p-trend</i> | 0.10 | | 0.63 | | | 0.12 |
| PhIP from red meat (ng/d) | | | | | | |
| <0.7 | 1.00 | ref. | 1.00 | ref. | | |
| 0.7 - <9.5 | 1.08 | 0.92 -1.27 | 1.00 | 0.81 -1.23 | | |
| 9.5 - <18.4 | 1.15 | 0.97 -1.35 | 1.06 | 0.86 -1.30 | | |
| 18.4 - <45.0 | 1.10 | 0.93 -1.30 | 0.95 | 0.78 -1.17 | | |
| ≥45.0 | 1.27 | 1.06 -1.52 | 1.05 | 0.85 -1.31 | | |
| <i>p-trend</i> | 0.02 | | 0.87 | | | 0.12 |
| PhIP from white meat (ng/d) | | | | | | |
| <14.4 | 1.00 | ref. | 1.00 | ref. | | |
| 14.4 - <29.9 | 1.09 | 0.93 -1.27 | 1.08 | 0.88 -1.32 | | |
| 29.9 - <56.3 | 1.11 | 0.95 -1.30 | 1.09 | 0.89 -1.33 | | |
| 56.3 - <110.4 | 1.10 | 0.93 -1.28 | 0.95 | 0.77 -1.16 | | |
| ≥110.4 | 1.15 | 0.98 -1.36 | 0.99 | 0.81 -1.21 | | |
| <i>p-trend</i> | 0.12 | | 0.42 | | | 0.29 |
| DiMelQx (ng/d) | | | | | | |
| <0.10 | 1.00 | ref. | 1.00 | ref. | | |
| 0.10 - <0.20 | 1.11 | 0.93 -1.33 | 1.21 | 0.97 -1.51 | | |
| 0.20 - <0.87 | 1.12 | 0.97 -1.30 | 1.09 | 0.92 -1.30 | | |
| 0.87 - <1.90 | 1.04 | 0.90 -1.20 | 0.91 | 0.76 -1.09 | | |
| ≥1.90 | 1.15 | 1.00 -1.34 | 0.98 | 0.83 -1.16 | | |
| <i>p-trend</i> | 0.11 | | 0.33 | | | 0.27 |
| MDM index (revertant colonies/day) | | | | | | |
| < 1222 | 1.00 | ref. | 1.00 | ref. | | |
| 1222- < 2256 | 1.12 | 0.96 -1.30 | 1.04 | 0.85 -1.27 | | |
| 2256 - < 3551 | 0.98 | 0.83 -1.15 | 1.09 | 0.89 -1.33 | | |
| 3551 - < 5663 | 1.13 | 0.96 -1.33 | 1.10 | 0.90 -1.33 | | |
| ≥ 5663 | 1.25 | 1.06 -1.47 | 1.01 | 0.83 -1.24 | | |
| <i>p-trend</i> | 0.02 | | 0.81 | | | 0.01 |

¹adjusted for age, race/ethnicity, pack-years smoked in the past ten years, family history of prostate cancer, updated history of diabetes mellitus, height, body mass index at age 21, updated vigorous leisure-time physical activity, updated current aspirin use, cumulative average updated intake of fish, and updated intake of energy, alcohol, vitamin E, α -linolenic acid, and total calcium

²p_i = p-interaction

Cancer Epidemiology, Biomarkers & Prevention

AACR American Association
for Cancer Research

Intake of meat mutagens and risk of prostate cancer in a cohort of U.S. health professionals

Sabine Rohrmann, Katharina Nimptsch, Rashmi Sinha, et al.

Cancer Epidemiol Biomarkers Prev Published OnlineFirst July 29, 2015.

Updated version Access the most recent version of this article at:
doi:[10.1158/1055-9965.EPI-15-0068-T](https://doi.org/10.1158/1055-9965.EPI-15-0068-T)

Author Manuscript Author manuscripts have been peer reviewed and accepted for publication but have not yet been edited.

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.