

Supplementary material

A gain-of-function mutation in the *CLCN2* chloride channel gene causes primary aldosteronism

Fabio L Fernandes-Rosa^{1,2,3*#}, Georgios Daniil^{1,2*}, Ian Orozco^{4,5*}, Corinna Göppner^{4,5\$}, Rami El Zein^{1,2\$}, Vandana Jain^{6^}, Sheerazed Boulkroun^{1,2^}, Xavier Jeunemaitre^{1,2,3}, Laurence Amar^{1,2,7}, Hervé Lefebvre^{8,9,10}, Thomas Schwarzmayr¹¹, Tim M Strom^{11,12}, Thomas J. Jentsch^{4,5#}, Maria-Christina Zennaro^{1,2,3#}

¹INSERM, UMRS_970, Paris Cardiovascular Research Center, Paris, France

²Université Paris Descartes, Sorbonne Paris Cité, Paris, France

³Assistance Publique-Hôpitaux de Paris, Hôpital Européen Georges Pompidou, Service de Génétique, Paris, France

⁴Leibniz-Forschungsinstitut für Molekulare Pharmakologie (FMP), Berlin, Germany

⁵Max-Delbrück-Centrum für Molekulare Medizin (MDC), Berlin, Germany

⁶Division of Pediatric Endocrinology, Department of Pediatrics, All India Institute of Medical Sciences, New Delhi, India

⁷Assistance Publique-Hôpitaux de Paris, Hôpital Européen Georges Pompidou, Unité Hypertension artérielle, Paris, France

⁸Normandie Univ, UNIROUEN, Rouen, France

⁹INSERM, DC2N, Rouen, France

¹⁰Department of Endocrinology, Diabetes and Metabolic Diseases, University Hospital of Rouen, Rouen, France

¹¹Institute of Human Genetics, Helmholtz Zentrum München, Neuherberg, Germany

¹²Institute of Human Genetics, Technische Universität München, Munich, Germany

*,\$,^equal contribution; #Corresponding author

Address correspondence to:

Maria-Christina Zennaro, MD, PhD

INSERM, U970

Paris Cardiovascular Research Center – PARCC

56, rue Leblanc,

75015 Paris – France

Tel : +33 (0)1 53 98 80 42

Fax : + 33 (0)1 53 98 79 52

e-mail : maria-christina.zennaro@inserm.fr

Fabio Fernandes Rosa, MD, PhD

INSERM, U970

Paris Cardiovascular Research Center – PARCC

56, rue Leblanc,

75015 Paris – France

Tel : +33 (0)1 53 98 80 43

Fax : + 33 (0)1 53 98 79 52

e-mail : fabio.fernandes-rosa@inserm.fr

Thomas J. Jentsch, MD, PhD

FMP / MDC

Robert-Rössle-Strasse 10

13125 Berlin – Germany

Tel: +49 30 9406 2961

Fax: +49 30 9406 2960
e-mail: Jentsch@fmp-berlin.de

Supplementary Table 1: Expression of plasma membrane chloride channels in the human adrenal cortex.

Gene name	protein	mRNA expression*
<i>CLCN2</i>	ClC-2	1.65±0.23/1.16±0.18**
<i>CLCN1</i>	ClC-1	0.51±0.07
<i>CFTR</i>	CFTR	0.46±0.05
<i>LRRC8A</i> ^{\$}	LRRC8A	1.09±0.22
<i>LRRC8B</i> [#]	LRRC8B	0.62±0.14/0.42±0.041**
<i>LRRC8C</i> [#]	LRRC8C	2.11±0.22
<i>LRRC8D</i> [#]	LRRC8D	9.95±0.84
<i>LRRC8E</i> [#]	LRRC8E	0.79±0.11
<i>TMEM16A</i>	Anoctamin-1	0.44±0.065

*mRNA expression was retrieved from a transcriptome study including 123 APA and 11 CA¹. Values represent median centred, log2-transformed and model-adjusted expression levels represented as mean±SEM. **Values represent expression levels detected by two different probes. ^{\$}Essential and [#]non-essential components of the volume-regulated anion channel (VRAC)².

Supplementary Table 2. Primers used for *CLCN2* sequence

Exon	Forward primer	Reverse primer
1	CAGGACAGAGCCCGGAACC	GGACAGGGATTAGGGTAGGCC
2	CATAAGCATGGTCCACTCCC	AGCAGCTCTAATGGCCTCTG
10	AGGCTCCTTTCACTCAGGT	CCTGTTTGACTGGGCCATT

Supplementary Table 3. Primers used for real-time RT-qPCR

Gene Symbol	Forward primer	Reverse primer
<i>18S</i>	CCCTGCCTTGTACACACC	CGATCCGAGGGCCTCACTA
<i>HPRT</i>	CTCAACTTTAACCTGGAAAGAATGTC	TCCTTTCAACCAGCAAGCT
<i>GAPDH</i>	TGCACCACCAACTGCTTAGC	GGCATGGACTGTGGTCATGAG
<i>CLCN2</i>	TTGATCCTGCTCCCTTCCAG	CATAAGCATGGTCCACTCCC
<i>StAR</i>	ATGAGTAAAGTGGTCCCAGATG	ACCTTGATCTCCTTGACATTGG
<i>CYP21A2</i>	GAGTAGTCTCCAAGGACAGGT	GTGGTGCTGAACTCCAAGAGGA
<i>CYP11B2</i>	GTGTGGAAGGAGCACTTGAGG	GATGCCTGTGTAGTGGAGGC

References

1. Boulkroun, S. *et al.* Prevalence, Clinical, and Molecular Correlates of KCNJ5 Mutations in Primary Aldosteronism. *Hypertension* **59**, 592-8 (2012).
2. Voss, F.K. *et al.* Identification of LRRC8 heteromers as an essential component of the volume-regulated anion channel VRAC. *Science* **344**, 634-8 (2014).