

# C/EBP $\beta$ Regulation of the Tumor Necrosis Factor $\alpha$ Gene

Richard M. Pope, Achim Leutz,\* and Scott A. Ness<sup>†</sup>

Department of Medicine, Division of Arthritis and Musculoskeletal Diseases, Northwestern University, and the Veterans Administration Lakeside Medical Center, Chicago, Illinois 60611; \*Zentrum für Molekulare Biologie, Universität Heidelberg, D-6900 Heidelberg, Germany; and <sup>†</sup>Department of Biochemistry, Molecular Biology and Cell Biology, Northwestern University, Evanston, Illinois 60208

## Abstract

**Activated macrophages contribute to chronic inflammation by the secretion of cytokines and proteinases. Tumor necrosis factor  $\alpha$  (TNF $\alpha$ ) is particularly important in this process because of its ability to regulate other inflammatory mediators in an autocrine and paracrine fashion. The mechanism(s) responsible for the cell type-specific regulation of TNF $\alpha$  is not known. We present data to show that the expression of TNF $\alpha$  is regulated by the transcription factor C/EBP $\beta$  (NF-IL6). C/EBP $\beta$  activated the TNF $\alpha$  gene promoter in cotransfection assays and bound to it at a site which failed to bind the closely related protein C/EBP $\alpha$ . Finally, a dominant-negative version of C/EBP $\beta$  blocked TNF $\alpha$  promoter activation in myeloid cells. Our results implicate C/EBP $\beta$  as an important regulator of TNF $\alpha$  by myelomonocytic cells. (*J. Clin. Invest.* 1994. 94:1449–1455.)** **Key words:** transcription factor • cytokine • promoter • inflammation

## Introduction

Activated macrophages produce an array of inflammatory cytokines including tumor necrosis factor  $\alpha$  (TNF $\alpha$ ), interleukins-1, -6, and -8, macrophage chemotactic protein-1, and macrophage inflammatory protein 1 $\alpha$  (MIP-1 $\alpha$ )<sup>1</sup> (1–6). Many of these cytokines are capable of both attraction and activation of additional inflammatory cells and induction of angiogenesis (7). In addition, in chronic inflammation, such as rheumatoid arthritis, macrophages contribute directly to joint destruction by producing proteases, such as collagenase (matrix metalloproteinase 1) and stromelysin (matrix metalloproteinase 3) (8, 9). TNF $\alpha$  may play a crucial role in the inflammatory process, since it is capable of inducing cytokines and matrix metalloproteinases in an autocrine and paracrine fashion (10, 11).

Achim Leutz's present address is Max Delbrück Center for Molecular Medicine, Robert-Rössly Str. 10, 13122 Berlin, Germany. Address correspondence to Richard M. Pope, M.D., Northwestern University Medical School, Department of Medicine, Division of Arthritis and Connective Tissue Diseases, Ward 3-315, 303 E. Chicago Avenue, Chicago, IL 60611.

Received for publication 18 February 1994 and in revised form 2 June 1994.

1. *Abbreviations used in this paper:* c, chicken; C/EBP, CCAAT/enhancer binding protein; cMGF, chicken monocyte growth factor; CMV, cytomegalovirus; DN, dominant-negative; EMSA, electrophoretic mobility shift assay; h, human; MIP-1 $\alpha$ , macrophage inflammatory protein 1 $\alpha$ ; r, rat.

A variety of stimuli can induce cells to produce TNF $\alpha$ , including infection by viruses or other agents, interaction with adhesion molecules, or treatment with phorbol esters, endotoxins, or cytokines such as IL-1 or even TNF $\alpha$  itself (12–14). Examination of the upstream promoter region of the human TNF $\alpha$  gene has revealed both cell type- and stimulus-specific regulatory elements, including potential binding sites for the transcription factors NF- $\kappa$ B and AP-1 (15–22). Although many cell types including lymphocytes, smooth muscle cells, mast cells, and others are capable of TNF $\alpha$  synthesis, activated monocytes/macrophages are the principal source of TNF $\alpha$  (12). Nonetheless, little is known about how the TNF $\alpha$  gene is regulated in a cell type-specific fashion.

C/EBP $\beta$  (CCAAT/enhancer binding protein  $\beta$ , also called NF-IL6, NF-M, LAP, IL6-DBP, AGP/EBP, and CRP2) is a member of a family of transcription factors termed bZip proteins which have a basic DNA-binding domain linked to a leucine zipper dimerization motif. It is closely related to the founding member of this family, C/EBP $\alpha$ , which has been shown to be intimately involved in the regulation of adipocyte differentiation (23). Earlier studies identified monocytes and macrophages as the principal source of C/EBP $\beta$  (24, 25). Cell type-specific gene regulation by C/EBP $\beta$  has been shown to depend on interactions with other transcription factors, such as NF- $\kappa$ B, c-jun, the glucocorticoid receptor, and the nuclear oncoprotein v-Myb (26–29). In addition, binding sites for C/EBP $\beta$  have been identified in the promoters for a large number of cytokine genes which are expressed in myeloid cells, including the genes for interleukins-1, -6, and -8, MIP-1 $\alpha$ , and chicken monocyte growth factor (cMGF) (24, 30–35), suggesting that the factor may be an important activator of inflammation and of growth regulatory genes. Finally, ectopic expression of Myb and C/EBP $\beta$  has been shown to induce the expression of myeloid-specific genes in heterologous cell types, implicating C/EBP $\beta$  as a key regulator of myeloid cell differentiation (27).

In this paper, we have investigated the relationship between C/EBP $\beta$  activity and TNF $\alpha$  expression. We identified a specific binding site for C/EBP $\beta$  in the TNF $\alpha$  gene promoter, suggesting that C/EBP $\beta$  directly regulates the expression of the cytokine. We tested this in cotransfection assays and found that C/EBP $\beta$ , more strongly than C/EBP $\alpha$ , activated a TNF $\alpha$  promoter-reporter construct. Furthermore, ectopic expression of a dominant-negative form of C/EBP $\beta$  was able to block activation of the TNF $\alpha$  promoter by endogenous transcription factors. Our results demonstrate that C/EBP $\beta$  is an important regulator of the TNF $\alpha$  gene and that it is responsible for the cell type-specific regulation of the TNF $\alpha$  gene in myelomonocytic cells.

## Methods

*Cell lines.* The human myelomonocytic cell lines U937 and THP-1 and the Jurkat human T cell line were obtained from the American Type

Culture Collection (Rockville, MD) and were maintained in RPMI medium supplemented with 10% heat-inactivated fetal bovine serum, glutamine, penicillin, and streptomycin. HD-11, an MC29 virus (v-myc oncogene) transformed chicken macrophage cell line, was maintained in Dulbecco's modified Eagle's medium supplemented with 8% fetal bovine serum, 3% chicken serum, glutamine, and antibiotics as above.

**Plasmids.** The human C/EBP $\beta$  expression vector (hC/EBP $\beta$ ), which expresses the NF-IL6 cDNA under control of the cytomegalovirus (CMV) promoter (from S. Akira, Osaka University, Japan), has been described (24), as have similar plasmids that express full-length chicken C/EBP $\beta$  (cC/EBP $\beta$ , also referred to as NF-M) and the 5D229 dominant-negative deletion mutant (27, 31). The latter plasmid expresses a protein having an NH<sub>2</sub>-terminal deletion so that only the last 229 amino acids remain. This protein is 92% identical and 99% homologous to its human counterpart, and the COOH-terminal 212 amino acids, representing the DNA binding and leucine zipper domains, are 99% identical. The TNF $\alpha$  promoter-luciferase reporter plasmids have been described (15, 22). The pCDM8 expression vector plasmid was purchased from Invitrogen (San Diego, CA).

**Transfections and reporter gene assays.** Jurkat T cells were transfected using DEAE-dextran as follows. On the day before transfection, the cells were split to  $\sim 0.7 \times 10^6$  cells/ml. For each sample,  $\sim 1 \times 10^7$  cells were washed twice, then resuspended in 2 ml STBS (25 mM Tris/HCl, pH 7.5, 137 mM NaCl, 5 mM KCl, 0.6 mM Na<sub>2</sub>PO<sub>4</sub>, 0.7 mM CaCl<sub>2</sub>, and 0.5 mM MgCl<sub>2</sub>) containing 750  $\mu$ g/ml DEAE-dextran and a total of 15  $\mu$ g of CsCl-purified plasmid DNA (specific plus carrier), for 30 min at 37°C, with gentle inversion every 5 min. The cells were then collected by centrifugation, resuspended in fresh medium, and allowed to recover for 3–5 h before treatment with 20 ng/ml PMA (Sigma Chemical Co., St. Louis, MO) as indicated. Cells were harvested 18–24 h later. The U937 cells were transfected by electroporation as described previously (15), using an electroporation apparatus from Invitrogen. DEAE-dextran-mediated transfection of HD-11 cells and luciferase assays have been described (27, 31). All transfection results were normalized for protein content, as determined using an assay reagent from Bio-Rad Laboratories (Richmond, CA). All experiments were performed a minimum of three times, and similar results were obtained using independent preparations of DNA.

**Electrophoretic mobility shift assays (EMSAs).** Nuclear extracts were obtained from U937, THP-1, and Jurkat cells as described (36). All buffers were supplemented with PMSF (20  $\mu$ g/ $\mu$ l), aprotinin, pepstatin A, and leupeptin (each 5  $\mu$ g/ml). For expression in insect cells, the cC/EBP $\beta$  cDNA was transferred to the transfer vector pVL1392 and transfected into Sf9 cells along with linearized Baculovirus DNA using a kit from PharMingen (San Diego, CA). The resulting virus was used to infect new Sf9 or High-5 cells, and nuclear extracts were prepared 2 d later as follows. All buffers were chilled and supplemented with protease inhibitors as described above. The cells were collected, washed twice with PBS, and resuspended in Tris-X buffer (20 mM Tris-Cl, pH 7.3, 50 mM KCl, 1 mM EDTA, 1.5 mM MgCl<sub>2</sub>, and 0.5 mM CaCl<sub>2</sub>) at a concentration of  $1 \times 10^7$ /ml. The cells were lysed by adding NP-40 to a final concentration of 0.5%, then the nuclei were recovered by centrifugation, resuspended at fourfold higher concentration of Tris-X buffer plus 0.3 M KCl, left 5 min on ice, and then centrifuged again. The supernatant was diluted with three volumes of Tris-X buffer (to make 0.085 M KCl final), and centrifuged again, and the resulting nuclear extract was stored at  $-80^\circ\text{C}$ . Bacterially expressed, purified rat C/EBP $\beta$  (rC/EBP $\beta$ ) and rat C/EBP $\alpha$  (rC/EBP $\alpha$ ) proteins were generously provided by P. Johnson (National Institutes of Health, Bethesda, MD). Bacterially expressed human NF- $\kappa$ B1 (p50) was purchased from Promega Corp. (Madison, WI).

The methods for radiolabeling the double-stranded oligonucleotide probes using Klenow polymerase and [ $\alpha$ -<sup>32</sup>P]dCTP and for performing the EMSAs have been described (27). Where indicated, antibodies or excess unlabeled competitor oligonucleotides were incubated with the proteins or nuclear extracts for 20 min on ice before the addition of the probe. The samples were then incubated for an additional 20 min before electrophoresis on nondenaturing polyacrylamide gels. Antibodies spe-

cific for hC/EBP $\beta$ , cC/EBP $\beta$ , and rC/EBP $\beta$  were obtained from S. Akira, A. Leutz, and P. Johnson, respectively.


**Synthetic oligonucleotides.** The following synthetic oligonucleotides were used to generate the  $-130/+35$  TNF $\alpha$  radiolabeled EMSA probe using the polymerase chain reaction: forward, 5'-GGCCACTACCG-CTTCCTC-3'; reverse, 5'-CCTCTGCTGTCCTTGCTGA-3'. The following radiolabeled oligonucleotides were used as probes in EMSAs: the  $-100/-74$  C/EBP $\beta$  binding site of the TNF $\alpha$  promoter, 5'-TCGACATGGGTTTCTCCACCAAGGAAGTTTTC-3', 3'-GTACCCAAAGAGGTGGTTCCTTCAAAAAGAGCT-5', the C/EBP $\beta$  binding sequence from the IL-6 promoter (24), 5'-TCGAGACATTGCACATCTG-3', 3'-CTGTAACGTCTTAGACAGCT-5', and the Ig/HIV promoter binding sequence for NF- $\kappa$ B (37), 5'-TCGAGAAGGGACTTTCCGGCTGGGGACTTTCCAGGGC-3', 3'-CTTCCCTGAAAGGCGACCCCTGAAAGGTCCCGAGCT-5'.

**DNaseI footprinting.** The  $-615$  TNF $\alpha$ -luciferase plasmid was digested with XhoI, radiolabeled by end-filling using Klenow polymerase and [ $\alpha$ -<sup>32</sup>P]dCTP, and then further digested with HindIII. The resulting fragment containing the TNF $\alpha$  promoter was purified after agarose gel electrophoresis using GeneClean (Bio101) and subjected to DNaseI footprinting as described (38).

## Results

**C/EBP $\beta$  activates the TNF $\alpha$  promoter.** The TNF $\alpha$  promoter has been shown to be strongly activated by treatment with PMA (15, 22), an activator of protein kinase C which also activates C/EBP $\beta$  activity (31, 39). We used transient transfection assays to determine whether C/EBP $\beta$  could cooperate with PMA treatment to activate the TNF $\alpha$  promoter. We chose the Jurkat human T cell line, which normally does not express either C/EBP $\beta$  or TNF $\alpha$ , under the conditions used in our assays (data not shown). A series of TNF $\alpha$  promoter-luciferase reporter gene constructs, containing portions of the promoter extending from just downstream to several hundred nucleotides upstream of the transcription start site (22), were transfected into Jurkat cells either with a control plasmid or with a plasmid expressing hC/EBP $\beta$ , under the control of the constitutive CMV promoter. Parallel experiments were performed with and without PMA treatment. As shown in Fig. 1, none of the TNF $\alpha$  promoter constructs were expressed at significant levels in the absence of PMA treatment, even in the presence of hC/EBP $\beta$ , and none were significantly activated by PMA treatment alone. However, PMA treatment of cells expressing C/EBP $\beta$  led to strong activation of all but the shortest promoter constructs tested ( $-36$  bp from the transcription start site), suggesting that PMA and C/EBP $\beta$  acted cooperatively to induce TNF $\alpha$  promoter activity. In addition, the data suggest that crucial PMA- and C/EBP $\beta$ -responsive elements must lie in the region bounded by positions  $-95$  and  $-36$  in the TNF $\alpha$  promoter. The promoter construct extending only to position  $-95$  was consistently less active than the longer versions ( $-120$  and  $-295$ ), suggesting that an important regulatory element(s) might lie near the  $-95$  site in the promoter.

**C/EBP $\beta$  binds a specific site in the TNF $\alpha$  promoter.** The experiments described above suggested that C/EBP $\beta$  could regulate the TNF $\alpha$  promoter. An EMSA was used to determine whether the promoter contained a bona fide C/EBP $\beta$  binding site. Based on our results with the promoter-reporter gene constructs (see above), we generated a radiolabeled fragment of the TNF $\alpha$  promoter bounded by positions  $-130$  to  $+35$  relative to the transcription start site. Fig. 2 A shows an EMSA experiment in which such an oligonucleotide was efficiently bound




**Figure 1.** *C/EBPβ* activates the *TNFα* promoter. Transfections of Jurkat T cells were performed as described in Methods, using 0.5  $\mu$ g of the h*C/EBPβ* plasmid and 3  $\mu$ g of the indicated *TNFα* promoter reporter construct. On the left of the bar graphs are diagrammed the *TNFα* reporter-luciferase constructs used. Potential binding sites for a number of transcription factors are indicated by labeled boxes. The relative light units have been corrected for the total protein in the cell lysates. Similar results have been obtained in > 10 experiments and with different plasmid preparations.


by insect cell-expressed chicken *C/EBPβ* (c*C/EBPβ*). Binding was inhibited completely by the addition of excess oligonucleotide containing the *C/EBPβ*-binding site from the human IL-6 promoter, and the addition of antibodies specific for c*C/EBPβ* resulted in a "supershift" of the protein-DNA complex (not shown), demonstrating the specificity of the binding reac-

tion. Similar complexes were also formed using nuclear extracts from the human myeloid cell line THP-1 (not shown). These results suggest that *C/EBPβ* could play a direct role in the regulation of the *TNFα* promoter.

A DNaseI footprint assay was used to more precisely localize the binding sites for *C/EBPβ* on the *TNFα* promoter. However, as shown in Fig. 2 *B*, the addition of THP-1 nuclear extracts to an end-labeled fragment of the *TNFα* promoter resulted in the formation of a new DNaseI hypersensitive site at position -90 relative to the start site of transcription, rather than a discrete protected region. These results suggest that the extracts contained a protein or complex that specifically interacted with this region of the promoter. Addition of antibodies specific for h*C/EBPβ*, but not nonimmune serum, blocked the hypersensitive site formation, indicating that *C/EBPβ*, or a protein complex containing *C/EBPβ*, was responsible for this change in the DNaseI sensitivity of the promoter.

Two potential *C/EBPβ*-binding sites, in opposite orientations, were noted extending from -97 to -79 bp relative to the transcription start site of the *TNFα* promoter, which conformed at 9 of 12 positions to the consensus *C/EBPβ*-binding site, T(T/G)NNGNAA(T/G) (24). As shown in Fig. 3, insect cell-expressed c*C/EBPβ* protein bound efficiently to a radiolabeled, synthetic 26-mer oligonucleotide containing this site. Again, *C/EBPβ* binding was inhibited completely when an excess of either unlabeled oligonucleotide used as the probe or one containing a previously identified *C/EBPβ*-binding site from the human IL-6 gene promoter was included. The complex was not inhibited by an unlabeled NF- $\kappa$ B binding oligonucleotide representing the Ig/HIV promoter (not shown), even though the oligonucleotide used as the probe contained a previously characterized binding site for NF- $\kappa$ B (data not shown; 17). Addition of antibodies specific for c*C/EBPβ*, but not control antibodies, resulted in a supershift of the protein-DNA complex. Similar results have also been obtained using recombinant, purified rat *C/EBPβ* (r*C/EBPβ*) (Fig. 4 *A*; data not shown). Although *C/EBPβ* bound quite efficiently to the *TNFα* promoter-derived binding site, the oligonucleotide derived from the IL-6 promoter was a better competitor when the -100/-74 *TNFα* oligonucleotide was used as probe, suggesting that the binding site in the *TNFα* promoter was relatively weak. We conclude that *C/EBPβ* regulates the human *TNFα* gene pro-





**Figure 3.** Binding of cC/EBP $\beta$  to the  $-100/-74$  human TNF $\alpha$  promoter oligonucleotide. The EMSA was performed with Baculovirus-expressed cC/EBP $\beta$  and with the radiolabeled  $-100/-74$  TNF $\alpha$  promoter oligonucleotide. The competitors included an unlabeled, irrelevant, control oligonucleotide (C), and unlabeled oligonucleotides representing the C/EBP $\beta$ -binding site of the IL-6 promoter (IL-6) and that used as the radiolabeled probe (TNF) (lanes 1–4). These competitors were used at 50- or 500-fold excess as indicated in the figure. Nonimmune (N) or immune (I), anti-cC/EBP $\beta$ , rabbit antisera were included where indicated (lanes 5 and 6).

moter by binding directly to a specific site and activating its transcription.

**Selectivity of binding to  $-100/-74$  TNF $\alpha$  oligonucleotide.** C/EBP $\beta$  and C/EBP $\alpha$  are both expressed in macrophage cell lines and are capable of heterodimerization and of binding to the same oligonucleotides (40–42). We next examined the ability of C/EBP $\alpha$  and C/EBP $\beta$  to interact differently with the TNF $\alpha$  promoter. Using an EMSA, purified recombinant C/EBP $\beta$  and C/EBP $\alpha$  both bound strongly to the oligonucleotide derived from the IL-6 promoter (Fig. 4 A). Binding by both isoforms to the IL-6 probe was inhibited by excess unlabeled IL-6, but not TNF $\alpha$ , promoter oligonucleotide. Similar results have been observed previously with oligonucleotides derived from the cMGF promoter (31, 43). In contrast, only C/EBP $\beta$  bound to the  $-100/-74$  TNF $\alpha$  oligonucleotide, suggesting that the site was specific for only one member of this highly related family of bZip proteins (Fig. 4 A).

A cotransfection assay was used to test whether both bZip proteins could activate the TNF $\alpha$  promoter. Jurkat T cells were cotransfected with TNF $\alpha$  or cMGF promoter-reporter constructs, plus either C/EBP $\beta$  or C/EBP $\alpha$  expression vectors. Transfected cells were then treated with PMA. Both transcription factors activated the cMGF promoter to approximately equal levels (Fig. 4 B). In contrast, C/EBP $\beta$  was a stronger activator of the TNF $\alpha$  promoter than C/EBP $\alpha$ , consistent with its stronger binding in the EMSA. When C/EBP $\alpha$  was coexpressed with C/EBP $\beta$ , the activation of the TNF $\alpha$  promoter was comparable with that observed with C/EBP $\beta$  alone, suggesting either that C/EBP $\alpha/\beta$  heterodimers were as active as C/EBP $\beta$  homodimers or that C/EBP $\beta$  was somehow dominant.

**Dominant-negative C/EBP $\beta$  inhibits the TNF $\alpha$  promoter.** Overexpression of C/EBP $\beta$  could affect the regulation of the TNF $\alpha$  promoter in two ways, either by binding to the promoter


and directly activating its transcription or by interfering with negative-regulatory effects of other bZip proteins such as C/EBP $\alpha$ , perhaps by forming inactive heterodimers (42). This was tested using a truncated, dominant-negative (DN) version of the chick C/EBP $\beta$  protein (27, 31), termed DN229. This version of C/EBP $\beta$  only contains the COOH-terminal 229-most amino acids. It therefore lacks the transactivation domain, leaving the dimerization and DNA-binding domains intact. It is capable of binding DNA, but has no transactivation activity, and forms inactive heterodimers with other C/EBP-like proteins (31). The DN229 construct has been shown to be quite effective at suppressing the expression of other C/EBP $\beta$ -regulated genes, such as the cMGF and *mim-1* genes (27, 31). When compared with control expression vectors without an insert, cotransfection of the DN229 plasmid was quite effective at inhibiting the ability of hC/EBP $\beta$  to activate the TNF $\alpha$  promoter in Jurkat cells, even though the cells were treated with PMA (Fig. 5). This indicates that the DN229 construct is not only unable to activate the TNF $\alpha$  promoter, but that expression of a truncated form of C/EBP $\beta$  blocks activation by the full-length protein.

A similar experiment was performed using the human myelomonocyte cell line U937, which expresses endogenous C/EBP $\beta$  protein and secretes a high concentration of TNF $\alpha$  in response to PMA (data not shown) to determine whether ectopically expressed DN229 could likewise block the expression of the TNF $\alpha$  promoter in myeloid cells. Fig. 6 shows that the TNF $\alpha$  promoter construct extending to position  $-120$ , the construct that consistently provided the greatest activation, was moderately active when transfected into U937 cells and that its expression was stimulated three- to fivefold when the cells were treated with PMA. (Similar results were obtained with other myeloid cell lines, including human THP-1 and chicken HD-11 cells.) Expression of the dominant-negative form of C/EBP $\beta$  (DN229) not only blocked activation by PMA, but also inhibited TNF $\alpha$  promoter activity to below basal levels. In contrast, the control plasmid containing only the CMV promoter had no effect on the activity of the TNF $\alpha$  promoter-reporter gene construct. We conclude that the human TNF $\alpha$  promoter is directly regulated by C/EBP $\beta$ , most probably by binding to the region between  $-95$  and  $-36$  relative to the transcription start site.

## Discussion

Activation of TNF $\alpha$  has been implicated in a wide variety of responses to inflammatory and infectious agents, and it is known to be highly expressed by activated macrophages, such as those found in the synovial lining of rheumatoid joints (44, 45). The results in this paper demonstrate that the transcription factor C/EBP $\beta$  is a key regulator of the TNF $\alpha$  gene. We have identified a specific C/EBP $\beta$ -binding site in the TNF $\alpha$  promoter, shown that the promoter responds dramatically to the expression of C/EBP $\beta$ , and used a dominant-negative version of C/EBP $\beta$  to inhibit the activation of the TNF $\alpha$  promoter.

The C/EBP $\beta$ -binding site identified by DNaseI footprinting and EMSA experiments maps to positions  $-100/-74$  of the human TNF $\alpha$  promoter. However, compared with the oligonucleotide from the IL-6 promoter, C/EBP $\beta$  bound the  $-100/-74$  site relatively weakly. Nevertheless, in DNaseI footprinting assays, which are generally much less sensitive than EMSAs, myeloid cell extracts induced the formation of a DNaseI hypersensitive site on the TNF $\alpha$  promoter at position  $-90$ ,



**Figure 4.** The TNF $\alpha$  promoter is specific for C/EBP $\beta$ . (A) Selective binding of C/EBP $\beta$ , but not C/EBP $\alpha$ , to the  $-100/-74$  TNF $\alpha$  promoter. The EMSA was performed with recombinant, purified rC/EBP $\alpha$  (lanes 3–6) and rC/EBP $\beta$  (lanes 1, 2, and 7–9), as indicated at the top of the figure. The radiolabeled probes, which included the C/EBP $\beta$ -binding site of the IL-6 promoter (IL-6) and the  $-100/-74$  TNF $\alpha$  promoter (TNF) oligonucleotides, and the competitor oligonucleotides, used at 100-fold excess, are also indicated at the top of the figure (symbols as in Fig. 3). Nonimmune (N) or monospecific, immune anti-rC/EBP $\beta$  (I) sera were added where indicated. The mobilities of the C/EBP $\alpha$  and C/EBP $\beta$  bound and the free, unbound radiolabeled  $-100/-74$  TNF $\alpha$  oligonucleotide are indicated on the left of the figure. (B) C/EBP $\beta$  preferentially activates the TNF $\alpha$  promoter. Jurkat cells were transfected as described in Methods. They were cotransfected with the TNF $\alpha$ -luciferase (3  $\mu$ g) or with cMGF-luciferase (4.5  $\mu$ g) constructs and with the hC/EBP $\beta$  or the rC/EBP $\alpha$  expression vectors in the concentrations indicated at the bottom of the figure. Where indicated, both C/EBP $\alpha$  and C/EBP $\beta$  were cotransfected at equal concentrations. The fold increase represents the fold increase over baseline induced by PMA. These experiments were repeated twice with similar results.


and this activity was inhibited by C/EBP $\beta$ -specific antisera. Thus, in the context of a nuclear extract, the C/EBP $\beta$  appears to interact with a relatively high affinity for the promoter, suggesting that the factor may be part of a multiprotein complex which binds the TNF $\alpha$  promoter in myeloid cells.

The TNF $\alpha$  promoter was differentially recognized by C/EBP $\beta$  and C/EBP $\alpha$ . Purified recombinant C/EBP $\beta$ , but not C/EBP $\alpha$ , bound to the TNF $\alpha$  promoter oligonucleotide in EMSA assays. Consistent with this difference, C/EBP $\beta$  was considerably more active than C/EBP $\alpha$  at activating the TNF $\alpha$  promoter construct. As a control, the cMGF promoter, which readily binds both C/EBP proteins, was activated equally well by each. Both C/EBP $\alpha$  and C/EBP $\beta$  are expressed in myelomonocytic cell lines, however, C/EBP $\alpha$  decreased while C/EBP $\beta$  increased with differentiation and activation (40, 41). These data suggest that the differential binding of the two transcription factors may be important in the regulation of TNF $\alpha$  in monocytes/macrophages where both bZip proteins are expressed.

The importance of the region of the TNF $\alpha$  promoter that contains the C/EBP $\beta$ -binding site (at position  $-90$ ) has also been noted by others. For example, three copies of a 43-nucleotide-long fragment of the promoter spanning this site was suffi-


cient to confer both elevated basal and TNF $\alpha$ -inducible activity to an otherwise minimally active heterologous promoter (21). More recently, a cyclosporin A- and FK506-sensitive factor in the murine T cell line Ar-5 was identified that functions through a portion of the promoter overlapping the C/EBP $\beta$ -binding site (17). Of interest, the  $-90$  site is adjacent to a potential binding site for NF- $\kappa$ B (17, 24), which has been shown to interact with C/EBP $\beta$  in vitro (26, 29). In addition, both C/EBP $\beta$  and NF- $\kappa$ B bound to the IL-6 and the IL-8 gene promoters, and the ratio of the two factors appeared to regulate the activities of those genes (34, 46). Our own preliminary data suggest that overexpression of NF- $\kappa$ B1 or RelA may inhibit C/EBP $\beta$ -mediated activation of the TNF $\alpha$  promoter in Jurkat cells (Pope, R., and S. A. Ness, unpublished observations). Further work will be required to determine whether these two transcription factors interact to regulate the TNF $\alpha$  gene.

An interesting feature of TNF $\alpha$  gene regulation in Jurkat cells was its apparent dependence on both C/EBP $\beta$  and PMA treatment, which presumably participates by stimulating protein kinase C. PMA treatment has been shown to lead to phosphorylation of C/EBP $\beta$ , induce its translocation to the nucleus, and enhance its ability to stimulate transcription (31, 39), sug-



**Figure 5.** Dominant-negative C/EBP $\beta$  inhibits TNF $\alpha$  promoter activation in Jurkat T cells. These experiments were performed as described in Fig. 2 except that in addition to the TNF $\alpha$  reporter (3  $\mu$ g) and the hC/EBP $\beta$  (0.5  $\mu$ g) expression plasmids, the DN C/EBP $\beta$  (DN229) or vector (pCDM8) plasmids were cotransfected in the indicated quantities. Relative light units were corrected for the total protein in the cell lysates. Percent suppression was determined by comparison with the control vector. These experiments were repeated three times with similar results.

gesting that PMA treatment may be required simply to convert the over-expressed C/EBP $\beta$  to a more active form. Alternatively, PMA may activate other transcription factors which cooperate with C/EBP $\beta$ , such as NF- $\kappa$ B or *c-jun* (47, 48). Several potential PMA-responsive elements have been identified in the TNF $\alpha$  promoter, including binding sites for the transcription factors AP-1 and AP-2 located at positions -63 and -34, respectively (22). Cotransfection with *c-jun* enhanced and mutation of the AP-1-binding site diminished TNF $\alpha$  gene expression, supporting the importance of the AP-1 site (22). Whether or not interaction between C/EBP $\beta$  and *c-jun* is important in



**Figure 6.** PMA-induced TNF $\alpha$  promoter activation in U937 cells is inhibited by DN C/EBP $\beta$ . The transfections were performed by electroporation as described in Methods, using 5  $\mu$ g of the -120 TNF $\alpha$  reporter construct, and 0.5  $\mu$ g of the DN C/EBP $\beta$  (DN229) or vector (pCDM8). Relative light units were corrected for the total protein in the cell lysates. These experiments were repeated four times with similar results.

TNF $\alpha$  activation (49), and whether the TNF $\alpha$  gene can be activated in the absence of C/EBP $\beta$ , for example in other cell types that do not express the factor, remains to be determined.

In addition to TNF $\alpha$ , a number of other cytokine genes, such as IL-1, IL-6, IL-8, and MIP-1 $\alpha$  also possess functional C/EBP $\beta$ -binding sites in their promoters (24, 30, 32-35). Many of these are considered important mediators of inflammation and are produced in response to TNF $\alpha$  (3-6). The expression of C/EBP $\beta$  may be important in chronic inflammation associated with a variety of disorders. The inhibition of C/EBP $\beta$  activity, perhaps mediated by ectopic expression of a dominant-negative form of the protein, presents a potential means of blocking the synthesis of cytokines like TNF $\alpha$  which contribute to chronic inflammation. Further work with dominant-negative forms of C/EBP $\beta$  will be required to determine if such therapy is possible.

## Acknowledgments

The authors commend Yi Shen for her excellent technical assistance and thank S. Akira, J. Economou, and P. Johnson for their generous gifts of plasmids, antisera, and bacterially expressed proteins used in this study.

R. M. Pope was supported by a National Institutes of Health Multi-purpose Arthritis and Musculoskeletal Diseases Center grant (P60-AR30692), by the Veterans Administration Research Service, and the Illinois Chapter of the Arthritis Foundation. S. A. Ness was supported by grants from the Leukemia Research Foundation and the Illinois Division of the American Cancer Society.

## References

- Firestein, G. S., J. M. Garcia-Alvaro, and R. Maki. 1990. Quantitative analysis of cytokine gene expression in rheumatoid arthritis. *J. Immunol.* 144:3347-3353.
- Guerne, P.-A., B. L. Zuraw, J. H. Vaughan, D. A. Carson, and M. Lotz. 1989. Synovium as a source of interleukin 6 in vitro. *J. Clin. Invest.* 83:585-592.
- Koch, A. E., S. L. Kunkel, J. C. Burrows, H. L. Evanoff, G. K. Haines, R. M. Pope, and R. M. Strieter. 1991. Synovial tissue macrophage as a source of the chemotactic cytokine IL-8. *J. Immunol.* 147:2187-2195.
- Koch, A. E., S. L. Kunkel, S. W. Chensue, G. K. Haines, and R. M. Strieter. 1992. Expression of interleukin-1 and interleukin-1 receptor antagonist by human rheumatoid synovial tissue macrophages. *Clin. Immunol. Immunopathol.* 65:23-29.
- Koch, A. E., S. L. Kunkel, L. A. Harlow, B. Johnson, H. L. Evanoff, G. K. Haines, M. D. Burdick, R. M. Pope, and R. M. Strieter. 1992. Enhanced production of monocyte chemoattractant protein-1 in rheumatoid arthritis. *J. Clin. Invest.* 90:772-779.
- Koch, A. E., S. L. Kunkel, L. A. Harlow, D. D. Mazarakis, G. K. Haines, M. D. Burdick, R. M. Pope, and R. M. Strieter. 1994. Macrophage inflammatory protein-1 $\alpha$ . A novel chemotactic cytokine for macrophages in rheumatoid arthritis. *J. Clin. Invest.* 93:921-928.
- Koch, A. E., P. J. Polverini, S. L. Kunkel, L. A. Harlow, L. A. DePietro, V. N. Elnor, S. G. Elnor, and R. M. Strieter. 1992. Interleukin-8 as a macrophage-derived mediator of angiogenesis. *Science (Wash. DC)*. 258:1798-1801.
- Gravallese, E. M., J. M. Darling, A. L. Ladd, J. N. Katz, and L. H. Glimcher. 1991. In situ hybridization studies of stromelysin and collagenase messenger RNA expression in rheumatoid synovium. *Arthritis Rheum.* 34:1076-1084.
- McCachren, S. S., B. F. Haynes, and J. E. Nield. 1990. Localization of collagenase mRNA in rheumatoid arthritis synovium by in situ hybridization histochemistry. *J. Clin. Immunol.* 10:19-24.
- Dayer, J.-M., B. Beutler, and A. Cerami. 1985. Cachectin/tumor necrosis factor stimulates collagenase and prostaglandin E<sub>2</sub> production by human synovial cells and dermal fibroblasts. *J. Exp. Med.* 162:2163-2168.
- DeForge, L. E., J. S. Kenney, M. L. Jones, J. S. Warren, and D. G. Remick. 1992. Biphasic production of IL-8 in lipopolysaccharide (LPS)-stimulated human whole blood. *J. Immunol.* 148:2133-2141.
- Vassalli, P. 1992. The pathophysiology of tumor necrosis factors. *Annu. Rev. Immunol.* 10:411-452.
- Webb, D. S. A., Y. Shimizu, G. A. Van Seventer, S. Shaw, and T. L.

- Gerrard. 1990. LFA-3, CD44, and CD45: physiologic triggers of human monocyte TNF and IL-1 release. *Science (Wash. DC)*. 249:1295-1297.
14. Yorochko, A. D., D. Y. Liu, D. Eierman, and S. Haskill. 1992. Integrins as a primary signal transduction molecule regulating monocyte immediate-early gene induction. *Proc. Natl. Acad. Sci. USA*. 89:9034-9038.
15. Economou, J. S., K. Rhoades, R. Essner, W. H. McBride, J. C. Gasson, and D. L. Morton. 1989. Genetic analysis of the human tumor necrosis factor  $\alpha$ /cachectin promoter region in a macrophage cell line. *J. Exp. Med.* 170:321-326.
16. Goldfeld, A. E., D. Doyle, and T. Maniatis. 1990. Human tumor necrosis factor  $\alpha$  gene regulation by virus and lipopolysaccharide. *Proc. Natl. Acad. Sci. USA*. 87:9769-9773.
17. Goldfeld, A. E., P. G. McCaffrey, J. L. Strominger, and A. Rao. 1993. Identification of a novel cyclosporin-sensitive element in the human tumor necrosis factor  $\alpha$  gene promoter. *J. Exp. Med.* 178:1365-1379.
18. Goldfeld, A. E., J. L. Strominger, and C. Doyle. 1991. Human tumor necrosis factor  $\alpha$  gene regulation in phorbol ester stimulated T and B cell lines. *J. Exp. Med.* 174:73-81.
19. Hensel, G., A. Meichle, K. Pfizenmaier, and M. Kronke. 1989. PMA-responsive 5' flanking sequences of the human TNF gene. *Lymphokine Res.* 8:347-351.
20. Leitman, D. C., E. R. Mackow, T. Williams, J. D. Baxter, and B. L. West. 1992. The core promoter region of the tumor necrosis factor  $\alpha$  gene confers phorbol ester responsiveness to upstream transcriptional activators. *Mol. Cell. Biol.* 12:1352-1356.
21. Leitman, D. C., R. C. J. Ribeiro, E. R. Mackow, J. D. Baxter, and B. L. West. 1991. Identification of a tumor necrosis factor-responsive element in the tumor necrosis factor  $\alpha$  gene. *J. Biol. Chem.* 266:9343-9346.
22. Rhoades, K. L., S. H. Golub, and J. S. Economou. 1992. The regulation of the human tumor necrosis factor  $\alpha$  promoter region in macrophage, T cell, and B cell lines. *J. Biol. Chem.* 267:22102-22107.
23. Cao, Z., R. M. Umek, and S. L. McKnight. 1991. Regulated expression of three C/EBP isoforms during adipose conversion of 3T3-L1 cells. *Genes & Dev.* 5:1538-1552.
24. Akira, S., H. Isshiki, T. Sugita, O. Tanabe, S. Kinoshita, Y. Nishio, T. Nakajima, T. Hirano, and T. Kishimoto. 1990. A nuclear factor for IL-6 expression (NF-IL6) is a member of a C/EBP family. *EMBO (Eur. Mol. Biol. Organ.) J.* 9:1897-1906.
25. Haas, J. G., M. Strobel, A. Leutz, P. Wendelga, C. Muller, E. Sterneck, G. Riethmuller, and H. W. Loms Ziegler-Heitbrock. 1992. Constitutive monocyte-restricted activity of NF-M, a nuclear factor that binds to a C/EBP motif. *J. Immunol.* 149:237-243.
26. LeClair, K. P., M. A. Blonar, and P. A. Sharp. 1992. The p50 subunit of NF- $\kappa$ B associates with the NF-IL6 transcription factor. *Proc. Natl. Acad. Sci. USA*. 89:8145-8149.
27. Ness, S. A., E. Kowenz-Leutz, T. Casini, T. Graf, and A. Leutz. 1993. Myb and MF-M: combinatorial activators of myeloid genes in heterologous cell types. *Genes & Dev.* 7:749-759.
28. Nishio, Y., H. Isshiki, T. Kishimoto, and S. Akira. 1993. A nuclear factor for interleukin-6 expression (NF-IL6) and the glucocorticoid receptor synergistically activate transcription of the rat  $\alpha$ 1-acid glycoprotein gene via direct protein-protein interaction. *Mol. Cell. Biol.* 13:1854-1862.
29. Stein, B., P. C. Cogswell, and A. S. Baldwin, Jr. 1993. Functional and physical associations between NF- $\kappa$ B and C/EBP family members: a rel domain-bZIP interaction. *Mol. Cell. Biol.* 13:3964-3974.
30. Grove, M., and M. Plumb. 1993. C/EBP, NF- $\kappa$ B, and c-Ets family members and transcriptional regulation of the cell-specific and inducible macrophage inflammatory protein 1a immediate-early gene. *Mol. Cell. Biol.* 13:5276-5289.
31. Katz, S., E. Kowenz-Leutz, C. Muller, K. Meese, S. A. Ness, and A. Leutz. 1993. The NF-M transcription factor is related to C/EBP $\beta$  and plays a role in signal transduction and leukemogenesis of avian myelomonocytic cells. *EMBO (Eur. Mol. Biol. Organ.) J.* 12:1321-1332.
32. Nomiyama, H., K. Keishima, K. Hirokawa, T. Hattori, K. Takatsuki, and R. Miura. 1993. Characterization of cytokine LD78 gene promoters: positive and negative transcriptional factors bind to a negative regulatory element common to LD78, interleukin-3, and granulocyte-macrophage colony-stimulating factor gene promoters. *Mol. Cell. Biol.* 13:2787-2801.
33. Shirakawa, F., K. Saito, C. A. Bonagura, D. L. Galson, M. J. Fenton, A. C. Webb, and P. E. Auron. 1993. The human prointerleukin 1 $\beta$  gene requires DNA sequences both proximal and distal to the transcription start site for tissue-specific induction. *Mol. Cell. Biol.* 13:1332-1344.
34. Stein, B., and A. S. Baldwin, Jr. 1993. Distinct mechanisms for regulation of the interleukin-8 gene involve synergism and cooperativity between C/EBP and NF- $\kappa$ B. *Mol. Cell. Biol.* 13:7191-7198.
35. Zhang, Y., and W. N. Rom. 1993. Regulation of the interleukin-1 $\beta$  (IL-1 $\beta$ ) gene by mycobacteria components and lipopolysaccharide is mediated by two nuclear factor-IL6 motifs. *Mol. Cell. Biol.* 13:3831-3837.
36. Schrieber, E., P. Matthias, M. M. Muller, and W. Schaffner. 1989. Rapid detection of octamer binding proteins with "mini-extracts" prepared from a small number of cells. *Nucleic Acids Res.* 17:6419.
37. Pomerantz, R. J., M. B. Feinberg, D. Trono, and D. Baltimore. 1990. Lipopolysaccharide is a potent monocyte/macrophage-specific stimulator of human immunodeficiency virus type 1 expression. *J. Exp. Med.* 172:253-261.
38. Bohmann, D., W. Keller, T. Dale, H. R. Scholer, G. Tabb, and I. Mattaj. 1987. A transcription factor which binds to the enhancers of SV40, immunoglobulin heavy chain and U2 snRNA genes. *Nature (Lond.)*. 325:268-272.
39. Trautwein, C., C. Caelles, P. van der Geer, T. Hunter, M. Karin, and M. Chojkier. 1993. Transactivation by NF-IL6/LAP is enhanced by phosphorylation of its activation domain. *Nature (Lond.)*. 364:544-547.
40. Natsuka, S., S. Akira, Y. Nishio, S. Hashimoto, T. Sugita, H. Isshiki, and T. Kishimoto. 1992. Macrophage differentiation-specific expression of NF-IL6, a transcription factor for interleukin-6. *Blood*. 79:460-466.
41. Scott, L. M., C. I. Civin, P. Rorth, and A. D. Friedman. 1992. A novel temporal expression pattern of three C/EBP family members in differentiating myelomonocytic cells. *Blood*. 80:1725-1735.
42. Williams, S. C., C. A. Cantwell, and P. F. Johnson. 1991. A family of C/EBP-related proteins capable of forming covalently linked leucine zipper dimers in vitro. *Genes & Dev.* 5:1553-1567.
43. Sterneck, E., C. Blattner, T. Graf, and A. Leutz. 1992. Structure of the chicken myelomonocytic growth factor gene and specific activation of its promoter in avian myelomonocytic cells by protein kinases. *Mol. Cell. Biol.* 12:1728-1735.
44. Chu, C. Q., M. Field, M. Feldmann, and R. N. Maini. 1991. Localization of tumor necrosis factor  $\alpha$  in synovial tissues and at the cartilage-pannus junction in patients with rheumatoid arthritis. *Arthritis Rheum.* 34:1125-1132.
45. MacNaul, K. L., N. I. Hutchinson, J. N. Parson, E. K. Bayne, and M. J. Tocci. 1990. Analysis of IL-1 and TNF- $\alpha$  gene expression in human rheumatoid synoviocytes and normal monocytes by in situ hybridization. *J. Immunol.* 145:4154-4166.
46. Matsusaka, T., K. Fujikawa, Y. Nishio, N. Mukaida, K. Matsushima, T. Kishimoto, and S. Akira. 1993. Transcription factors NF-IL6 and NF- $\kappa$ B synergistically activate transcription of the inflammatory cytokines, interleukin 6 and interleukin 8. *Proc. Natl. Acad. Sci. USA*. 90:10193-10197.
47. Angel, P., M. Imagawa, R. Chiu, B. Stein, R. J. Imbra, H. J. Rahmsdorf, C. Jonat, P. Herrlich, and M. Karin. 1987. Phorbol ester-inducible genes contain a common cis element recognized by a TPA-modulated trans-activating factor. *Cell*. 49:729-739.
48. Beg, A. A., T. S. Finco, P. V. Nantermet, and A. S. Baldwin, Jr. 1993. Tumor necrosis factor and interleukin-1 lead to phosphorylation and loss of I $\kappa$ B $\alpha$ : a mechanism for NF- $\kappa$ B activation. *Mol. Cell. Biol.* 13:3301-3310.
49. Hsu, W., T. K. Kerppola, P.-L. Chen, T. Curran, and S. Chen-Kiang. 1994. Fos and Jun repress transcription activation by NF-IL6 through association at the basic zipper region. *Mol. Cell. Biol.* 14:268-276.