

Preoperative comparison of three anticholinergic drug scales in older adult patients and development of postoperative delirium: a prospective observational study

Drugs & Aging

Maria Heinrich, Anika Müller, Andela Cvijan, Rudolf Mörgeli, Jochen Kruppa,

Georg Winterer, Arjen J.C. Slooter, Claudia D. Spies

on behalf of the BioCog Consortium

Corresponding Author

Claudia Spies, MD

Full Professor of Anesthesiology and Intensive Care Medicine

Department of Anesthesiology and Operative Intensive Care Medicine (CCM, CVK)

Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health

Augustenburger Platz 1, D-13353 Berlin, Germany

Email: claudia.spies@charite.de

Table S1: Overview of surgical sites (n=837)

Surgical Site	Frequency (Percentage)
Nervous System	41 (4.9%)
Endocrine Glands	10 (1.2%)
Eyes	31 (3.7%)
Ears	8 (1%)
Nose and the Paranasal Sinuses	15 (1.8%)
Face and in the Oral Cavity	22 (2.6%)
Pharynx, Larynx and Trachea	5 (0.6%)
Lung and Bronchus	19 (2.3%)
Heart	42 (5%)
Blood Vessels	15 (1.8%)
Lymphatic Vessels	19 (2.3%)
Digestive Tract	155 (18.5%)
Urinary Organs	47 (5.6%)
Male Genital Organs	52 (6.2%)
Female Genital Organs	51 (6.1%)
Jaw and Craniofacial Bone	14 (1.7%)
Locomotor System Organs	233 (27.8%)
Mamma	11 (1.3%)
Skin and Underlying Tissue	17 (2%)
Diagnostic Procedures (in general anesthesia)	26 (3.1%)