

**Repository of the Max Delbrück Center for Molecular Medicine (MDC)
in the Helmholtz Association**

<https://edoc.mdc-berlin.de/20218/>

**The ART of bringing extinction to a freeze - history and future of species
conservation, exemplified by rhinos.**

Hildebrandt T.B., Hermes R., Goeritz F., Appeltant R., Colleoni S., de Mori B., Diecke S., Drukker M., Galli C., Hayashi K., Lazzari G., Loi P., Payne J., Renfree M., Seet S., Stejskal J., Swegen A., Williams S.A., Zainuddin Z.Z., Holtze S.

This is a copy of the final article (publisher's version) which was originally published with a CC-BY-NC-ND 4.0 license in:

Theriogenology
2021 JUL 15 ; 169: 76-88
2021 APR 18 (first published online)
doi: [10.1016/j.theriogenology.2021.04.006](https://doi.org/10.1016/j.theriogenology.2021.04.006)

Publisher: [Elsevier](#)



Copyright © 2022 The Authors. Published by Elsevier Inc. This article is available under the [Creative Commons CC-BY-NC-ND license](#) and permits non-commercial use of the work as published, without adaptation or alteration provided the work is fully attributed.



The ART of bringing extinction to a freeze – History and future of species conservation, exemplified by rhinos



Thomas B. Hildebrandt ^a, Robert Hermes ^a, Frank Goeritz ^a, Ruth Appeltant ^b, Silvia Colleoni ^c, Barbara de Mori ^{d,e}, Sebastian Diecke ^{f,g}, Micha Drukker ^h, C. Galli ^{c,i}, K. Hayashi ^j, G. Lazzari ^c, Pasqualino Loi ^k, J. Payne ^l, Marilyn Renfree ^m, Steven Seet ^a, Jan Stejskal ⁿ, A. Swegen ^b, Suzannah A. Williams ^b, Zainal Zahari Zainuddin ^l, Susanne Holtze ^{a,*}

^a The Leibniz Institute for Zoo and Wildlife Research, Alfred-Kowalke-Straße 17, 10315, Berlin, Germany

^b Nuffield Department of Women's & Reproductive Health, University of Oxford, John Radcliffe Hospital, Oxford, OX3 9DU, United Kingdom

^c Avantea srl, Laboratory of Reproductive Technologies, Via Porcellasco 7/F, 26100, Cremona, Italy

^d Department of Comparative Biomedicine and Food Science, University of Padua, 35020, Padua, Italy

^e Ethics Laboratory for Veterinary Medicine, Conservation and Animal Welfare, University of Padua, 35020, Padua, Italy

^f Technology Platform Pluripotent Stem Cells, Max Delbrück Center for Molecular Medicine in the Helmholtz Association (MDC), Robert-Rössle-Str. 10, 13092 Berlin, Germany

^g Berlin Institute of Health, Anna-Louisa-Karsch-Straße 2, 10178 Berlin, Germany

^h Institute of Stem Cell Research, German Research Center for Environmental Health, Helmholtz Center Munich, Ingolstaedter Landstraße 1, 85764, Neuherberg, Germany

ⁱ Fondazione Avantea, 26100, Cremona, Italy

^j Faculty of Medical Sciences, Kyushu University, Maidashi 3-1-1, Higashi-ku, Fukuoka, 812-8582, Japan

^k Faculty of Veterinary Medicine, University of Teramo, Campus Coste San Agostino, 64100, Teramo, Italy

^l Borneo Rhino Alliance, Sabah, Malaysia

^m School of BioSciences, The University of Melbourne, Victoria, 3010, Australia

ⁿ Zoo Dvůr Králové, Štefánikova 1029, 544 01, Dvůr Králové nad Labem, Czech Republic

ARTICLE INFO

Article history:

Received 31 October 2020

Received in revised form

12 April 2021

Accepted 14 April 2021

Available online 18 April 2021

Keywords:

Cryopreservation

Self-sustaining population

Stem cell technologies

In-vitro-gametogenesis

Biobanking

ABSTRACT

The ongoing mass extinction of animal species at an unprecedented rate is largely caused by human activities. Progressive habitat destruction and fragmentation is resulting in accelerated loss of biodiversity on a global scale. Over decades, captive breeding programs of non-domestic species were characterized by efforts to optimize species-specific husbandry, to increase studbook-based animal exchange, and to improve enclosure designs. To counter the ongoing dramatic loss of biodiversity, new approaches are warranted. Recently, new ideas, particularly the application of assisted reproduction technologies (ART), have been incorporated into classical zoo breeding programs. These technologies include semen and oocyte collection, artificial insemination, and in-vitro embryo generation. More futuristic ideas of advanced ART (aART) implement recent advances in biotechnology and stem-cell related approaches such as cloning, inner cell mass transfer (ICM), and the stem-cell-associated techniques (SCAT) for the generation of gametes and ultimately embryos of highly endangered species, such as the northern white rhinoceros (*Ceratotherium simum cottoni*) of which only two female individuals are left. Both, ART and aART greatly depend on and benefit from the rapidly evolving cryopreservation techniques and bio-banking not only of genetic, but also of viable cellular materials suitable for the generation of induced pluripotent stem cells (iPSC). The availability of cryopreserved materials bridges gaps in time and space, thereby optimizing the available genetic variability and enhancing the chance to restore viable populations.

© 2021 Published by Elsevier Inc.

* Corresponding author.

E-mail address: holtze@izw-berlin.de (S. Holtze).

1. Introduction

Cryopreservation in liquid nitrogen at $-196\text{ }^{\circ}\text{C}$ brings most biological processes to a halt, thereby greatly expanding the possibilities of assisted reproduction technologies (ART). The visionary physician and researcher, Kurt Benirschke, in 1975 created the “Frozen Zoo®”, the first large-scale systematic cryobank for blood products, DNA samples, tissue, cells and, reproductive material of exotic species at the San Diego Zoo. In anticipation of future technologies, he already intended to extend biological knowledge and to sustain biodiversity, stating that “You must collect things for reasons you don’t yet understand.” [1]. This collection has expanded since, counting 10,000 cell lines in 2020 [2]. Indeed, biotechnological possibilities have greatly expanded, opening a wealth of new tools and possibilities, many of which are still being implemented or are at the boundaries of our current imagination.

The Anthropocene – the human epoch – is characterized by the dominance of *Homo sapiens* over the planet. Overexploitation of limited natural resources bears the risk of significant negative economical and societal consequences [3] as we fundamentally depend not only on their steady supply but also on the invaluable ‘ecosystem services’ which intact ecosystems provide [4]. The current unprecedented rate of species extinction is estimated at 100 to 1000 times higher than natural background rates [5,6] (Fig. 1a and b). This earth’s sixth great extinction event has already driven 22% of all mammalian species close to being lost forever. Ecological interactions are extremely complex and intertwined so that the loss of a single species may have much larger implications than we can foresee (so-called vortex-effect [7]). In our own interest [5,8], it becomes more and more pressing to contain and ideally reverse this development by protecting ecosystems, species and genetic variability.

Important scientific advances have been achieved in the recent past and will continue to open new pathways for the future of biodiversity conservation. After summarizing the historical, present and future possibilities, limitations, and success stories of species conservation, we will explore the new possibilities arising from advancing cryopreservation methods. Many important examples are connected with rhinoceros species, which grants them an inherent focus in our review; nevertheless – wherever relevant and possible – we attempt to adequately mention examples across all animal taxa.

2. Species conservation strategies

The most fundamental level of species conservation is the

protection of natural resources, as habitat loss or degradation for logging and agriculture presents the greatest current threat to biodiversity [9]. Starting in the late 18th century and encouraged by scientists such as Alexander von Humboldt [10], the Yellowstone National Park became the first public protected area (Fig. 2a and b). Nevertheless, large viable habitats are being emptied of wildlife by hunting [11,12], pollution [13], and by invasive species [14]. In many instances now, however, restoring habitat, stopping poaching and removing pollution and invasive species will not solve the problem because remaining clusters of wild animals are too small to be viable [15]. One of the earliest species conservation success stories concerns the southern white rhinoceros (SWR, *Ceratotherium simum*) which recovered from near extinction [16], through hunting bans and habitat protection [17], as well as strategic translocation [18]. Translocations have since become an important tool in wildlife conservation [19–21]. This success renders the SWR – despite lately declining numbers [22,23] – the most abundant rhinoceros species today. In contrast, the sister taxon northern white rhinoceros (NWR, *Ceratotherium simum cottoni*) has dwindled to only two female individuals, as poaching for rhino horn [24,25], leaves the Rhinocerotidae one of the most threatened mammalian families [26,27].

Insurance populations and ex-situ breeding by zoological institutions, crucial to species conservation plans [28,29], have led to at least 13 to 19 well-documented success stories among vertebrates [30–33], comprising the European bison (*Bos bonasus*) [34] (Fig. 2c), the Przewalski horse (*Equus ferus przewalskii*) [35] (Fig. 2c), the Arabian oryx (*Oryx leucoryx*) [36], and the red wolf (*Canis rufus*) [37,38]. Currently ongoing reintroduction efforts include for example the Scimitar-horned oryx (*Oryx dammah*) [39], and the Spix’s macaw (*Cyanopsitta spixii*) [40]. Despite considerable challenges and drawbacks, these encouraging examples illustrate the feasibility re-establishing wild populations that had already gone extinct.

To halt mass extinction and irreversible loss of keystone species before suitable solutions are found or habitats restored requires elaborate methods to enhance genetic management. Especially if complicated by small numbers of founder individuals, infertility due to old age, diseases, spatial distance between individuals or suboptimal husbandry, assisted reproduction technology (ART) [41–43] or advanced ART (aART) involving state-of-the-art stem cell and biotechnology (see Fig. 1c) will be necessary. Therein, making use of cryopreserved tissues, cells and reproductive materials is crucial, providing the opportunity to preserve and convey genetic diversity across time and space (Fig. 1d).




Fig. 1. Different scenarios of population development due to human impact.

Fig. 1. (a) Three different scenarios of the population development after human impact. (b) Reality for a large number of mammals, birds, and reptiles e.g. dodo, Steller’s sea cow since 1600s. (c) Interventions as demonstrated in (a)–(d) can help to rescue critically endangered species. (d) The role of cryopreservation if current technologies are not sufficient.




Fig. 2. Different levels of human intervention for species conservation.

Fig. 2. (a) Intact nature prior anthropogenic impact. (b) Situation after foundation of 1st national park (Yellow-Stone National Park) 1872. (c) Captive breeding for reintroduction programs (e.g. European bison, Przewalski horse). (d) Implementation of assisted reproduction (e.g. black footed ferret, Californian condor). (e) *BioRescue* – an international program for saving the northern white rhino from extinction will use SCAT for artificial gamete generation. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

2.1. Assisted reproduction technology (ART)

ART in wildlife species has been extensively reviewed elsewhere (e.g. Refs. [44–46]) and will be only briefly summarized here to illustrate the importance of cryopreservation. Many of the established techniques have been developed and optimized for livestock production. In-depth-knowledge of reproductive anatomy and physiology are an important prerequisite, but only available for approximately 250 species, mainly mammals and birds [47]. Taxonomically closely related domestic or less endangered model species may serve as a blue-print for developing suitable protocols for endangered wildlife species [42]. The first species to profit from the use of ART is the California condor (*Gymnogyps californianus*). Mainly due to lead poisoning it went extinct in the wild in 1987. Artificial incubation, foster parenting, or hand-rearing, state-of-the-art molecular genetics [48,49] and a ban of lead ammunition have restored a largely independent wild population [50] (Fig. 2d).

2.1.1. Hormone monitoring and administration

Endocrine monitoring of steroid metabolites is indispensable for monitoring ovarian cyclicity, infertility, seasonality of testicular activity, pregnancies, and for the determination of optimal timing for reproductive interventions, whereas monitoring corticosteroid levels can help to improve husbandry by assessing stress [51]. Hormonal stimulation is crucial to induce ovulation and manipulate reproductive activities (e.g., superovulation, or estrus synchronization in preparation for artificial insemination (AI), embryo transfer (ET), or contraception) and requires different protocols in

different taxa, e.g. Refs. [52–55]. Cryopreservation of the relevant samples allows for long-term collection, systematic evaluation, and safely shipping samples to laboratories.

2.1.2. Semen collection

Semen collection is fundamental to fertility assessment, ART, and gamete cryo-banking [56]. Relevant methods comprise massaging, phantom use, postcoital collection, e.g. Refs. [57,58], electroejaculation (EE) [59] urethral catheterization [56,60], or obtaining epididymal sperm by castration, biopsy, aspiration, or post mortem. Most frequently used methods in wildlife species are EE (from bats [61] to rhinoceros [62]), as well as urethral catheterization (mainly used in carnivores [63]). Semen collection opens up the possibility to move sperm instead of animals between facilities or between the wild and captivity. This permits for the inclusion of individuals into the gene pool that do not naturally mate owing to physical or behavioural handicaps, the use of aliquots for disease screening, or sex-sorting [64,65]. Semen cryopreservation removes the limitations imposed by generations and time, involving even long deceased individuals in reproduction [42,52,66].

2.1.3. Artificial insemination (AI)

AI is the most frequently used ART and has produced viable offspring in more than 50 wildlife species [45], comprising 14 bovid, seven cervid [53,67], three cetacean [68–72], two rhinoceros [73–75], and various wild cat species, including ocelot, Pallas's cat, fishing cat, sand cat, tiger, and clouded leopard [76,77]. Building on

knowledge of the poultry industry, AI supports the recovery of the whooping crane, peregrine falcon [78], houbara bustard [79,80], and Spix' Macaw [81], and was successful in numerous species of raptors, cranes, waterfowl, psittacines, and passerines [82].

Relevant contributions of AI to species recovery programs are limited to (i) the black-footed ferret (*Mustela nigripes*), by increasing genetic diversity while reducing inter-generational time [[83]]. Laparoscopic AI with fresh or frozen semen, even 20 years after their cryopreservation [42], has generated more than 8000 offspring and 4400 releases [83–85] (Fig. 2d). (ii) Giant panda (*Ailuropoda melanoleuca*) populations, decreasing due to habitat loss, poaching, bamboo flowering [86], low reproductive success [87] improved – besides by hand rearing of twin cubs [88] – due to AI with fresh and frozen-thawed semen [89], (Fig. 2d). (iii) AI in captive Asian (*Elephas maximus*) and African elephants (*Loxodonta africana*) [90–93] helps to avoid breeding-related transfers and increase genetic exchange, also with wild populations [94].

2.1.4. Ovum pick up (OPU)

The collection of oocytes (OPU) is more invasive, complex, and costly than that of semen. It is either performed timely post mortem, laparotomically, transcutaneously under ultrasound-guidance, transvaginally, or transrectally. OPU is regularly performed in domestic or laboratory species such as cattle, deer, horses, and macaques. It has been rarely applied in wildlife [45], where it is reported mainly for black [95] and Sumatran rhinoceros [96], and recently for the northern and southern white rhinoceros [52,96].

2.1.5. In vitro fertilization (IVF) and embryo transfer (ET)

In vitro fertilization (IVF) with fresh or frozen-thawed semen has been developed and optimized for humans, laboratory animals, and livestock [45] to increase the female genetic contribution to the gene pool. IVF is followed by in vitro culture of embryos and ET. Obstacles are oocyte or embryo retrieval, the vulnerability of the large oocyte to cryo-damage, more complex handling and culture, and costly equipment. Moreover, synchronization of embryo development and facilitating foeto-maternal recognition is a major obstacle to overcome. Exact knowledge of reproductive cycles is crucial for successful reimplantation. Consequently, as compared to AI, relatively few live offspring have been produced following IVF-ET.

Since the first successful mammalian embryo transfer in the rabbit in 1890 [97], the first non-domestic species to give birth to live offspring after ET was the baboon (*Papio cynocephalus*) [98]. Successful interspecies embryo transfers were achieved from eland (*Tragelaphus oryx*) and gaur (*Bos gaurus*) to domestic cow (*Bos Taurus*) [99,100], and from bongo (*Tragelaphus euryceros*) to eland [101]. Despite standard use in domestic species, and despite further successes in IVF and ET of felid [102–107], bovine [108], deer [109–111], and primate species as models for human ART [112], this technique has not played a major role in the genetic management of mammalian wildlife species so far [113]. Successful hormonally induced gamete harvesting, *in-vitro* fertilization, and embryo development have generated large numbers of viable amphibian offspring, including the endangered Wyoming toad (*Bufo baxteri*) and Mississippi gopher frogs (*Rana sevososa*), which were released to the wild [114,115].

2.2. Advanced assisted reproduction technology (aART)

The term advanced assisted reproduction technology was coined in 2004 [116] and refers to methods that require extensive laboratory equipment and expertise which lie beyond those needed for the more “classical” ART methods. While some of these advanced techniques have been available for decades, recent

developments have opened up new pathways which may present the only hope for critically endangered species such as the northern white rhinoceros or the Bornean subspecies of the Sumatran rhinoceros (*Dicerorhinus sumatrensis harrissoni*). In both cases the effective founder population is already too small for established species recovery programs. Therefore, the incorporation of new cellular resources in combination with cryopreservation, such as tissue biopsies, blood samples, or fibroblast cultures can open up a new avenue in conservation. Contrary to cloning, this cellular material may be subjected to stem-cell associated techniques (SCAT), yielding artificial gamete cultures and thereby widen the genetic pool by including – with the help of cryopreservation – samples from deceased or completely infertile individuals [117,118] (Fig. 2e). A strategic roadmap was outlined in “Rewinding the process of mammalian extinction” [66]. Along these lines, high quality hybrid embryos from SWR and NWR gametes [52], as well as pure NWR blastocysts [119] were successfully generated. The blastocysts were also the basis for the establishment of two embryonic stem-cell lines. In addition, we were able to produce integration-free naive and primed iPSC-like cells derived from cryopreserved fibroblast cultures of a deceased northern white rhino. First steps towards the transformation of these primordial germ cells (PGC) were very promising. They expressed typical PGC marker genes *Blimp*, *Stella*, *Sox17*, and *Oct4* [120]. These achievements provide the basis for the second phase: The production of artificial oocytes and spermatozoa originally derived from simple fibroblast cultures.

2.2.1. Intracytoplasmic sperm injection (ICSI)

Intracytoplasmic sperm injection (ICSI), the injection of sperm into the egg cytoplasm through a micropipette, is extensively used in humans [121], but remains rare in domestic and livestock reproduction owing to low success rates, and the requirement of expensive equipment, and skills [122,123] ICSI nevertheless offers a great advantage when semen characteristics are insufficient for IVF. Attempts of implementing ET and in southern and northern white and Sumatran rhinoceros following ICSI are currently undertaken [52,66].

2.2.2. Somatic cell nuclear transfer (SCNT)

SCNT, alias somatic cloning, has suffered a substantial loss in reputation owing to a range of problems, such as early and late abortions, compromised immune systems, circulatory and respiratory problems, and a high rate of foetal death, probably primarily mediated by atypical epigenetic re-programming [124].

Nevertheless, it has been successfully performed not only in domestic species such as cattle, horse, pig, and sheep [125], but also in non-domestic red deer (*Cervus elaphus*) [126] and cynomolgus monkey (*Macaca fascicularis*) [127]. Interspecies somatic nuclear transfer has been achieved in several wildlife species such as the Gaur (*Bos gaurus* to *Bos taurus*) [128], including the endangered mouflon (*Ovis orientalis musimon* in *Ovis aries*) [129], African wildcat (*Felis silvestris lybica* in *Felis catus*) [130], grey wolf (*Canis lupus* in *Canis familiaris*) [131,132], sand cat (*Felis margarita* in *Felis catus*) [133], the extinct Pyrenean Ibex (*Capra pyrenaica* in *Capra hircus*) [134], and the Russian sturgeon (*Acipenser gueldenstaedtii* in *Acipenser ruthenus*) [135].

Although somatic cloning and genomic approaches in mammals may become a last desperate option for species conservation [136], it is not very efficient in generating live offspring and major technical and ecological challenges remain unsolved [113]. Cloning may, however, be an interesting option for non-mammalian vertebrates such as amphibians (in which the technique was pioneered) or species where breeding is threatened or that have even gone extinct [137].

2.2.3. Inner cell mass (ICM) exchange

The inner cell mass (ICM) of an early stage mammalian embryo contains the cells that will determine the developing organism, whereas the trophoblastic vesicle surrounding it gives rise to the placenta. Transferring the ICM of an endangered mammalian species into a trophoblastic vesicle derived from a surrogate female of a different species facilitates implantation and successful gestation of offspring of the endangered species. The foster mother's species will be selected to be taxonomically closely related, but less endangered. This is a highly experimental approach, which has so far been successfully applied between sheep and goat [138,139], and between sheep embryos [140], but may represent a viable option in the future – especially if combined with artificial gametes derived from induced pluripotent stem cells [46]. ICM exchange has not been attempted so far as a solution to the pressing problem of lacking reproductively healthy recipients in critically endangered species.

2.2.4. Stem cell-associated techniques (SCAT)

aART in combination with techniques for generating gametes from stem cells provide a new conceptual strategy for saving critically endangered or practically extinct species. This ambitious and novel approach is currently being developed for rhinoceros species [52,66]. In the future it may serve as a blue-print for applying the potential of in-vitro-derived gametes [139–145] created from induced pluripotent stem cells to the conservation of further species on the brink of extinction.

While SCNT generates copies of existing genotypes, recent advances in stem cell technology [120] have opened a new promising path, using live cells to establish induced pluripotent stem cell (iPSC)-derived gametes. As this approach comprises meiosis, it is capable of generating an enormous variety of new genotypes. By crossbreeding arbitrary individuals (theoretically, iPSC of male donors can be used to produce oocytes, which requires silencing of Y-chromosome-linked gene(s) [142], a pedigree can be designed in the Petri dish to optimally exploit the available genetic diversity. So far, viable offspring from iPSC-derived gametes has been generated only in the mouse (*Mus musculus*; [141,143–145], but efforts are currently undertaken to extend this approach for saving highly endangered taxa, specifically the northern white rhinoceros (*Ceratotherium simum cottoni*). iPSC have been successfully established for several domestic and laboratory species [146], but also for several wildlife species such as the quail [147], several feline species [148], e.g. the endangered snow leopard [149] and for the critically endangered northern white rhinoceros [150,151] (Fig. 2e).

2.3. Cryopreservation

Cryopreservation crucially enhances the possibilities of ART by rendering the use of biological materials independent of time and space.

2.3.1. Cryopreserved materials and associated methods

Various materials can be preserved using cryopreservation for extended, potentially indefinite, periods of time [152]. Different methods have been developed for the long-term cryopreservation of biological samples, with a strong focus on vertebrate species, whereas many other taxa remain unstudied. The major difficulty to overcome is vulnerability to cryo-damage, which depends on cell membrane composition, its permeability regarding both, water and the cryoprotectant, cryoprotectant toxicity, tolerance to osmotic changes, and resistance to cooling and freezing temperatures [153]. Thus, suitable protocols differ substantially between species and material.

Blood samples are suitable for biochemical analyses, as well as

for medical inquiries, and may serve as a source of DNA for molecular biological investigations, and – if adequately preserved – also as a source of viable lymphocytes suitable for iPSC protocols.

Tissue samples e.g. of liver or spleen are useful for the extraction of enzymes, and genomic and mitochondrial DNA, and for establishing primary cell cultures. Viable cells for subsequent cell culture can be preserved by freezing tissues with an adequate cryoprotectant such as glycerol or dimethylsulfoxide (DMSO).

Cell culture-based methods have gained considerable importance in basic research and biomedical applications over the past decades. Besides storing nuclear genetic information, cells further contain viable cell organelles and represent an amplifiable source of biological material. Over the past decades, cell lines from a wide range of different taxa [152] have been successfully established and banked. In the context of ART, owing to the development of advanced methods such as SCNT, ICM and in vitro gameteogenesis, cultured and cryopreserved cells will play a crucial role in maintaining and improving population viability of rare and endangered species.

Cell culture is nevertheless expensive and requires well-equipped laboratory space, sterile working conditions and skilled personnel. The most accessible samples are fibrous tissue such as skin or gingiva, and the less abundant lymphocytes that may be recovered from blood; viable cells have even been successfully isolated also from ejaculate, milk [154], and feces [155]. Skin biopsies can be obtained at zoological institutions, e.g. from ear-notches, or opportunistically during handling for quarantine, veterinary interventions such as castrations, surgery, transport, or ultimately during necropsy. If adequately cooled and stored, such tissue may remain viable for several days before further processing and may be either cryopreserved or cultured immediately. Primary cells can be obtained by centrifugation of blood, by mechanical disaggregation or enzymatic digestion of tissues, or by explant culture [156]. The hence cultured cell lines can be frozen, protected with glycerol or DMSO, and stored in liquid nitrogen containers for many years [1]. Such cryopreserved tissues and cell lines provide a viable and expandable source of genetic material and living cells that offer manifold possibilities for molecular and basic research.

Growing fibroblasts to sufficient numbers requires time, which increases material costs and risk of contamination, and repeated passaging of cell cultures may result in decreased viability. Unless primary cells are derived from neoplastic tissues or immortalized by mutation or specific intended modifications [157], their lifespan is limited due to undergoing senescence and ceasing to proliferate after a certain number of divisions [158]. Another limitation is the accumulation of mutations and chromosomal aberrations (aneuploidy) that may arise under culture conditions [159]. Therefore, for somatic cell-based technologies it is crucial to closely monitor the quality of the original cell lines and ensure genome integrity including chromosomal stability and the absence of relevant epigenetic alterations during prolonged culture and differentiation, which occur randomly and unpredictably [160]. Cell lines differ in quality based on culture conditions, age of the cell material donor, number of passage and many more factors, rendering quality control crucial to improve the observed low rates of reprogramming following SCNT and iPSC. Much work needs to be done for banked cell lines to become a valuable resource for offspring production. Cell lines can be reprogrammed to iPSC [120], and – if exposed to suitable factors – their pluripotent nature permits them to differentiate into various tissues including gametes.

Reproductive material suitable for cryopreservation comprises gametes, embryos, gonadal tissues, as well as embryonal (ES) and induced pluripotent stem cells (iPSC), which are gaining importance in preserving species biodiversity, as the associated technology evolves. However, progress is hampered due to significant

physiological variations among species and lack of fundamental knowledge in germplasm cryobiology.

Spermatozoa of approximately 116 mammalian species have been cryopreserved, accounting for 2% of all mammals, from which in approximately 45 species live births have been achieved following AI [161] since the first successful cryopreservation in 1949 [162,163] and the first calf produced from cryopreserved semen [163]. This renders cryopreserved semen the most successful and practical resource for propagating endangered species. Cryopreserved sperm of genetically valuable individuals can be stored for decades as a backup for possible future losses in genetic diversity. Ram semen has been reported to sire viable offspring after 50 years of storage without any decrease in pregnancy rates compared to recently frozen semen [164]. Even if thawed sperm quality is poor, it may generate viable offspring using ICSI [165]. Two of the most successful recent conservation programs, the giant panda [166] and the black-footed ferret have incorporated AI using fresh and frozen-thawed semen [167]. For the black-footed-ferret, cryopreserved semen of six of the last 18 survivors was employed to produce eight live offspring by AI, some of which up to 20 years after initial cryopreservation. Their relevance is illustrated nicely by their significant contribution to genetic diversity and heterozygosity in the population, lowering measures of inbreeding by 5.8% [83].

In contrast to mammals, bird sperm cells are fragile and knowledge is limited, complicating semen collection and processing [78]. Due to poor outcomes of cryopreservation in cranes [168], AI was not implemented in crane genetic management. Frozen-thawed sperm has generated viable offspring in a few amphibians [169] and fish [170] species, but so far not in reptiles, in which only artificial insemination with chilled semen has been accomplished so far [171].

One obstacle to the broad application of sperm cryopreservation is the variability in sensitivity to cryoprotectants and low temperatures from species to species. Tolerance to glycerol ranges from less than 2% in mice to 6% in chinchillas [153,172]. Sperm cryopreservation protocols are continuously extended and improved [173], e.g. by slow-freezing [174], optimizing freezing of small [175] and large [176] volumes; directional freezing [167,177,178] and double-freezing [179]. Even more challenging are amphibian and fish studies, as their sperm remain immotile as long as they are in seminal plasma. Motility is induced and quickly exhausted when they are released into a lower osmolarity environment, in the case of frogs, in urine [54,180,181]. Thus, amphibian and fish cryopreservation protocols need to keep sperm inactivated during handling and storage by mimicking the testicular environment. Anuran sperm has been successfully cryopreserved following non-invasive collection via hormonal induction [54]. Lately, sperm of a variety of coral species has been cryopreserved for the in vitro production of larvae and restoration of reefs [182].

Oocytes are a scarce resource compared to sperm cells, and less accessible. They can be obtained only invasively or post mortem [183], and are more difficult to handle [153]. Therefore, they are underrepresented in cryobanks and oocytes of only a few species have been preserved so far using vitrification [45,184]. Due to their large size, structure and lipid-rich yolk composition, frozen-thawed oocytes are more vulnerable to cryo-damage and until now, have not yielded any offspring in wild mammalian, fish, or amphibian species [167]. Survival and function of frozen-thawed oocytes has improved with the development of minimum volume vitrification (MVB), exhibiting extremely fast cooling rates that solidifies the sample into a glass-like state, thereby avoiding harmful intra- and extracellular ice crystal formation [185]. Using this method, oocytes of four carnivore species (Mexican grey wolf [186], serval, and Pallas's cat [187], and lion [188]), one antelope (chousingha [189]),

and one marsupial (Tasmanian devil [190]) have been cryopreserved, but the developmental competence of the cryopreserved oocytes has only partially been reported.

Embryos of approximately 51 species (1% of mammals) have been cryopreserved, the success of which in terms of post-thaw viability and further development is not always reported [161]. Live births after ET of frozen-thawed embryos have been achieved in merely 25 species [161]. Similar to oocytes, differences in cryoprotection and freezability of fertilized eggs among different taxa will require considerable research. Vitrification [184,191] has been found to preserve the developmental potential of human embryos better than the earlier developed slow-freezing [192,193] or ultra-rapid freezing [194]. As of now, our bank contains few embryos with thus far unproven viability. Additionally, purebred SWR and purebred NWR embryos have been generated using harvested oocytes that have been fertilized with cryopreserved sperm of dead NWR males [52,66,195] and are currently stored in liquid nitrogen [119].

- Gonadal tissues are designed to mature functional gametes, but they also continue to contain gamete precursor cells of various different developmental stages. These spermatogenic and oogenic cells are another viable resource for obtaining functional spermatozoa and oocytes [153]. As they lack a metaphase spindle, are smaller and metabolically less active compared to mature gametes and contain low amounts of lipids [196], they are more cryo-resistant. Hormonal treatment of the female to stimulate follicle development for OPU results in just a handful of oocytes for collection under general anaesthesia. Therefore, collecting oocytes by stimulating the ovary to develop them in vivo brings limited opportunities for successful population recovery. One approach is to harness the full potential of the ovary, which contains thousands of immature oocytes in primordial follicles – which can be obtained opportunistically during OPU or by using a needle-derived ovarian cortex biopsy – by developing an in vitro follicle culture method. The exceptional power of this technique is that, even when animals have died, their cryopreserved ovarian tissue can still lead to offspring. Establishing methods to culture and grow these follicles in the lab would vastly increase the chance of successful in vitro embryo production for endangered species. A first step in this process is to xenotransplant ovarian tissue into mice to verify the health and growth potential of collected frozen-thawed ovarian tissue (e.g. for endangered rhinoceros species, Fig. 3). By xenografting or in vitro culture of banked ovarian and testicular tissue, including that derived from neonatal and prepubertal individuals [196–198], mature gametes may be obtained, although this has not yet been achieved in non-domestic species.

Ovaries or ovarian tissue of a variety of mammalian species have been cryopreserved, e.g. of several felines [199,200] and the black-footed ferret (*Mustela nigripes*), amongst others [45].

Cryopreserved ovarian tissues from wombats, elephants, wallabies, and lions have been transplanted into immunodeficient mice, consistently resulting in the formation of morphologically normal secondary or antral follicles [200]. While it is still a long way to obtain viable mature oocytes for successful IVF, nevertheless, this approach represents an important step towards the inclusion of female gametes into ART.

For amphibians, direct cryopreservation of immature ovarian follicles may be a viable option, but would require methods such as xeno-transplantation to obtain mature, ovulated oocytes. For obtaining viable oocytes from cryo-preserved primordial germ cells, the generation of chimeras would be required to produce adults that can yield viable gametes; this may be a feasible future possibility also in fish and birds [55,167].

Comparable to ovarian tissue, in theory germ cells contained in testicular tissue can resume spermatogenesis to produce viable




Fig. 3. Xenotransplantation of rhinoceros ovarian tissue into mice to verify the health and growth potential of collected frozen-thawed ovarian tissue.

spermatozoa, even in tissue of neonatal and prepubertal individuals [183]. To achieve this, transplantation or grafting are needed. ICSI of sperm recovered from grafted cryopreserved testicular tissue has generated live offspring in the mouse and rabbit [201], pig [202,203], and rhesus macaque (*Macaca mulatta*) [204], in birds (e.g. quails, *Coturnix japonica*) [205], and in fishes, such as hatched larvae of the critically endangered cyprinid hommoroko (*Gnathopogon caeruleus*) [206]. Testicular tissue of various mammalian wildlife species has been cryopreserved [207] without reported successful generation of live offspring so far.

2.4. Cryobanks

Genome resource banking (GRB) is the systematic collection, storage, and redistribution of biomaterials in an organized, logistical, and secure manner, which together with associated genomic information, are essential for progression of biomedicine, health, and basic research. Especially animal germplasm (sperm, eggs, embryos, ovarian, and testicular tissues) employed in combination with ART offers great potential to decelerate the loss of gene diversity in captive populations.

Unfortunately, currently there are no suitable alternatives to cryobanking samples at preferably $-196\text{ }^{\circ}\text{C}$ in liquid nitrogen. The associated disadvantages comprise the risk of cross-contamination with pathogens in liquid nitrogen, danger of accidents, as well as high energetic costs and dependence on a constant energy supply. Alternatives such as dry biobanking by exsiccating samples via freeze- or vacuum-drying (i.e. lyophilization), have yielded some successes, such as fertilizing ovine oocytes with freeze-dried spermatozoa using ICSI [208]. However, reliable protocols are not established as of now to an extent that the return to biological activity after rehydration is ensured, but the technology may offer a much cheaper and more stable solution for biodiversity storage in the future [161].

Biobanks offer countless possibilities. Their main advantage is that they maintain genetic variability across time and space. Cryopreservation of viable cells can keep otherwise extinct species suspended and buy time until the methodology and technology are in place to bring the species back to regular existence [209]. Cryobanks of endangered animal species represent an extremely valuable backup of today's biodiversity. Original genetic material can be maintained without removing genetically valuable individuals from the wild, and decrease the interval between generations [165,167]. Naturally, they are also confronted with constraints of space and resources given that storage tanks, maintenance and regular liquid nitrogen supply are costly. Long-term commitment of a cell bank sponsor is needed to secure the perpetuity of the collection, and mirroring of the bank, preferably in geographically different sites is highly recommended [1]. Optimally, an automated, digitised sample storage system is implemented. Whereas the majority of existent cryobanks established at zoos aim at preserving mainly genetic material or other non-alive items, extending this classical concept to living materials, and using advanced cryopreservation techniques, cellular methods and state-of-the-art stem cell technologies will open up many novel possibilities.

Preservation of genetic material from wildlife is mostly derived from zoos due to the ease of access and include many species that are threatened, extinct in the wild, or completely extinct. The oldest cryobank for wildlife-derived samples is the San Diego Frozen Zoo® with more than 10,000 cell lines across 11 mammalian orders, oocytes, sperm and embryos representing almost 1000 different taxa [1,2,152], established in 1975. In addition, the Frozen Ark Consortium [210,211], was founded. In the 1990s, the biobank at IZW Berlin was established with now more than 150 different living cell lines and tissues and reproductive material of approximately 250 different exotic species, amongst them cryopreserved semen from a total of 45 species [210,212]. Further initiatives include the

National Institute for Environmental Studies in Japan founded in 2002, containing cell lines of many avian and mammalian species, the amphibian ark (www.amphibianark.org), the Israel–German Ark of Life (IGAL), founded in 2016, that preserves tissues and cells of wildlife and exotic species from the largest wildlife clinic in the Middle East at Ramat Gan Safari, as well as the frozen Zoo in Australia (<http://www.australianfrozenzoo.org.au/>). Most recently, CryoArks [213] and in 2020, the biobank Nature’s SAFE (Saving Animals From Extinction) were established in the UK (<https://www.natures-safe.com/>). Unusually, Nature’s SAFE is an independent charity and is dedicated to collecting samples from endangered species with the view to regeneration when required. Several of these banks have been established when the generally accepted paradigm still excluded the possibility of reprogramming differentiated cells, thus before the recently arising possibilities could be foreseen. It therefore seems safe to assume that future applications will be just as much beyond our current imagination.

3. Discussion

At the current rate of species extinction, urgent action is needed to preserve as much biodiversity as possible. However, when recapitulating the initial and recent encouraging success stories of species recovery by measures ranging from simple habitat protection to the incorporation of increasing levels of assisted reproduction technology, and with the recent emergence of state-of-the-art stem cell biotechnology, we may gain hope for the future. The incorporation of increasingly sophisticated genetic and reproductive tools has offered us strategies for minimizing the occurrence of genetic diseases, as in the California condor [48], retain high genetic diversity in black-footed ferrets using frozen-thawed semen that had been stored for 20 years [83] and re-build viable species from population sizes that would otherwise doom the species to extinction (Fig. 1b–d).

Threats to nature may be increasing - but so are the opportunities and tools at our disposition.

This is illustrated best with the northern white rhinoceros (*Ceratotherium simum cottoni*), of which only two female individuals are left. Cryopreserved tissue samples, somatic cell lines, iPSCs, and spermatozoa of 12 (5.7) individuals have been stored [66] and have, so far, allowed us to create so far five blastocysts of a species with no male individual alive. Encouraging for the prospect of recovering the NWR from these available resources, the genomes of the preserved specimen show levels of heterozygosity comparable to the SWRs, with higher levels of genome-wide heterozygosity and slightly lower levels of autozygosity in the NWR compared to the SWR [214], sparking hope for this charismatic keystone species. We may still be in time to someday reintroduce this landscape architect back to its now empty home in the central African bushland.

Zoos and zoological research institutions are key players for conserving genetic variability and provide reliable access to valuable material. Sample collection would optimally be implemented into the routine of zoo veterinarian work. A global network of cell culture repositories is missing. Although the number of GRB has been increasing over the past decades, their integration into common strategies, a unified data base, and concerted contribution to the management of captive populations could be further extended. Accessible and interconnected databases are needed to combine the available range of samples of a given species, pooling the existing resources for successful conservation efforts and research, thereby lifting the impact for biodiversity preservation on a higher level. It is further highly desirable to render the collections safe by mirroring across several distant locations. A further important impediment to globally concerted research efforts and

international cooperation is grounded in the exchange of samples and specimens. The many national and international legislations and rules intended to prevent importation of exotic diseases and misuse – the Nagoya protocol, CITES, TRACES, national export- and import permits, to name a few – can render sample exchange very costly in terms of time and money, sometimes almost impossible. With the help of governments and legislation, conservation research should be facilitated in this respect.

4. Conclusion

Cryopreservation is an indispensable tool. In combination with assisted reproduction technology, it can become an integral part of successful species conservation management. The entire information of an organism is contained in each of its cells and this information can be preserved in liquid nitrogen for decades or even centuries. Recent advances in stem-cell technology may allow for reprogramming these cells to gametes, ultimately resulting in entirely new individuals, conceived from the material of a small skin biopsy. Many obstacles are still in our way before we can apply these techniques to one, and a hopefully further on to a broad range of animals. As of now, we will not be able prevent the extinction of many populations; but by preserving cells of as many species as possible, we may be having the chance to bring them back in the future. Now, as much as in 1984 [1], we must go on and “collect things for reasons we don’t yet understand.”

Acknowledgements

This study was funded by the BMBF project “BioRescue” (01LC1902A), by the Nadace ĀEZ project PR21/46514, by Richard McLellan and Merck KGaA, Darmstadt, Germany to C.G., and by Fondation Hoffmann to S. A. Williams. The Authors like to thank the German Society of Cryobanks (GDK) for support of our conservation research. Thanks to Derek Ott and Jan Zwilling for carefully reading the manuscript.

References

- [1] Benirschke K. The frozen zoo concept. *Zoo Biol* 1984;3(4):325–8. <https://doi.org/10.1002/zoo.1430030405>.
- [2] San Diego Zoo Institute for Conservation Research. Frozen zoo. <https://institute.sandi-egozoo.org/resources/frozen-zoo>. [Accessed 10 October 2020].
- [3] Crutzen PJ. The “Anthropocene”. In: Ehlers E, Krafft T, editors. *Earth system science in the anthropocene*. Berlin: Springer; 2006. p. 13–8.
- [4] Ehrlich PR, Ehrlich AH. In: *Extinction: the causes and consequences of the disappearance of species*. vol. 1. Random House; 1981.
- [5] Ceballos G, Ehrlich PR, Barnosky AD, García A, Pringle RM, Palmer TM. Accelerated modern human-induced species losses: entering the sixth mass extinction. *Sci. Adv.* 2015;1(5). <https://doi.org/10.1126/sciadv.1400253>. e1400253.
- [6] Lamkin M, Miller AI. On the challenge of comparing contemporary and deep-time biological-extinction rates. *Bioscience* 2016;66(9):785–9. <https://doi.org/10.1093/biosci/biw088>.
- [7] Lacy RC. VORTEX: a computer simulation model for population viability analysis. *Wildl Res* 1993;20(1):45. <https://doi.org/10.1071/wr9930045>.
- [8] Ceballos G, Ehrlich PR. The misunderstood sixth mass extinction. *Science* 2018;360(6393):1080–1. <https://doi.org/10.1126/science.aau0191>.
- [9] Joppa LN, O’Connor B, Visconti P, Smith C, Geldmann J, Hoffmann M, et al. Big Data and biodiversity. Filling in biodiversity threat gaps. *Science* 2016;352(6284):416–8. <https://doi.org/10.1126/science.aaf3565>.
- [10] Editorial. Humboldt’s legacy. *Nat. Ecol. Evol.* 2019;3(9):1265–6. <https://doi.org/10.1038/s41559-019-0980-5>.
- [11] Gray TNE, Hughes AC, Laurance WF, Long B, Lynam AJ, O’Kelly H, et al. The wildlife snaring crisis: an insidious and pervasive threat to biodiversity in Southeast Asia. *Biodivers Conserv* 2018;27(4):1031–7. <https://doi.org/10.1007/s10531-017-1450-5>.
- [12] Ripple WJ, Abernethy K, Betts MG, Chapron G, Dirzo R, Galetti M, et al. Bushmeat hunting and extinction risk to the world’s mammals. *R. Soc. Open Sci.* 2016;3(10):160498. <https://doi.org/10.1098/rsos.160498>.
- [13] IUCN. The IUCN red list of threatened species.: version 2016-2. <http://www.iucnredlist.org>; 2016.

- [14] Tilman D, Clark M, Williams DR, Kimmel K, Polasky S, Packer C. Future threats to biodiversity and pathways to their prevention. *Nature* 2017;546(7656):73–81. <https://doi.org/10.1038/nature22900>.
- [15] Payne John. To Prevent Extinction of Endangered Large Mammals: Changing Perceptions and Practices, 1960–2020, and Proposals for a future approach. *Malayan Nature Journal* 2021:269–81. 81st Anniversary Special Issue.
- [16] Emslie RH, Brooks M. How many southern white rhinos were there? A response to Kees Rookmaaker. *PACHYDERM* 2002;33:100–1.
- [17] Rookmaaker K. The alleged population reduction of the southern white rhinoceros (*Ceratotherium simum simum*) and the successful recovery. *Säugetierkundliche Mitteilungen* 2000;45(2):55–70.
- [18] Player I. Translocation of white rhinoceros in South Africa. *Oryx* 1967;9(2):137–50. <https://doi.org/10.1017/S0030605300006165>.
- [19] Griffith B, Scott JM, Carpenter JW, Reed C. Translocation as a species conservation tool: status and strategy. *Science* 1989;245(4917):477–80. <https://doi.org/10.1126/science.245.4917.477>.
- [20] Berger-Tal O, Blumstein DT, Swaisgood RR. Conservation translocations: a review of common difficulties and promising directions. *Anim Conserv* 2020;23(2):121–31. <https://doi.org/10.1111/acv.12534>.
- [21] Amin R, Thomas K, Emslie RH, Foose TJ, van Strien N. An overview of the conservation status of and threats to rhinoceros species in the wild. *Int Zoo Yearbk* 2006;40(1):96–117. <https://doi.org/10.1111/j.1748-1090.2006.00096.x>.
- [22] SaveTheRhino.org. Rhino population figures. www.savetherhino.org/rhino-info/population-figures/. [Accessed 16 May 2015].
- [23] SaveTheRhino.org. Rhino population figures. www.savetherhino.org/rhino-info/population-figures/. [Accessed 28 October 2020].
- [24] Haas TC, Ferreira SM. Conservation risks: when will rhinos be extinct? *IEEE Trans. Cybern.* 2016;46(8):1721–34. <https://doi.org/10.1109/TCYB.2015.2470520>.
- [25] Traffic. TRAFFIC's engagement on African rhinoceros conservation and the global trade in rhinoceros horn: cambridge Conservation Biology. <http://www.traffic.org/rhinos/accesse>. [Accessed 1 December 2016].
- [26] Di Minin E, Laitila J, Montesino-Pouzols F, Leader-Williams N, Slotow R, Goodman PS, et al. Identification of policies for a sustainable legal trade in rhinoceros horn based on population projection and socioeconomic models. *Conserv Biol* 2015;29(2):545–55. <https://doi.org/10.1111/cobi.12412>.
- [27] Emslie RH, Milliken T, Talukdar B. African and Asian rhinoceroses – status, conservation and trade.: a report from the IUCN Species Survival Commission (IUCN/SSC) African and Asian rhino specialist groups and TRAFFIC to the CITES secretariat pursuant to resolution Conf. 9.14 (Rev. CoP15). <http://rhinos.org/wp-content/uploads/2015/07/final-cop16-rhino-rpt.pdf>; 2013.
- [28] Snyder NFR. Limitations of captive breeding in endangered species recovery. 1996.
- [29] Guidelines on the use of ex situ management for species conservation. Version 2.0. 2014.
- [30] Butchart SH, Stattersfield AJ, Collar NJ. How many bird extinctions have we prevented? *Oryx* 2006;40(3):266–78. <https://doi.org/10.1017/S0030605306000950>.
- [31] Gusset M, Dick G. 'Building a Future for Wildlife'? Evaluating the contribution of the world zoo and aquarium community to in situ conservation. *Int Zoo Yearbk* 2010;44(1):183–91. <https://doi.org/10.1111/j.1748-1090.2009.00101.x>.
- [32] Hoffman M, Hilton-Taylor C, Angulo A, Böhm M, Al ET, Cavanagh R. The impact of conservation on the status of the world's vertebrates. 2010.
- [33] Conde DA, Flessness N, Colchero F, Jones OR, Scheuerlein A. Conservation. An emerging role of zoos to conserve biodiversity. *Science* 2011;331(6023):1390–1. <https://doi.org/10.1126/science.1200674>.
- [34] Pucek Z, editor. European bison: status survey and conservation action plan. IUCN; 2004. <http://www.loc.gov/catdir/enhancements/fy0739/2005412156-d.html>.
- [35] King SRB, Boyd L, Zimmermann W, Kendall BE. *Equus ferus*. The IUCN red list of threatened species. 2015.
- [36] Iucn Ssc Antelope Specialist Group. *Oryx leucoryx*. IUCN Red List of Threatened Species 2017:e.T15569A50191626. <https://www.iucnredlist.org/species/15569/50191626>.
- [37] Murray DL, Bastille-Rousseau G, Adams JR, Waits LP. The challenges of red wolf conservation and the fate of an endangered species recovery program. *CONSERVATION LETTERS* 2015;8(5):338–44. <https://doi.org/10.1111/conl.12157>.
- [38] Murphy SM, Adams JR, Cox JJ, Waits LP. Substantial red wolf genetic ancestry persists in wild canids of southwestern Louisiana. *CONSERVATION LETTERS* 2019;12(2). <https://doi.org/10.1111/conl.12621>. e12621.
- [39] Lignereux L, Chaber A-L, Saegerman C, Heath L, Knowles NJ, Wadsworth J, et al. Foot-and-mouth disease outbreaks in captive scimitar-horned oryx (*Oryx dammah*). *Transbound. Emerg. Dis.* 2020;67(4):1716–24. <https://doi.org/10.1111/tbed.13502>.
- [40] Marcuk V, Purchase C, Boer D de, Bürkle M, Scholtyssek K. Qualitative description of the submission and agonistic behavior of the Spix's Macaw (*Cyanopsitta spixii*, Spix 1824), with special reference to the displacement displays. *J Ethol* 2020. <https://doi.org/10.1007/s10164-020-00650-6>.
- [41] Comizzoli P, Brown JL, Holt WV. Reproductive science as an essential component of conservation biology: new edition. *Adv Exp Med Biol* 2019;1200:1–10. https://doi.org/10.1007/978-3-030-23633-5_1.
- [42] Howard JG, Wildt DE. Approaches and efficacy of artificial insemination in felids and mustelids. *Theriogenology* 2009;71(1):130–48. <https://doi.org/10.1016/j.theriogenology.2008.09.046>.
- [43] Lasley BL, Loskutoff NM, Anderson GB. The limitation of conventional breeding programs and the need and promise of assisted reproduction in nondomestic species. *Theriogenology* 1994;41(1):119–32. [https://doi.org/10.1016/S0093-691X\(05\)80057-3](https://doi.org/10.1016/S0093-691X(05)80057-3).
- [44] Herrick JR. Assisted reproductive technologies for endangered species conservation: developing sophisticated protocols with limited access to animals with unique reproductive mechanisms. *Biol Reprod* 2019;100(5):1158–70. <https://doi.org/10.1093/biolre/iz025>.
- [45] Mastromonaco GF, Songsasen N. Reproductive technologies for the conservation of wildlife and endangered species. In: Presicce GA, editor. *Reproductive technologies in animals*. Amsterdam: Academic Press; 2020. p. 99–117.
- [46] Saragusty J, Ajmone-Marsan P, Sampino S, Modlinski JA. Reproductive biotechnology and critically endangered species: merging in vitro gametogenesis with inner cell mass transfer. *Theriogenology* 2020;155:176–84. <https://doi.org/10.1016/j.theriogenology.2020.06.009>.
- [47] Reproductive sciences in animal conservation: progress and prospects. In: Holt WV, Brown JL, Comizzoli P, editors. *Advances in experimental medicine and biology*. vol. 753. Springer; 2014.
- [48] Ryder O, Miller W, Ralls K, Ballou JD, Steiner CC, Mittelberg A, et al. Whole genome sequencing of California condors is now utilized for guiding genetic management. *Plant & Animal Genome Conference 2016;XXIV:W741*.
- [49] Hunter ME, Hoban SM, Bruford MW, Segelbacher G, Bernatchez L. Next-generation conservation genetics and biodiversity monitoring. *Evol. Appl.* 2018;11(7):1029–34. <https://doi.org/10.1111/eva.12661>.
- [50] Bishop JA, Dunn JL, Grantham C, McClure C. Parental behavior of California condors in wild and captive populations: 2020 undergraduate research showcase. https://scholarworks.boisestate.edu/under_showcase_2020/15/; 2020.
- [51] Schwarzenberger F. The many uses of non-invasive faecal steroid monitoring in zoo and wildlife species. *Int Zoo Yearbk* 2007;41(1):52–74. <https://doi.org/10.1111/j.1748-1090.2007.00017.x>.
- [52] Hildebrandt TB, Hermes R, Colleoni S, Diecke S, Holtze S, Renfree MB, et al. Embryos and embryonic stem cells from the white rhinoceros. *Nat Commun* 2018;9(1):2589. <https://doi.org/10.1038/s41467-018-04959-2>.
- [53] Morrow CJ, Penfold LM, Wolfe BA. Artificial insemination in deer and non-domestic bovids. *Theriogenology* 2009;71(1):149–65. <https://doi.org/10.1016/j.theriogenology.2008.09.001>.
- [54] Kouba AJ, Lloyd RE, Houck ML, Silla AJ, Calatayud N, Trudeau VL, et al. Emerging trends for biobanking amphibian genetic resources: the hope, reality and challenges for the next decade. *Biol Conserv* 2013;164:10–21. <https://doi.org/10.1016/j.biocon.2013.03.010>.
- [55] Ciulow J, Trudeau VL, Kouba AJ. Amphibian declines in the twenty-first century: why we need assisted reproductive technologies. New York; Imprint: Springer. In: Holt WV, Brown JL, Comizzoli P, editors. *Reproductive sciences in animal conservation: progress and prospects*. first ed. New York, NY: Springer; 2014. p. 275–316. *Advances in experimental medicine and biology*; vol. 753.
- [56] Lueders I, Luther I, Scheepers G, van der Horst G. Improved semen collection method for wild felids: urethral catheterization yields high sperm quality in African lions (*Panthera leo*). *Theriogenology* 2012;78(3):696–701. <https://doi.org/10.1016/j.theriogenology.2012.02.026>.
- [57] Landowski V, Gill J. Einige Beobachtungen über das Sperma des Indischen Elefanten (*Elephas maximus* L.). *Zool Gart* 1964;(29):205.
- [58] O'Brien JK, Roth TL. Post-coital sperm recovery and cryopreservation in the Sumatran rhinoceros (*Dicerorhinus sumatrensis*) and application to gamete rescue in the African black rhinoceros (*Diceros bicornis*). *Reproduction* 2000; 263–71. <https://doi.org/10.1530/reprod/118.2.263>.
- [59] Brindley GS. Electroejaculation: its technique, neurological implications and uses. *J Neurol Neurosurg Psychiatry* 1981;44(1):9–18. <https://doi.org/10.1136/jnnp.44.1.9>.
- [60] Zambelli D, Prati F, Cunto M, Iacono E, Merlo B. Quality and in vitro fertilizing ability of cryopreserved cat spermatozoa obtained by urethral catheterization after medetomidine administration. *Theriogenology* 2008;69(4):485–90. <https://doi.org/10.1016/j.theriogenology.2007.10.019>.
- [61] Fasel NJ, Helfenstein F, Buff S, Richner H. Electroejaculation and semen buffer evaluation in the microbat *Carollia perspicillata*. *Theriogenology* 2015;83(5):904–10. <https://doi.org/10.1016/j.theriogenology.2014.11.030>.
- [62] Roth TL, Stoops MA, Atkinson MW, Blumer ES, Campbell MK, Cameron KN, et al. Semen collection in rhinoceroses (*Rhinoceros unicornis*, *Diceros bicornis*, *Ceratotherium simum*) by electroejaculation with a uniquely designed probe. *J Zoo Wildl Med* 2005;36(4):617–27. <https://doi.org/10.1638/05-019.1>.
- [63] Mackie P, Chan B, Franke M, Mastromonaco GF. Urethral catheterization as an alternative method for collecting sperm in the black-footed ferret (*Mustela nigripes*). *Conserv. Physiol.* 2020;8(1). <https://doi.org/10.1093/conphys/coaa078>. coaa078.
- [64] Behr B, Rath D, Hildebrandt TB, Goeritz F, Blottner S, Portas TJ, et al. Germany/Australia index of sperm sex sortability in elephants and rhinoceros. *Reprod Domest Anim* 2009;44(2):273–7. <https://doi.org/10.1111/j.1439-0531.2007.01056.x>.
- [65] Hermes R, Behr B, Hildebrandt TB, Blottner S, Sieg B, Frenzel A, et al. Sperm sex-sorting in the Asian elephant (*Elephas maximus*). *Anim Reprod Sci*

- 2009;112(3–4):390–6. <https://doi.org/10.1016/j.anireprosci.2008.05.007>.
- [66] Saragusty J, Diecke S, Drukker M, Durrant B, Friedrich Ben-Nun I, Galli C, et al. Rewinding the process of mammalian extinction. *Zoo Biol* 2016;35(4):280–92. <https://doi.org/10.1002/zoo.21284>.
- [67] Roldan ERS, Gomendio M, Gardé JJ, Espeso G, Ledda S, Berlinguer F, et al. Inbreeding and reproduction in endangered ungulates: preservation of genetic variation through the Organization of Genetic Resource Banks. *Reprod Domest Anim* 2006;41(Suppl 2):82–92. <https://doi.org/10.1111/j.1439-0531.2006.00772.x>.
- [68] Robeck TR, Steinman KJ, Greenwell M, Ramirez K, van Bonn W, Yoshioka M, et al. Seasonality, estrous cycle characterization, estrus synchronization, semen cryopreservation, and artificial insemination in the Pacific white-sided dolphin (*Lagenorhynchus obliquidens*). *Reproduction* 2009;138(2):391–405. <https://doi.org/10.1530/REP-08-0528>.
- [69] Robeck TR, Steinman KJ, Montano GA, Katsumata E, Osborn S, Dalton L, et al. Deep intra-uterine artificial inseminations using cryopreserved spermatozoa in beluga (*Delphinapterus leucas*). *Theriogenology* 2010;74(6):989–1001. <https://doi.org/10.1016/j.theriogenology.2010.04.028>.
- [70] Robeck TR, Steinman KJ, Yoshioka M, Jensen E, O'Brien JK, Katsumata E, et al. Estrous cycle characterisation and artificial insemination using frozen-thawed spermatozoa in the bottlenose dolphin (*Tursiops truncatus*). *Reproduction* 2005;129(5):659–74. <https://doi.org/10.1530/rep.1.00516>.
- [71] Robeck TR, Montano GA, Steinman KJ, Smolensky P, Sweeney J, Osborn S, et al. Development and evaluation of deep intra-uterine artificial insemination using cryopreserved sexed spermatozoa in bottlenose dolphins (*Tursiops truncatus*). *Anim Reprod Sci* 2013;139(1–4):168–81. <https://doi.org/10.1016/j.anireprosci.2013.04.004>.
- [72] O'Brien JK, Steinman KJ, Schmitt T, Robeck TR. Semen collection, characterisation and artificial insemination in the beluga (*Delphinapterus leucas*) using liquid-stored spermatozoa. *Reprod Fertil Dev* 2008;20(7):770–83. <https://doi.org/10.1071/RD08031>.
- [73] Hildebrandt TB, Hermes R, Walzer C, Sós E, Molnar V, Mezösi L, et al. Artificial insemination in the anoestrous and the postpartum white rhinoceros using GnRH analogue to induce ovulation. *Theriogenology* 2007;67(9):1473–84. <https://doi.org/10.1016/j.theriogenology.2007.03.005>.
- [74] Hermes R, Goeritz F, Saragusty J, Sós E, Molnar V, Reid CE, et al. First successful artificial insemination with frozen-thawed semen in rhinoceros. *Theriogenology* 2009;71(3):393–9. <https://doi.org/10.1016/j.theriogenology.2008.10.008>.
- [75] Stoops MA, Campbell MK, DeChant CJ, Hauser J, Kottwitz J, Pairan RD, et al. Enhancing captive Indian rhinoceros genetics via artificial insemination of cryopreserved sperm. *Anim Reprod Sci* 2016;172:60–75. <https://doi.org/10.1016/j.anireprosci.2016.07.003>.
- [76] Swanson WF. Practical application of laparoscopic oviductal artificial insemination for the propagation of domestic cats and wild felids. *Reprod Fertil Dev* 2018;31(1):27–39. <https://doi.org/10.1071/RD18350>.
- [77] Jewgenow K, Songsasen N. Reproduction and advances in reproductive studies in carnivores. *Adv Exp Med Biol* 2014;753:205–39. https://doi.org/10.1007/978-1-4939-0820-2_10.
- [78] Blanco JM, Wildt DE, Höfle U, Voelker W, Donoghue AM. Implementing artificial insemination as an effective tool for ex situ conservation of endangered avian species. *Theriogenology* 2009;71(1):200–13. <https://doi.org/10.1016/j.theriogenology.2008.09.019>.
- [79] Saint Jalme M, Gaucher P, Paillat P. Artificial insemination in Houbara bustards (*Chlamydotis undulata*): influence of the number of spermatozoa and insemination frequency on fertility and ability to hatch. *J Reprod Fertil* 1994;100(1):93–103. <https://doi.org/10.1530/jrf.0.1000093>.
- [80] Rabier R, Robert A, Lacroix F, Lesobre L. Genetic assessment of a conservation breeding program of the houbara bustard (*Chlamydotis undulata undulata*) in Morocco, based on pedigree and molecular analyses. *Zoo Biol* 2020. <https://doi.org/10.1002/zoo.21569>.
- [81] Ministério do Meio Ambiente. Action Plan For The Conservation Of The Spix's Macaw - Cyanopsitta Spixii: Captive Program Protocol - Version 2018.
- [82] Gee GF. Artificial insemination and cryopreservation of semen from nondomestic birds.: Proceedings. In: Bakst MR, editor. International symposium on the artificial insemination of poultry: proceedings. Savoy, Ill.; 1995. p. 262–79.
- [83] Howard JG, Lynch C, Santymire RM, Marinari PE, Wildt DE. Recovery of gene diversity using long-term cryopreserved spermatozoa and artificial insemination in the endangered black-footed ferret. *Anim Conserv* 2016;19(2):102–11. <https://doi.org/10.1111/acv.12229>.
- [84] Santymire R, Graves G. Black-footed Ferret SAFE program action plan 2019–2021. https://assets.speakcdn.com/assets/2332/2019-2021_black-footed_ferret_safe_program_plan-updated_032020.pdf; 2019.
- [85] Santymire RM, Lavin SR, Branvold-Faber H, Kreeger J, Che-Castaldo J, Rafacz M, et al. Influence of vitamin E and carcass feeding supplementation on fecal glucocorticoid and androgen metabolites in male black-footed ferrets (*Mustela nigripes*). *PLoS One* 2020;15(10). <https://doi.org/10.1371/journal.pone.0241085>. e0241085.
- [86] Zhu L, Hu Y, Qi D, Wu H, Zhan X, Zhang Z, et al. Genetic consequences of historical anthropogenic and ecological events on giant pandas. *Ecology* 2013;94(10):2346–57. <https://doi.org/10.1890/12-1451.1>.
- [87] Martin-Wintle MS, Shepherdson D, Zhang G, Huang Y, Luo B, Swaisgood RR. Do opposites attract? Effects of personality matching in breeding pairs of captive giant pandas on reproductive success. *Biol Conserv* 2017;207:27–37. <https://doi.org/10.1016/j.biocon.2017.01.010>.
- [88] Wei R, Zhang G, Yin F, Zhang H, Liu D. Enhancing captive breeding in giant pandas (*Ailuropoda melanoleuca*): maintaining lactation when cubs are rejected, and understanding variation in milk collection and associated factors. *Zoo Biol* 2009;28(4):331–42. <https://doi.org/10.1002/zoo.20232>.
- [89] Huang Y, Li D, Zhou Y, Zhou Q, Li R, Wang C, et al. Factors affecting the outcome of artificial insemination using cryopreserved spermatozoa in the giant panda (*Ailuropoda melanoleuca*). *Zoo Biol* 2012;31(5):561–73. <https://doi.org/10.1002/zoo.20421>.
- [90] Brown JL, Goeritz F, Pratt-Hawkes N, Hermes R, Galloway M, Graham LH, et al. Successful artificial insemination of an asian elephant at the national zoological park. *Zoo Biol* 2004;23(1):45–63. <https://doi.org/10.1002/zoo.10116>.
- [91] Hildebrandt TB, Goeritz F, Hermes R, Reid CE, Dehnhard M, Brown JL. Aspects of the reproductive biology and breeding management of Asian and African elephants *Elephas maximus* and *Loxodonta africana*. *Int Zoo Yearbk* 2006;40(1):20–40. <https://doi.org/10.1111/j.1748-1090.2006.00020.x>.
- [92] Hildebrandt TB, Hermes R, Jewgenow K, Goeritz F. Ultrasonography as an important tool for the development and application of reproductive technologies in non-domestic species. *Theriogenology* 2000;53(1):73–84. [https://doi.org/10.1016/s0093-691x\(99\)00241-1](https://doi.org/10.1016/s0093-691x(99)00241-1).
- [93] Hermes R, Goeritz F, Streich WJ, Hildebrandt TB. Assisted reproduction in female rhinoceros and elephants—current status and future perspective. *Reprod Domest Anim* 2007;42(Suppl 2):33–44. <https://doi.org/10.1111/j.1439-0531.2007.00924.x>.
- [94] Hildebrandt TB, Hermes R, Saragusty J, Potier R, Schwammer HM, Balfanz F, et al. Enriching the captive elephant population genetic pool through artificial insemination with frozen-thawed semen collected in the wild. *Theriogenology* 2012;78(6):1398–404. <https://doi.org/10.1016/j.theriogenology.2012.06.014>.
- [95] Hermes R, Goeritz F, Portas TJ, Bryant BR, Kelly JM, MacLellan LJ, et al. Ovarian superstimulation, transrectal ultrasound-guided oocyte recovery, and IVF in rhinoceros. *Theriogenology* 2009;72(7):959–68. <https://doi.org/10.1016/j.theriogenology.2009.06.014>.
- [96] Galli C, Hermes R, Goeritz F, Colleoni S, Diecke S, Drukker M, et al. First results of oocyte maturation and in-vitro-fertilisation (IVF) in Sumatran and northern white rhinoceroses. In: Proceedings of the scientific program of the 15th international elephant & rhino conservation and research symposium. 51; 2016. Singapore.
- [97] Heape Walter. III. Preliminary note on the transplantation and growth of mammalian ova within a uterine foster-mother. *Proc Royal Soc* 1891;48:292–5.
- [98] Kraemer DC, Moore GT, Kramen MA. Baboon infant produced by embryo transfer. *Science* 1976;192(4245):1246–7. <https://doi.org/10.1126/science.818710>.
- [99] Dresser BL, Kramer L, Pope CE, Dahlhausen RD, Blausler C. Superovulation of African eland (*Taurotragus oryx*) and interspecies embryo transfer to Holstein cattle. *Theriogenology* 1982;17(1):86.
- [100] Stover J, Evans J. Interspecies embryo transfer from gaur (*Bos gaurus*) to domestic Holstein cattle (*Bos taurus*) at the New York Zoological Park. In: 10th international congress on animal reproduction and artificial insemination; 1984. p. 10–4.
- [101] Dresser BL, Pope CE, Kramer L, Kuehn G, Dahlhausen RD, Maruska EJ, et al. Birth of bongo antelope (*Tragelaphus euryceros*) to eland antelope (*Tragelaphus oryx*) and cryopreservation of bongo embryos. *Theriogenology* 1985;23(1):190. [https://doi.org/10.1016/0093-691X\(85\)90096-2](https://doi.org/10.1016/0093-691X(85)90096-2).
- [102] Swanson WF. Laparoscopic oviductal embryo transfer and artificial insemination in felids—challenges, strategies and successes. *Reprod Domest Anim* 2012;47(Suppl 6):136–40. <https://doi.org/10.1111/rda.12069>.
- [103] Donoghue AM, Johnston LA, Seal US, Armstrong DL, Tilton RL, Wolf P, et al. In vitro fertilization and embryo development in vitro and in vivo in the tiger (*Panthera tigris*). *Biol Reprod* 1990;43(5):733–44. <https://doi.org/10.1095/biolreprod43.5.733>.
- [104] Pope CE, Gómez MC, Dresser BL. In vitro embryo production and embryo transfer in domestic and non-domestic cats. *Theriogenology* 2006;66(6–7):1518–24. <https://doi.org/10.1016/j.theriogenology.2006.01.026>.
- [105] Swanson WF. Reproductive biotechnology and conservation of the forgotten felids—the small cats. *Reproductive biotechnology and conservation of the forgotten felids—the small cats*. In: Proceedings of the 1st international symposium on assisted reproductive technologies for conservation and genetic management of wildlife. vols. 17–8; 2001.
- [106] Herrick JR, Mehrdadfar F, Campbell MK, Levens G, Leiske K, Swanson WF. Birth of sand cat (*Felis margarita*) kittens following in vitro fertilization and embryo transfer. *Biol Reprod* 2010;83(Suppl_1):28. <https://doi.org/10.1093/biolreprod/83.s1.28>.
- [107] Pope CE, Gómez MC, Dumas C, MacLean RA, Crichton E, Armstrong D, et al. 122 birth of Black-Footed Cat kittens after transfer of cryopreserved embryos produced by in vitro-fertilization of oocytes with cryopreserved sperm. *Reprod Fertil Dev* 2012;24(1):173. <https://doi.org/10.1071/RDv24n1Ab122>.
- [108] Johnston LA, Parrish JJ, Monson R, Leibfried-Rutledge L, Susko-Parrish JL, Northey DL, et al. Oocyte maturation, fertilization and embryo development in vitro and in vivo in the gaur (*Bos gaurus*). *J Reprod Fertil* 1994;100(1):131–6. <https://doi.org/10.1530/jrf.0.1000131>.
- [109] Locatelli Y, Hendriks A, Vallet J-C, Baril G, Duffard N, Bon N, et al. Assessment LOPU-IVF in Japanese sika deer (*Cervus nippon nippon*) and application to

- Vietnamese sika deer (*Cervus nippon pseudaxis*) a related subspecies threatened with extinction. *Theriogenology* 2012;78(9):2039–49. <https://doi.org/10.1016/j.theriogenology.2012.07.025>.
- [110] Smithsonian Conservation Biology Institute. First Eld's deer born from in vitro fertilization with help of Smithsonian Conservation Biology Institute scientists. <https://www.si.edu/newsdesk/releases/first-eld-s-deer-born-vitro-fertilization-help-smithsonian-conservation-biologyinstitutes>. [Accessed 31 August 2019].
- [111] Berg DK, Beaumont SE, Berg MC, Asher GW. Red deer (*Cervus elaphus*) calves born from in vitro-produced blastocysts fertilized and cultured in deer synthetic oviduct fluid. *Reprod Fertil Dev* 2004;16(2):222. <https://doi.org/10.1071/RDv16n1Ab201>.
- [112] Bavister Barry D, Boatman Dorothy E. IVF in nonhuman primates: current status and future directions. In: *In vitro fertilization and embryo transfer in primates*. New York, NY: Springer; 1993. p. 30–45. https://link.springer.com/chapter/10.1007/978-1-4612-2716-8_2.
- [113] Steven L. Monfort. "Mayday mayday mayday", the millennium ark is sinking!. In: *Reproductive sciences in animal conservation*. New York, NY: Springer; 2014. p. 15–31. https://link.springer.com/chapter/10.1007/978-1-4939-0820-2_2.
- [114] Kouba AJ, Vance CK, Willis EL. Artificial fertilization for amphibian conservation: current knowledge and future considerations. *Theriogenology* 2009;71(1):214–27. <https://doi.org/10.1016/j.theriogenology.2008.09.055>.
- [115] Kouba AJ, Willis EL, Vance CK, Hasenstab S, Reichling S, Krebs J, et al. Development of assisted reproduction technologies for the endangered Mississippi gopher frog (*Rana sevosia*) and sperm transfer for in vitro fertilization. *Reprod Fertil Dev* 2012;24(1):170. <https://doi.org/10.1071/RDv24n1Ab116>.
- [116] Baldassarre H, Karatzas CN. Advanced assisted reproduction technologies (ART) in goats. *Anim Reprod Sci* 2004;82–83:255–66. <https://doi.org/10.1016/j.anireprosci.2004.04.027>.
- [117] Callaway E. Stem-cell plan aims to bring rhino back from brink of extinction. *Nature* 2016;533(7601):20–1. <https://doi.org/10.1038/533020a>.
- [118] Hikabe O, Hamazaki N, Nagamatsu G, Obata Y, Hirao Y, Hamada N, et al. Reconstitution in vitro of the entire cycle of the mouse female germ line. *Nature* 2016;539(7628):299–303. <https://doi.org/10.1038/nature20104>.
- [119] Hildebrandt TB, Holtze S, Biasetti P, Colleoni S, Mori B de, Diecke S, et al. Conservation research in times of covid-19 – the rescue of the northern white rhino. *Journal of Applied Animal Ethics Research* 2021. Accepted for publication 17.12.2020.
- [120] Yamanaka S. Induction of pluripotent stem cells from mouse fibroblasts by four transcription factors. *Cell Prolif* 2008;41(Suppl 1):51–6. <https://doi.org/10.1111/j.1365-2184.2008.00493.x>.
- [121] Fishel S, Aslam I, Lisi F, Rinaldi L, Timson J, Jacobson M, et al. Should ICSI be the treatment of choice for all cases of in-vitro conception? *Hum Reprod* 2000;15(6):1278–83. <https://doi.org/10.1093/humrep/15.6.1278>.
- [122] Magata F, Tsuchiya K, Okubo H, Ideta A. Application of intracytoplasmic sperm injection to the embryo production in aged cows. *J Vet Med Sci* 2019;81(1):84–90. <https://doi.org/10.1292/jvms.18-0284>.
- [123] Onishi A, Perry ACF. Livestock production via micromanipulation. In: *In vitro fertilization*. Cham: Springer; 2019. p. 939–43. https://link.springer.com/chapter/10.1007/978-3-319-43011-9_79.
- [124] Ibtisham F. Animal cloning applications and issues. *Russ J Genet*.
- [125] Walton M. Commercial applications of SCNT in livestock. In: *Animal biotechnology 2*. Cham: Springer; 2018. p. 21–35. https://link.springer.com/chapter/10.1007/978-3-319-92348-2_2.
- [126] Berg DK, Li C, Asher G, Wells DN, Oback B. Red deer cloned from antler stem cells and their differentiated progeny. *Biol Reprod* 2007;77(3):384–94. <https://doi.org/10.1095/biolreprod.106.058172>.
- [127] Liu Z, Cai Y, Wang Y, Nie Y, Zhang C, Xu Y, et al. Cloning of macaque monkeys by somatic cell nuclear transfer. *Cell* 2018;174(1):245. <https://doi.org/10.1016/j.cell.2018.01.036>.
- [128] Lanza RP, Cibelli JB, Diaz F, Moraes CT, Farin PW, Farin CE, et al. Cloning of an endangered species (*Bos gaurus*) using interspecies nuclear transfer. *Cloning* 2000;2(2):79–90. <https://doi.org/10.1089/152045500436104>.
- [129] Loi P, Ptak G, Barboni B, Fulka J, Cappai P, Clinton M. Genetic rescue of an endangered mammal by cross-species nuclear transfer using post-mortem somatic cells. *Nat Biotechnol* 2001;19(10):962–4. <https://doi.org/10.1038/nbt1001-962>.
- [130] Gómez MC, Pope CE, Giraldo A, Lyons LA, Harris RF, King AL, et al. Birth of African Wildcat cloned kittens born from domestic cats. *Clon Stem Cell* 2004;6(3):247–58. <https://doi.org/10.1089/clo.2004.6.247>.
- [131] Kim MK, Jang G, Oh HJ, Yuda F, Kim HJ, Hwang WS, et al. Endangered wolves cloned from adult somatic cells. *Clon Stem Cell* 2007;9(1):130–7. <https://doi.org/10.1089/clo.2006.0034>.
- [132] Oh HJ, Kim MK, Jang G, Kim HJ, Hong SG, Park JE, et al. Cloning endangered gray wolves (*Canis lupus*) from somatic cells collected postmortem. *Theriogenology* 2008;70(4):638–47. <https://doi.org/10.1016/j.theriogenology.2008.04.032>.
- [133] Gómez MC, Pope CE, Kutner RH, Ricks DM, Lyons LA, Ruhe M, et al. Nuclear transfer of sand cat cells into enucleated domestic cat oocytes is affected by cryopreservation of donor cells. *Clon Stem Cell* 2008;10(4):469–83. <https://doi.org/10.1089/clo.2008.0021>.
- [134] Folch J, Cocero MJ, Chesné P, Alabart JL, Domínguez V, Coggié Y, et al. First birth of an animal from an extinct subspecies (*Capra pyrenaica pyrenaica*) by cloning. *Theriogenology* 2009;71(6):1026–34. <https://doi.org/10.1016/j.theriogenology.2008.11.005>.
- [135] Fatira E, Havelka M, Labbé C, Depincé A, Pšenička M, Saito T. A newly developed cloning technique in sturgeons; an important step towards recovering endangered species. *Sci Rep* 2019;9(1):10453. <https://doi.org/10.1038/s41598-019-46892-4>.
- [136] Ryder OA, Benirschke K. The potential use of "cloning" in the conservation effort. *Zoo Biol* 1997;16(4):295–300. [https://doi.org/10.1002/\(SICI\)1098-2361\(1997\)16:4<295::AID-ZOO1>3.0.CO;2-5](https://doi.org/10.1002/(SICI)1098-2361(1997)16:4<295::AID-ZOO1>3.0.CO;2-5).
- [137] Holt WW, Pickard AR, Prather RS. Wildlife conservation and reproductive cloning. *Reproduction* 2004;127(3):317–24. <https://doi.org/10.1530/rep.1.00074>.
- [138] Polzin VJ, Anderson DL, Anderson GB, BonDurant RH, Butler JE, Pashen RL, et al. Production of sheep-goat chimeras by inner cell mass transplantation. *J Anim Sci* 1987;65(1):325–30. <https://doi.org/10.2527/jas1987.651325x>.
- [139] Meinecke-Tillmann S, Meinecke B. Experimental chimaeras-removal of reproductive barrier between sheep and goat. *Nature* 1984;307(5952):637–8. <https://doi.org/10.1038/307637a0>.
- [140] Loi P, Galli C, Lazzari G, Matsukawa K, Fulka J, Goeritz F, et al. Development to term of sheep embryos reconstructed after inner cell mass/trophoblast exchange. *J Reprod Dev* 2018;64(2):187–91. <https://doi.org/10.1262/jrd.2017-109>.
- [141] Sato T, Katagiri K, Kubota Y, Ogawa T. In vitro sperm production from mouse spermatogonial stem cell lines using an organ culture method. *Nat Protoc* 2013;8(11):2098–104. <https://doi.org/10.1038/nprot.2013.138>.
- [142] Taketo T. The role of sex chromosomes in mammalian germ cell differentiation: can the germ cells carrying X and Y chromosomes differentiate into fertile oocytes? *Asian J Androl* 2015;17(3):360–6. <https://doi.org/10.4103/1008-682x.143306>.
- [143] Hayashi K, Ogushi S, Kurimoto K, Shimamoto S, Ohta H, Saitou M. Offspring from oocytes derived from in vitro primordial germ cell-like cells in mice. *Science* 2012;338(6109):971–5. <https://doi.org/10.1126/science.1226889>.
- [144] Sato T, Katagiri K, Yokonishi T, Kubota Y, Inoue K, Ogonuki N, et al. In vitro production of fertile sperm from murine spermatogonial stem cell lines. *Nat Commun* 2011;2:472. <https://doi.org/10.1038/ncomms1478>.
- [145] Hayashi K, Saitou M. Generation of eggs from mouse embryonic stem cells and induced pluripotent stem cells. *Nat Protoc* 2013;8(8):1513–24. <https://doi.org/10.1038/nprot.2013.090>.
- [146] Stanton MM, Tzatzalos E, Donne M, Kolundzic N, Helgason I, Ilic D. Prospects for the use of induced pluripotent stem cells in animal conservation and environmental protection. *Stem Cells Transl. Med.* 2019;8(1):7–13. <https://doi.org/10.1002/sctm.18-0047>.
- [147] Lu Yangqing, West D Franklin, Jordan J Brian, Mumaw L Jennifer, Jordan T Erin, Gallegos-Cardenas Amalia, et al. Avian-induced pluripotent stem cells derived using human reprogramming factors. *Stem Cells Dev* 2012;21:394–403.
- [148] Verma Rajneesh, Verma J Paul. Using stem cells to study and preserve biodiversity in endangered big cats. *Stem Cells in Animal Species: From Pre-clinic to Biodiversity*. Cham: Humana Press; 2014. p. 109–17.
- [149] Verma R, Holland MK, Temple-Smith P, Verma PJ. Inducing pluripotency in somatic cells from the snow leopard (*Panthera uncia*), an endangered felid. *Theriogenology* 2012;77(1):220–8. <https://doi.org/10.1016/j.theriogenology.2011.09.022>. 228.e1-2.
- [150] Ben-Nun IF, Montague SC, Houck ML, Ryder O, Loring JF. Generation of induced pluripotent stem cells from mammalian endangered species. *Methods Mol Biol* 2015;1330:101–9. https://doi.org/10.1007/978-1-4939-2848-4_10.
- [151] Korody ML, Pivaroff C, Nguyen TD, Peterson SE, Ryder OA, Loring JF. Four new induced pluripotent stem cell lines produced from northern white rhinoceros with non-integrating reprogramming factors. <https://doi.org/10.1101/202499>; 2017.
- [152] Ryder OA, Onuma M. Viable cell culture banking for biodiversity characterization and conservation. *Annu. Rev. Anim. Biosci.* 2018;6:83–98. <https://doi.org/10.1146/annurev-animal-030117-014556>.
- [153] Comizzoli P, Songsasen N, Hagedorn M, Wildt DE. Comparative cryobiological traits and requirements for gametes and gonadal tissues collected from wildlife species. *Theriogenology* 2012;78(8):1666–81. <https://doi.org/10.1016/j.theriogenology.2012.04.008>.
- [154] Nel-Themaat L, Gómez MC, Damiani P, Wirtu G, Dresser BL, Bondioli KR, et al. Isolation, culture and characterisation of somatic cells derived from semen and milk of endangered sheep and eland antelope. *Reprod Fertil Dev* 2007;19(4):576–84. <https://doi.org/10.1071/RD06153>.
- [155] Iyengar V, Albaugh GP, Lohani A, Nair PP. Human stools as a source of viable colonic epithelial cells. *Faseb J* 1991;5(13):2856–9. <https://doi.org/10.1096/fasebj.5.13.1655550>.
- [156] Ganjibakhsh M, Aminishakib P, Farzaneh P, Karimi A, Fazeli SAS, Rajabi M, et al. Establishment and characterization of primary cultures from Iranian oral squamous cell carcinoma patients by enzymatic method and explant culture. *J Dent (Tehran)* 2017;14(4):191–202.
- [157] Yeager TR. Constructing immortalized human cell lines. *Curr Opin Biotechnol* 1999;10(5):465–9. [https://doi.org/10.1016/S0958-1669\(99\)00011-7](https://doi.org/10.1016/S0958-1669(99)00011-7).
- [158] Hayflick L, Moorhead PS. The serial cultivation of human diploid cell strains. *Exp Cell Res* 1961;25:585–621. [https://doi.org/10.1016/0014-4827\(61\)90192-6](https://doi.org/10.1016/0014-4827(61)90192-6).
- [159] Gómez MC, Pope CE, López M, Dumas C, Giraldo A, Dresser BL. Chromosomal

- aneuploidy in African Wildcat somatic cells and cloned embryos. *Clon Stem Cell* 2006;8(2):69–78. <https://doi.org/10.1089/clo.2006.8.69>.
- [160] Rebuzzini P, Zuccotti M, Redi CA, Garagna S. Achilles' heel of pluripotent stem cells: genetic, genomic and epigenetic variations during prolonged culture. *Cell Mol Life Sci* 2016;73(13):2453–66. <https://doi.org/10.1007/s00018-016-2171-8>.
- [161] Saragusty J, Anzalone DA, Palazzese L, Arav A, Patrizio P, Gosálvez J, et al. Dry biobanking as a conservation tool in the Anthropocene. *Theriogenology* 2020;150:130–8. <https://doi.org/10.1016/j.theriogenology.2020.01.022>.
- [162] Polge C, Smith AU, Parkes AS. Revival of spermatozoa after vitrification and dehydration at low temperatures. *Nature* 1949;164(4172):666. <https://doi.org/10.1038/164666a0>.
- [163] Stewart D. Storage of bull spermatozoa at low temperature. *Vet Rec* 1951;63:65–6. <https://ci.nii.ac.jp/naid/10019971791/>.
- [164] Smith R. World's oldest frozen semen: trialled in merino sire evaluation and results are now available. *Beyond the bale* 2020;(83):48.
- [165] Wildt DE, Roth TL. Assisted reproduction for managing and conserving threatened felids. *Int Zoo Yearbk* 1997;35(1):164–72. <https://doi.org/10.1111/j.1748-1090.1997.tb01207.x>.
- [166] Comizzoli P. Birth of a Giant Panda cub after artificial insemination with frozen-thawed semen: a powerful reminder about the key role of bio-preservation and biobanking for wildlife conservation. *Biopreserv Biobanking* 2020;18(5):349–50. <https://doi.org/10.1089/bio.2020.29076.pjc>.
- [167] Comizzoli P, Holt WV. Recent advances and prospects in germplasm preservation of rare and endangered species. *Adv Exp Med Biol* 2014;753:331–56. https://doi.org/10.1007/978-1-4939-0820-2_14.
- [168] Brown ME, Singh RP, Pukazhenthi B, Keefer CL, Songsasen N. Cryopreservation effects on sperm function and fertility in two threatened crane species. *Cryobiology* 2018;82:148–54. <https://doi.org/10.1016/j.cryobiol.2018.01.010>.
- [169] Browne RK, Silla AJ, Upton R, Della Togna G, Marcec-Greaves R, Shishova NV, et al. Sperm collection and storage for the sustainable management of amphibian biodiversity. *Theriogenology* 2019;133:187–200. <https://doi.org/10.1016/j.theriogenology.2019.03.035>.
- [170] Nusbaumer D, Da Marques Cunha L, Wedekind C. Sperm cryopreservation reduces offspring growth. *Proc Biol Sci* 2019;286(1911):20191644. <https://doi.org/10.1098/rspb.2019.1644>.
- [171] Strand J, Thomsen H, Jensen JB, Marcussen C, Nicolajsen TB, Skriver MB, et al. Biobanking in amphibian and reptilian conservation and management: opportunities and challenges. *Conservation Genet Resour* 2020;12(4):709–25. <https://doi.org/10.1007/s12686-020-01142-y>.
- [172] Comizzoli P, Mermillod P, Maugé R. Reproductive biotechnologies for endangered mammalian species. *Reprod Nutr Dev* 2000;40(5):493–504. <https://doi.org/10.1051/rnd:2000113>.
- [173] Prieto MT, Sanchez-Calabuig MJ, Hildebrandt TB, Santiago-Moreno J, Saragusty J. Sperm cryopreservation in wild animals. *Eur J Wildl Res* 2014;60(6):851–64. <https://doi.org/10.1007/s10344-014-0858-4>.
- [174] Saragusty J, Arav A. Current progress in oocyte and embryo cryopreservation by slow freezing and vitrification. *Reproduction* 2011;141(1):1–19. <https://doi.org/10.1530/REP-10-0236>.
- [175] Hermes R, Hildebrandt TB, Goeritz F, Fasel NJ, Holtze S. First cryopreservation of phyllostomid bat sperm. *Theriogenology* 2019;131:28–31. <https://doi.org/10.1016/j.theriogenology.2019.03.014>.
- [176] Saragusty J, Osmers J-H, Hildebrandt TB. Controlled ice nucleation—is it really needed for large-volume sperm cryopreservation? *Theriogenology* 2016;85(7):1328–33. <https://doi.org/10.1016/j.theriogenology.2015.12.019>.
- [177] Arav A, Saragusty J. Directional freezing of sperm and associated derived technologies. *Anim Reprod Sci* 2016;169:6–13. <https://doi.org/10.1016/j.anireprosci.2016.02.007>.
- [178] Hermes R, Saragusty J, Goeritz F, Bartels P, Potier R, Baker B, et al. Freezing African elephant semen as a new population management tool. *PLoS One* 2013;8(3). <https://doi.org/10.1371/journal.pone.0057616>. e57616.
- [179] Si W, Hildebrandt TB, Reid CE, Krieger R, Ji W, Fassbender M, et al. The successful double cryopreservation of rabbit (*Oryctolagus cuniculus*) semen in large volume using the directional freezing technique with reduced concentration of cryoprotectant. *Theriogenology* 2006;65(4):788–98. <https://doi.org/10.1016/j.theriogenology.2005.06.010>.
- [180] Poo S, Hinkson KM. Applying cryopreservation to anuran conservation biology. *Conserv Sci and Prac* 2019;1(9). <https://doi.org/10.1111/csp2.91>.
- [181] Cabrita E, Sarasquete C, Martínez-Páramo S, Robles V, Beirão J, Pérez-Cerezales S, et al. Cryopreservation of fish sperm: applications and perspectives. *J Appl Ichthyol* 2010;26(5):623–35. <https://doi.org/10.1111/j.1439-0426.2010.01556.x>.
- [182] Hagedorn M, Spindler RE. The reality, use and potential for cryopreservation of coral reefs. *Adv Exp Med Biol* 2014;753:317–29. https://doi.org/10.1007/978-1-4939-0820-2_13.
- [183] Pukazhenthi B, Comizzoli P, Travis AJ, Wildt DE. Applications of emerging technologies to the study and conservation of threatened and endangered species. *Reprod Fertil Dev* 2006;18(1–2):77–90. <https://doi.org/10.1071/RD05117>.
- [184] Rall WF, Fahy GM. Ice-free cryopreservation of mouse embryos at -196 degrees C by vitrification. *Nature* 1985;313(6003):573–5. <https://doi.org/10.1038/313573a0>.
- [185] Vajta G, Kuwayama M. Improving cryopreservation systems. *Theriogenology* 2006;65(1):236–44. <https://doi.org/10.1016/j.theriogenology.2005.09.026>.
- [186] Boutelle S, Lenahan K, Krisher R, Bauman KL, Asa CS, Silber S. Vitrification of oocytes from endangered Mexican gray wolves (*Canis lupus baileyi*). *Theriogenology* 2011;75(4):647–54. <https://doi.org/10.1016/j.theriogenology.2010.10.004>.
- [187] Nowak A, Kochan J, Prochowska S, Partyka A, Młodawska W, Witarski W, et al. The viability of serval (*Leptailurus serval*) and Pallas cat (*Felis manul*) oocytes after cryopreservation using the rapid-I method. *Cryo Lett* 2019;40(4):226–30. <https://pubmed.ncbi.nlm.nih.gov/31278403/>.
- [188] Zahmel J, Jänsch S, Jewgenow K, Sandgreen D-M, Skalborg Simonsen K, Colombo M. Maturation and fertilization of African lion (*Panthera leo*) oocytes after vitrification. *Cryobiology* 2020. <https://doi.org/10.1016/j.cryobiol.2020.11.011>.
- [189] Rao BS, Mahesh YU, Suman K, Charan KV, Lakshmikantam U, Gibence HRW, et al. Meiotic maturation of vitrified immature chousingha (*Tetracerus quadricornis*) oocytes recovered postmortem. *Cryobiology* 2011;62(1):47–52. <https://doi.org/10.1016/j.cryobiol.2010.12.002>.
- [190] Czarny NA, Rodger JC. Vitrification as a method for genome resource banking oocytes from the endangered Tasmanian devil (*Sarcophilus harrisii*). *Cryobiology* 2010;60(3):322–5. <https://doi.org/10.1016/j.cryobiol.2010.02.007>.
- [191] Fahy GM, MacFarlane DR, Angell CA, Meryman HT. Vitrification as an approach to cryopreservation. *Cryobiology* 1984;21(4):407–26. [https://doi.org/10.1016/0011-2240\(84\)90079-8](https://doi.org/10.1016/0011-2240(84)90079-8).
- [192] Whittingham DG. Survival of mouse embryos after freezing and thawing. *Nature* 1971;233(5315):125–6. <https://doi.org/10.1038/233125a0>.
- [193] Kolibianakis EM, Venetis CA, Tarlatzis BC. Cryopreservation of human embryos by vitrification or slow freezing: which one is better? *Curr Opin Obstet Gynecol* 2009;21(3):270–4. <https://doi.org/10.1097/GCO.0b013e3283297dd6>.
- [194] Abdelhafez FF, Desai N, Abou-Setta AM, Falcone T, Goldfarb J. Slow freezing, vitrification and ultra-rapid freezing of human embryos: a systematic review and meta-analysis. *Reprod Biomed Online* 2010;20(2):209–22. <https://doi.org/10.1016/j.rbmo.2009.11.013>.
- [195] Hermes R, Hildebrandt TB, Goeritz F. Cryopreservation in rhinoceros—setting a new benchmark for sperm cryosurvival. *PLoS One* 2018;13(7). <https://doi.org/10.1371/journal.pone.0200154>. e0200154.
- [196] Gougeon A. Human ovarian follicular development: from activation of resting follicles to preovulatory maturation. *Ann Endocrinol (Paris)* 2010;71(3):132–43. <https://doi.org/10.1016/j.ando.2010.02.021>.
- [197] Fassbender M, Hildebrandt TB, Paris MCJ, Colenbrander B, Jewgenow K. High-resolution ultrasonography of xenografted domestic cat ovarian cortex. *J Reprod Dev* 2007;53(5):1023–34. <https://doi.org/10.1262/jrd.19021>.
- [198] Santos RR, Amorim C, Cecconi S, Fassbender M, Imhof M, Lornage J, et al. Cryopreservation of ovarian tissue: an emerging technology for female germline preservation of endangered species and breeds. *Anim Reprod Sci* 2010;122(3–4):151–63. <https://doi.org/10.1016/j.anireprosci.2010.08.010>.
- [199] Wiedemann C, Zahmel J, Jewgenow K. Short-term culture of ovarian cortex pieces to assess the cryopreservation outcome in wild felids for genome conservation. *BMC Vet Res* 2013;9(1):37. <https://doi.org/10.1186/1746-6148-9-37>.
- [200] Comizzoli P, Songsasen N, Wildt DE. Protecting and extending fertility for females of wild and endangered mammals. *Canc Treat Res* 2010;156:87–100. https://doi.org/10.1007/978-1-4419-6518-9_7.
- [201] Shinohara T, Inoue K, Ogonuki N, Kanatsu-Shinohara M, Mikai H, Nakata K, et al. Birth of offspring following transplantation of cryopreserved immature testicular pieces and in-vitro microinsemination. *Hum Reprod* 2002;17(12):3039–45. <https://doi.org/10.1093/humrep/17.12.3039>.
- [202] Kaneko H, Kikuchi K, Tanihara F, Noguchi J, Nakai M, Ito J, et al. Normal reproductive development of pigs produced using sperm retrieved from immature testicular tissue cryopreserved and grafted into nude mice. *Theriogenology* 2014;82(2):325–31. <https://doi.org/10.1016/j.theriogenology.2014.04.012>.
- [203] Kaneko H, Kikuchi K, Men NT, Noguchi J. Embryo production by intracytoplasmic injection of sperm retrieved from Meishan neonatal testicular tissue cryopreserved and grafted into nude mice. *Anim Sci J* 2019;90(2):158–66. <https://doi.org/10.1111/asj.13138>.
- [204] Fayomi AP, Peters K, Sukhwani M, Valli-Pulaski H, Shetty G, Meistrich ML, et al. Autologous grafting of cryopreserved prepubertal rhesus testis produces sperm and offspring. *Science* 2019;363(6433):1314–9. <https://doi.org/10.1126/science.aav2914>.
- [205] Liu J, Cheng KM, Silversides FG. Production of live offspring from testicular tissue cryopreserved by vitrification procedures in Japanese quail (*Coturnix japonica*). *Biol Reprod* 2013;88(5):124. <https://doi.org/10.1095/biolreprod.113.108951>.
- [206] Higaki S, Kuwata N, Tanaka K, Tooyama I, Fujioka Y, Sakai N, et al. Successful vitrification of whole juvenile testis in the critically endangered cyprinid honmoroko (*Gnathopogon caeruleus*). *Zygote* 2017;25(5):652–61. <https://doi.org/10.1017/S0967199417000430>.
- [207] Thuwanut P, Thongphakdee A, Sommanustweechai A, Siriaroonrat B, Chatdarong K. A case report concerning male gametes rescued from a Siamese Eld's deer (*Rucervus eldi siamensis*): post-thawed testicular and epididymal sperm quality and heterologous zona pellucida binding ability. *J Vet Med Sci* 2013;75(1):123–5. <https://doi.org/10.1292/jvms.11-0491>.
- [208] Olaciregui M, Luño V, Domingo P, González N, Gil L. In vitro developmental ability of ovine oocytes following intracytoplasmic injection with freeze-dried spermatozoa. *Sci Rep* 2017;7(1):1096. <https://doi.org/10.1038/>

- s41598-017-00583-0.
- [209] Leibo SP, Songsasen N. Cryopreservation of gametes and embryos of non-domestic species. *Theriogenology* 2002;57(1):303–26. [https://doi.org/10.1016/s0093-691x\(01\)00673-2](https://doi.org/10.1016/s0093-691x(01)00673-2).
- [210] Clarke AG. The Frozen Ark Project: the role of zoos and aquariums in preserving the genetic material of threatened animals. *Int Zoo Yearbk* 2009;43(1):222–30. <https://doi.org/10.1111/j.1748-1090.2008.00074.x>.
- [211] Costa Mafalda, Bruford Michael W. The Frozen Ark Project: biobanking and endangered animal samples for conservation and research: inside Ecology. <https://insideecology.com/2018/01/12/the-frozen-ark-project-biobanking-endangered-animal-samples-for-conservation-and-research/>; 2018.
- [212] Fickel J, Wagener A, Ludwig A. Semen cryopreservation and the conservation of endangered species. *Eur J Wildl Res* 2007;53(2):81–9. <https://doi.org/10.1007/s10344-007-0089-z>.
- [213] U.K. Research and Innovation. BBR-CryoArks: enhancing frozen collections for non-model and endangered animal taxa. <https://gtr.ukri.org/projects?ref=BB%2FR015260%2F1>. [Accessed 1 October 2020].
- [214] Tunstall T, Kock R, Vahala J, Diekhans M, Fiddes I, Armstrong J, et al. Evaluating recovery potential of the northern white rhinoceros from cryopreserved somatic cells. *Genome Res* 2018;28(6):780–8. <https://doi.org/10.1101/gr.227603.117>.