

## Perspective

## Epigenetic regulatory layers in the 3D nucleus

Andréa Willemin,<sup>1,2,\*</sup> Dominik Szabó,<sup>1,2</sup> and Ana Pombo<sup>1,2,\*</sup><sup>1</sup>Max-Delbrück-Center for Molecular Medicine in the Helmholtz Association (MDC), Berlin Institute for Medical Systems Biology (BIMSB), Epigenetic Regulation and Chromatin Architecture Group, Berlin, Germany<sup>2</sup>Humboldt-Universität zu Berlin, Institute for Biology, Berlin, Germany\*Correspondence: [andrea.willemin@mdc-berlin.de](mailto:andrea.willemin@mdc-berlin.de) (A.W.), [ana.pombo@mdc-berlin.de](mailto:ana.pombo@mdc-berlin.de) (A.P.)<https://doi.org/10.1016/j.molcel.2023.12.032>

## SUMMARY

Nearly 7 decades have elapsed since Francis Crick introduced the *central dogma of molecular biology*, as part of his *ideas on protein synthesis*, setting the fundamental rules of sequence information transfer from DNA to RNAs and proteins. We have since learned that gene expression is finely tuned in time and space, due to the activities of RNAs and proteins on regulatory DNA elements, and through cell-type-specific three-dimensional conformations of the genome. Here, we review major advances in genome biology and discuss a set of *ideas on gene regulation* and highlight how various biomolecular assemblies lead to the formation of structural and regulatory features within the nucleus, with roles in transcriptional control. We conclude by suggesting further developments that will help capture the complex, dynamic, and often spatially restricted events that govern gene expression in mammalian cells.

## INTRODUCTION

In 1956, 3 years after the publication of the double helix structure of DNA, Crick introduced a concept that became known as the *central dogma of molecular biology*. In *ideas on protein synthesis*,<sup>1</sup> Crick sets the major axiom of the directionality of sequence information transfer during protein biogenesis—that is, how information is passed from the DNA to messenger RNA (mRNA) and to proteins (Figure 1A). Crick's central dogma states that “once information has got into a protein it can't get out again,” where the term “information” refers to a given sequence and its transfer from one substrate, being either deoxyribonucleotides (in DNA), ribonucleotides (in RNA), or amino acids (in proteins), into another. Although DNA can be copied into RNA by the process of transcription, proteins are generated by the translation of an RNA intermediate (generally mRNA), which converts the nucleotide sequence into its amino acid counterpart, following a correspondence referred to as the genetic code. Other types of information transfer exist, namely, DNA replication, reverse transcription of RNA into DNA (such as for retroviruses prior to their integration in the genome), and RNA self-replication.

Whereas the central dogma still explains the directional transfer of sequence information between DNA, RNA, and protein, we are far from understanding how regulatory mechanisms arise from the activity of nuclear factors within the complex nuclear landscape (Figure 1B). Genome regulation involves crosstalks and feedback loops between DNA, RNAs, and proteins that occur in the three-dimensional (3D) cell nucleus (Figure 1C). Chromatin adopts a hierarchy of folding states, from whole chromosomes occupying preferred territories to specific chromatin interactions, and to local nucleosome positioning.<sup>2,3</sup> Nuclear macroscale properties also contribute to gene regulation and include diverse membraneless microenvironments composed

of proteins and RNAs. Finding general rules of genome regulation remains especially difficult because 3D genome structure and gene regulation are highly dependent on the genomic context of each gene, are specific of cell type, developmental and disease states, and can vary over time. Deciphering how the homeostatic control of gene activation and repression is achieved in a timely and spatially defined manner is fundamental because the first events that dictate cellular function and identity occur through gene regulation and have the greatest impact on development, disease onset, and progression.

In this article, we explore how proteins, RNAs, and regulatory DNA elements organize the genome and regulate the expression of genes. The distinction between co-observed events with functional, cause-and-effect implications, and coincidental co-occurrences remains challenging. The advent of multimodal approaches that simultaneously measure different parameters in the same cell promises to deliver molecular information that can connect, at the single-cell level, genome structure and function to address current challenges.

## DISCOVERING REGULATORY DNA ELEMENTS AND FUNCTIONAL ncRNAs


Core to our current knowledge of genome function are the seminal discoveries on DNA sequence elements, or *cis*-regulatory elements (CREs), which hold regulatory activity and are found throughout the genome (Figure 2). The release of the first assemblies of mammalian genomes (e.g., the human genome<sup>4</sup>) made evident that protein-coding sequences account for only a minor fraction of the genome. The number of protein-coding genes in humans is currently estimated to about 20,000. Within genes, only a small fraction of DNA sequences code for amino acids (within exons), and the remainder (introns) can contain CREs that regulate the same or other genes. Less than 1.2% of the total




**A** Ideas on protein synthesis (Crick, 1956) Transfer = Sequence



**B** Ideas on gene regulation (1885-present) Transfer = Transcriptional regulation



**C** Molecular and structural parameters of gene regulation



**Figure 1. From sequence information transfer from DNA to RNA and proteins to genome regulation**

(A) Schematic representation of sequence information transfer between DNA, RNAs, and proteins (adapted from Crick's *ideas on protein synthesis*, 1956). DNA can be replicated or transcribed into RNA. Translation generates proteins from mRNA templates. Solid arrows indicate the protein biogenesis route and its directionality. Dashed arrows point to DNA replication, RNA reverse transcription, and RNA self-replication.

(B) Abundant inputs from proteins, RNAs, and DNA elements organize the genome and control gene expression at the transcriptional level (*ideas on gene regulation*, collection of evidence from 1885, with Carl Rabl's landmark publication on the territorial organization of chromosomes, to present). Note the reverse directionality of regulatory information transfer compared with sequence information transfer. CREs, *cis*-regulatory elements; DNAm, DNA methylation; ncRNAs, non-coding RNAs; CTCF, CCCTC-binding factor; NPCs, nuclear pore complexes; HP/CBX, heterochromatin/chromobox proteins; RNA Pol II, RNA polymerase II; PTMs, post-translational modifications; TFs, transcription factors.

(C) Main molecular (left) and structural (right) parameters of gene regulation.

3 billion base pairs in the haploid human genome sequence encodes proteins.<sup>5</sup> In contrast, model organisms, such as fly, worm, and yeast, have smaller genomes more densely packed with protein-coding sequences, and their gene expression is more locally regulated.

The development of microarray and next generation sequencing technologies has been instrumental in deciphering how gene regulatory information is stored in the genome. Major efforts such as in the Encyclopedia of DNA Elements (ENCODE) Consortium have revealed that most (>90%) of the genome is transcribed in at least one of the analyzed cell types, although sometimes at low levels.<sup>6</sup> Notably, the fraction of the genome transcribed in individual cell types is around 15%.<sup>7</sup> Genome-wide studies also found that the transcribed genome comprises many newly discovered non-coding RNAs (ncRNAs) that hold regulatory functions, such as long ncRNAs, enhancer RNAs, microRNAs, and circular RNAs.


Apart from the major types of ncRNAs, around 95% of all transcribed genomic locations serve as a template for transcripts whose possible functions remain obscure.<sup>6,8</sup> Tens of millions of distinct transcripts emanating from several hundreds of thousands of genomic regions have been identified across all the assayed tissue types. Pervasive intergenic transcripts (i.e., not falling in exons nor introns) display similar tissue-specific expression patterns between human and chimpanzee, suggesting that such transcription events are under positive selection and have biological functions.<sup>9</sup> Moreover, short pervasive transcripts associated with gene promoters and termini correlate with the expression of the protein-coding genes that they delineate

and have conserved synteny (maintenance of linear genomic arrangement) between human and mouse.<sup>10</sup> The potential biological significance of “pervasive transcription” remains highly debated owing to the difficulty in assessing it unequivocally. It is possible that the process of transcription, rather than the transcripts themselves, holds regulatory roles, for instance, through chromatin remodeling as a result of RNA polymerase II (RNA Pol II) transcription.<sup>7</sup> New approaches are needed to assess the function of pervasive transcription from a mechanistic viewpoint and in a genome-wide manner, to distinguish whether specific ncRNAs or their transcription regulate gene expression, or are simply by-products of other events.

**TOWARD COMPLETE ATLASES OF CIS-REGULATORY DNA ELEMENTS**

The ENCODE project has mapped candidate CREs in the human genome. Most CREs were categorized as promoters or enhancers, if they are directly adjacent to their target genes or regulate distal genes, respectively.<sup>11</sup> CREs are generally bound by multiple transcription factors (TFs) and other chromatin-associated proteins.<sup>12–14</sup> In the current model of CRE function, specific TFs known as pioneer factors bind to high-affinity DNA motifs, thought as “anchors,” to trigger the opening of chromatin, forming a “core.” Subsequent binding and assembly of other TFs and co-factors around the core, through weaker protein-DNA interactions, multivalent associations between proteins, or with ncRNAs (Figure 3), are thought to further modulate the transcriptional dynamics of specific genes.<sup>13</sup>

The latest ENCODE report assigned approximately 8% of the human genome as putative *cis*-regulatory sequences.<sup>15</sup> The



**Figure 2. Gene regulatory information code**  
CREs cover a substantial fraction of the genome and regulate transcription of target genes. CREs can be grouped into enhancers and promoters, which regulate transcription of target genes across large linear genomic distances. CREs are either cell-type specific or common to multiple cell types and can be located in extragenic or intragenic genomic regions.

total fraction of the genome that contains *cis*-regulatory information remains under intense debate, partly because we lack a complete understanding of how different regulatory activities act either directly on DNA (e.g., TF binding), through transcription itself or non-coding transcripts.<sup>5,7,16</sup>

In sum, evidence accumulated in the last 20 years suggests that most of the genome, rather than being “junk,” has regulatory functions in at least one cell type or tissue, either by encoding functional RNAs, providing binding sites for TFs, or through other less understood genome regulatory effects, such as pervasive transcription. Regulatory regions are selectively deployed in individual cellular or tissue contexts to modulate specific protein-coding gene transcription and provides a fertile ground for new regulations to evolve. Understanding the regulatory genome is also crucial beyond basic research, for healthcare applications because disease-associated sequence variants often overlap with cell-type-specific putative CREs in non-coding DNA.<sup>17</sup>

### SPATIAL ORGANIZATION OF THE GENOME: WHEN 3D STRUCTURE MEETS FUNCTION

Transcriptional regulation occurs within the 3D space of the nucleus, where hierarchically organized layers of genome conformation have evolved to allow for physical contacts between non-coding regulatory DNA elements and their cognate promoters, for spatial segregation of active and inactive chromatin regions, and for preferred positions of genomic sequences relative to different biochemical nuclear microenvironments. These layers ensure appropriate expression of housekeeping and cell-identity genes in time and space and allow for adequate cellular responses to differentiation cues and other stimuli. In the following sections, we introduce different layers of 3D genome structure and review their roles in gene regulation.

#### TADs

The development of genome-wide methods that combine ligation of proximal DNA fragments with high-throughput sequencing has led to the discovery of topologically associating domains (TADs).<sup>18–20</sup> TADs are sub-megabase to megabase-sized genomic regions with preferred local chromatin interactions.<sup>19–22</sup> Each TAD is flanked by boundary regions, which are enriched for binding sites of the TF CCCTC-binding factor (CTCF) and for cohesin binding, short repetitive sequences, and housekeeping genes.<sup>18,19,23</sup> Most studies on TAD formation have focused on the roles of the cohesin complex and CTCF.<sup>24,25</sup> Cohesin is a multi-protein complex that extrudes DNA<sup>26</sup> until it finds DNA-bound CTCF especially at binding sites

with convergent orientations, which are often present at TAD boundaries,<sup>27</sup> thereby forming loops.

TADs were initially proposed to be cell-type invariant with conservation across mammalian species,<sup>18,28,29</sup> but wide variations between cell types are reported in recent comparisons of TAD border positions.<sup>30–32</sup> Single-cell imaging shows that TAD-like structures exist in single cells; yet, they are positioned at a wide range of different genomic locations in different cells with average positions that match ligation-based results from bulk cell populations.<sup>33</sup> This suggests that TADs reflect conformations of highest probability, while allowing contacts between genomic regions located in different TADs to some degree. Supporting this view, cohesin depletion leads to loss of preferred TAD positions, whereas variable TAD-like domains are still seen in single cells.


#### E-P communication within and between TADs

The contact enrichment of TADs and barrier activity of their boundaries is thought to relate with the capacity of insulator DNA sequences to restrict enhancer-promoter (E-P) communication.<sup>34</sup> In this view, enhancers preferentially communicate with genes in the same TAD while being isolated from genes in adjacent or distant TADs.<sup>35</sup> Elegant examples at developmental gene loci show how specific genomic rearrangements that affect TADs and/or their boundaries can alter insulation and lead to abnormal E-P communication and gene misregulation.<sup>36–38</sup> However, the TAD-centric view for general *cis*-regulation is challenged by observations that many E-P (and promoter-promoter) interactions occur across domain boundaries and can efficiently activate gene expression.<sup>32,39,40</sup> Furthermore, TAD disruption after acute cohesin or CTCF depletion in the range of hours results in only modest changes in nascent transcription in mouse embryonic stem cells (approximately 50 differentially expressed genes, mostly associated with differentiation and pluripotency),<sup>41</sup> whereas cohesin or CTCF removal for several days affects the expression of hundreds to, at most, few thousands of genes.<sup>24,25</sup> In line with these observations, loop extrusion may promote contacts between only a subset of gene promoters, especially those directly bound by CTCF, and their enhancers, whereas a number of enhancers and promoters interact independently of CTCF/cohesin.<sup>42,43</sup>


*Cis*-regulatory communication is increasingly recognized as a complex phenomenon often specific of genomic context.<sup>44</sup> E-P interactions can occur in the absence of gene expression, independently of tissue or developmental stage,<sup>45,46</sup> implying that productive E-P communication relies on additional events, such as enhancer activation from primed or poised enhancer states. Conversely, enhancer activity is often broader in time and space than the activity of the target genes, and gene

### Mechanisms class I: Regulation via *cis*-regulatory sequences



#### *De novo* recruitment of Pol II



#### Activation of poised Pol II



#### Regulatory transcription events



**Figure 3. Transcriptional regulation mechanisms class I: regulation via CREs**

CREs are binding sites for transcription factors (TFs), which bring together enhancers and their target gene promoters. *De novo* recruitment of transcriptional machinery including RNA polymerase II (RNA Pol II) and co-factors leads to transcription of the target gene. RNA Pol II can be bound to gene promoters and enhancers in a poised state. Through recruitment of release factors, poised promoters become active to allow transcriptional activation. Regulatory activity can be achieved through transcription of non-coding RNAs, which can modulate chromatin-modifying enzymes, influence local transcription regulators, or promote the formation of multi-molecular assemblies in the nucleus. Enhancer-promoter communication is dynamic, and gene activity is not always correlated with protein- or RNA-mediated spatial proximity.

activation at the right time and place can depend on physical E-P interactions.<sup>47,48</sup> Gene activation may also occur without changes in E-P proximity,<sup>49</sup> or coincide with increased E-P distance.<sup>50</sup> Spatiotemporal E-P regulatory communication is far from being fully elucidated, may be more transient or dynamic than initially thought, and likely consists of a combination of different context-dependent molecular mechanisms governed by diverse chromatin factors, including both proteins and RNAs.

### Heterochromatin and euchromatin states

A growing body of work suggests that membraneless nuclear sub-compartments play important roles in organizing chromatin and are associated with gene activation or repression (Figure 4).<sup>51</sup>

On a global level, chromatin is often classified as euchromatin and heterochromatin. Euchromatin represents the transcriptionally active, decompacted form of chromatin and tends to physically localize toward the nuclear interior, including around splicing speckles, or associates with nuclear pore complexes (NPCs).<sup>52–54</sup> The main epigenetic features of euchromatin are acetylation of histone H3 at residue K27 and trimethylation of H3K4. Transcriptionally active genes can associate with splicing speckles, including genes from different chromosomes<sup>55–57</sup> or with each other in transcription factories.<sup>58–62</sup>

Heterochromatin is characterized by higher levels of compaction, tends to contain silent genes, and acts as a repressive epigenetic environment.<sup>63</sup> In most cell types, heterochromatin predominantly associates with the nuclear lamina, a mesh of proteins located at the inner nuclear membrane, and with the surface of nucleoli.<sup>64</sup> Heterochromatin is further divided into constitutive and facultative sub-types, which have distinctive molecular properties and sub-nuclear localization.

### Constitutive heterochromatin

Constitutive heterochromatin is present around centromeric and telomeric regions of chromosomes, is enriched in repetitive sequences, and is largely invariant between cell types or throughout differentiation.<sup>63</sup> At the molecular level, constitu-


tive heterochromatin is dependent on H3K9 methylation and the chromodomain-containing heterochromatin protein 1 (HP1).<sup>65,66</sup> HP1 proteins interact with Suv39h1 and Suv39h2, the enzymes that deposit H3K9me3, thereby stabilizing them onto chromatin for efficient heterochromatin formation and/or maintenance.<sup>67</sup> In mammalian cells, the constitutive heterochromatin state is orchestrated by the HP1 $\alpha$  homolog (chromobox 5 [CBX5]),<sup>68</sup> a protein also likely involved in physically connecting constitutive heterochromatin with the nuclear lamina through lamin B receptor binding.<sup>69–71</sup> HP1 $\alpha$  interacts with H3K9me3 with an affinity and specificity that are modulated by phosphorylation of an intrinsically disordered region (IDR) in the HP1 $\alpha$  protein,<sup>72,73</sup> which also interacts with nuclear RNAs.<sup>74,75</sup> The proper targeting of HP1 $\alpha$  to chromatin depends on the interaction of HP1 $\alpha$  with H3K9me3 and with repeat-containing RNAs at telomeric and pericentromeric regions.<sup>75–78</sup>

HP1 $\alpha$  can form phase separated liquid droplets and compact DNA *in vitro*, which are also dependent on phosphorylation of its IDR.<sup>79</sup> The phase separation potential of HP1 $\alpha$  has been proposed to confer a single mechanism by which constitutive heterochromatin forms and excludes other factors, namely, components of the transcriptional machinery,<sup>80</sup> depending on their chemical properties.<sup>81</sup> However, recent *in vivo* evidence using cultured mouse fibroblasts shows that HP1 $\alpha$  can efficiently mediate repression of a reporter gene independently of droplet formation.<sup>82</sup> The compacted state of heterochromatin may instead result from bridging-type interactions that generate a collapsed chromatin conformation through polymer phase separation.<sup>83,84</sup>

### Facultative heterochromatin and Polycomb repression mechanisms

In contrast with constitutive heterochromatin, facultative heterochromatin is more scattered throughout the genome, is highly specific of cell type, and dynamically changes during development and in disease.<sup>54</sup> Facultative heterochromatin occurs at genomic regions occupied by Polycomb-group proteins and their associated histone modifications, H3K27me3 and

**Mechanisms class II: Regulation via chromatin state and biophysical properties**



**Figure 4. Transcriptional regulation mechanisms class II: regulation via chromatin state and biophysical properties**

Heterochromatin and euchromatin exhibit distinct compaction states, epigenetic signatures, host different regulatory complexes and associate with different nuclear environments, which fine-tune embedded *cis*-regulatory elements. Proximity to repressive nuclear landmarks such as the lamina and the nucleolus preserves gene inactivity. Proximity to active nuclear landmarks such as nuclear speckles that contain RNA binding proteins and splicing factors is correlated with transcription potential. Chromatin occurs in loosely packed, decondensed (or “melted”) state or in a more compact condensed state relating with transcriptional activity.

H2AK119ub1, and is a typical feature of developmentally regulated genes (e.g., *Hox* loci) that are dynamically activated or repressed in response to developmental cues.<sup>85</sup> CBX2, a subunit of Polycomb repressive complex 1 (PRC1), which contains a chromodomain, is thought to have an orchestrating role in facultative heterochromatin, similar to the role of HP1 $\alpha$  in constitutive heterochromatin.<sup>86,87</sup> Canonical PRC1 complexes contain one of the five different CBX proteins: CBX2, CBX4, CBX6, CBX7, or CBX8.<sup>88</sup> CBX7/8 are recruited to H3K27me3, which is deposited by PRC2, and mediate PRC1 targeting to this mark.<sup>89</sup> PRC1 catalyzes the monoubiquitination of histone H2A at residue K119 (H2AK119ub1), which stimulates PRC2 recruitment and further deposition of H3K27me3,<sup>90</sup> creating a feedback loop that promotes the formation of facultative heterochromatin domains.<sup>91</sup> PRCs also widely co-associate with a poised form of RNA Pol II at developmental and metabolic genes poised for activation, often at their promoters but, in some cases, also throughout their coding regions.<sup>92–94</sup> The interplay between DNA, chromatin

modifiers, and RNAs is a known feature of PRC1 recruitment, as reported in the context of developmental regulator gene repression in mouse embryonic stem cells.<sup>95,96</sup> Long ncRNAs (lncRNAs) have also been implicated in Polycomb function<sup>97</sup> and may promote the initial chromatin recruitment of PRC1/2, similar to *de novo* HP1 $\alpha$  recruitment at prospective constitutive heterochromatin.<sup>98–100</sup>

Phase separation has also been shown to participate in facultative heterochromatin through CBX2. The IDR of CBX2 is required for the formation of liquid-like PRC1 condensates *in vivo*,<sup>87,101</sup> chromatin compaction *in vitro*,<sup>102</sup> and is essential for normal axial patterning in mice,<sup>103</sup> although loss of PRC1 condensates does not interfere with pre-existing compacted chromatin.<sup>86</sup> Collectively, these observations suggest a stepwise process in which PRC1-CBX2 condensates indirectly mediate the compaction of facultative heterochromatin. First, PRC1 is recruited to H3K27me3-marked regions via CBX7/8 (or via lncRNAs), and subsequently mobilized into nearby

PRC1-CBX2 condensates.<sup>87</sup> This second event is thought to be mediated by another component of canonical PRC1, the Polyhomeotic subunits, which can bridge PRC1 complexes with one another.<sup>104,105</sup> Lastly, deposition of H2AK119ub1 by PRC1 and/or further H3K27me3 through feedback with PRC2 within the condensate would trigger chromatin compaction.<sup>86</sup> A complete characterization of the compaction mechanism remains to be established, especially whether chromatin segments harboring repressive histone modifications spontaneously compact *in vivo*, or require molecular adaptors to bridge modified nucleosomes.

### Mapping the nuclear position of heterochromatin domains

The sub-nuclear spatial distribution of heterochromatic domains has been extensively investigated thanks to sequencing techniques such as DNA adenine methyltransferase identification (DamID), which uses a fusion between a DNA adenine methyltransferase and a protein of interest to detect its close chromatin association in a genome-wide manner.<sup>106</sup> Genomic stretches that preferentially localize close to the nuclear lamina, referred to as “lamina-associated domains” or LADs, were mapped using DamID of nuclear lamins or their interacting proteins.<sup>107</sup> LADs are contiguous genomic stretches of 0.01–10 Mb in length in mammals, which can cover 30%–50% of the genome depending on cell type.<sup>108</sup> LADs contain repressed or lowly expressed genes and are characterized by high levels of H3K9me2/3, and the presence of H3K27me3 mainly at their borders with inter-LAD regions.<sup>109–111</sup> LADs are also partially related to domains that associate with the surface of nucleoli, called “nucleolus-associated domains” or NADs,<sup>111–114</sup> which were initially identified by DNA sequencing methods in biochemical isolated nucleoli.<sup>63,115</sup> For brevity, we focus below on the case of LADs, although similar principles are thought to apply to NADs.

LADs can be classified according to their variability between cell types, as constitutive LADs (cLADs) or facultative LADs (fLADs).<sup>116–118</sup> fLADs represent more than half of all LADs, contain on average more genes than cLADs, and their cell-type-specific dynamics likely play important roles in gene regulation. The association or detachment of genomic regions with or from the lamina often correlate with down- or upregulation of the genes they contain, respectively.<sup>118–121</sup> However, artificially tethering loci to the lamina has only modest effects on gene expression, often leading to incomplete repression,<sup>122</sup> variable outcomes from gene to gene,<sup>123</sup> or no effect.<sup>124</sup> Furthermore, gene promoters located at LADs that are less tightly embedded within the lamina can evade repression with varying degrees.<sup>125</sup> Lastly, forcing expression of genes within LADs causes detachment of those transcription units from the lamina, whereas, conversely, gene inactivation can result in their lamina engagement, suggesting that the act of gene transcription may also counteract local lamina association.<sup>126</sup> An appealing possibility is that the global organization of LADs (and possibly of NADs) could result from the biophysical properties of heterochromatin and the sub-nuclear environment, such as mutual affinity between H3K9me2/3-rich chromatin and the lamina,<sup>51</sup> which would nonetheless remain susceptible to additional forces at the local level, especially transcriptional activity. The nuclear

(or nucleolar) periphery may also convey a non-homogeneous medium for gene repressiveness, at different depths or locations along the periphery, which could explain the variable consequences of lamina association or dissociation on gene activity.

### Radial gene positioning and proximity to splicing speckles

Expression at specific genes also correlates more generally with nuclear radial positioning of loci.<sup>62,127,128</sup> Typical examples are the progressive relocation of the  $\beta$ -globin gene to a more internal nuclear position during erythroid maturation, when its expression increases,<sup>62</sup> the immunoglobulin genes and other immune loci during lymphocyte differentiation,<sup>129,130</sup> and the brain-derived neurotrophic factor (*Bdnf*) gene during neuronal activation.<sup>131</sup> Other studies did not find a clear correlation between radial chromatin positioning and gene expression status,<sup>132</sup> suggesting that it is not a general determinant for all genes or in all systems and requires further study.

The relative distance of genomic regions to the nuclear lamina and to nuclear splicing speckles has been recently mapped genome-wide by tyramide signal amplification sequencing (TSA-seq), a sequencing-based approach that uses antibody-enzyme conjugates directed against specific nuclear landmarks and quantitatively labels DNA according to its physical distance to the landmark.<sup>133,134</sup> TSA-seq showed that distances of genes to nuclear splicing speckles are inversely correlated with expression levels in various human cell types, with even small distance shifts being closely correlated with gene expression changes between different cell types. From all genes upregulated in one cell type compared with another, about 10% map to regions that increase their proximity to speckles in the cell type in which they are most expressed.

### Chromosome territories

Chromosome territories represent the largest hierarchical layer of chromatin organization in the eukaryotic cell nucleus. On the verge of the 20<sup>th</sup> century, Carl Rabl and Theodor Boveri found using light microscopy that chromosomes occupy preferred locations, largely separated from each other,<sup>135,136</sup> which was later confirmed by laser-UV-microirradiation and more directly by DNA fluorescence *in situ* hybridization (DNA-FISH) experiments.<sup>137–139</sup>

The radial positions of chromosome territories in both pluripotent and differentiated cells correlate with gene density in nearly spherical nuclei (e.g., embryonic stem cells and lymphocytes)<sup>140–142</sup> or chromosome size in non-spherical nuclei.<sup>137</sup> Gene-poor chromosomes (or chromosomal sub-regions) are positioned toward the periphery of the nucleus, whereas gene-rich chromosomes of similar size are generally located in the nuclear interior.<sup>132,143–145</sup>

Chromosome territories are not strictly separated in mammalian cells, where they intermingle with one another as seen by FISH performed on ultrathin nuclear cryosections combined with confocal imaging or electron microscopy,<sup>2</sup> and by chromosome conformation capture experiments followed by polymer modeling.<sup>146</sup> Intermingling varies between cell types, being lower in human embryonic stem cells and cancer cell lines compared with peripheral blood lymphocytes.<sup>2,146</sup> The extent

and specificity of chromosome intermingling changes during activation of resting peripheral blood lymphocytes and upon transcriptional inhibition.<sup>2,147</sup> It remains unclear whether transcription occurs preferentially at the surface or throughout the volume of chromosome territories<sup>2,148,149</sup> and to which degree is trans-chromosomal intermingling cell-type specific<sup>147</sup> or variable within cell populations.

Preferred proximities between different chromosomes have been associated with the propensity for chromosomal translocations in specific cell types,<sup>2,150</sup> a process relevant for unstable cancer genomes and autism spectrum disorders.<sup>151</sup> The mechanisms that drive preferential chromosome territory arrangements may include nuclear shape, preferred associations to the nuclear lamina, nucleoli, or speckles, and transcriptional activity. Further investigation is required to elucidate the specificity and general mechanisms of trans-interactions between chromosomes and their functional implications.

### Gene positioning relative to chromosome territories

The position of individual genes relative to their chromosome territories changes with gene activation and depends on cell type.<sup>148,152,153</sup> Several studies found that a number of highly expressed genes or gene clusters (including *Hox*, *major histocompatibility complex [MHC]*, *epidermal differentiation complex [EDC]*, and *T-cell receptor beta chain [TCRb]*) are present at the surface, or even excluded, from their chromosome territory in functionally relevant cell types.<sup>148,153–156</sup> At the territory surface, loci may be exposed to neighboring chromosomes,<sup>2,153</sup> to the nuclear lamina, or to membraneless nuclear bodies, such as the nucleolus or speckles.

Although gene looping out from chromosome territories can coincide with transcriptional activation and extensive decondensation of the protruding chromatin segment, the three processes are not strictly coupled.<sup>148,153,157</sup> For example, the insertion of the *β-globin* locus control region (LCR) to a gene-rich genomic region of the *Rad23a* locus was sufficient to reposition the locus away from the periphery of its chromosome territory, resulting in increased expression of some, but not all, genes adjacent to the LCR insertion.<sup>158</sup> Conversely, the activation of the *plasminogen activator, urokinase (PLAU)* gene in hepatoblastoma-derived HepG2 cells treated with phorbol esters led to the looping of *PLAU* out from its chromosome territory but without leading to increased probability for transcriptional activity.<sup>148</sup> In contrast, the transcriptional silencing of *PLAU* was highly correlated with its internal location within its territory. These findings suggest that there is more complexity in the relationship of territory positioning of genes and their transcriptional regulation. For some loci, repositioning within their territory may be actively regulated and core to the activation process, possibly by modulating the exposure of specific genomic loci to regulatory cues present in different nuclear microenvironments.<sup>159</sup> The process of gene activation or transcription may itself be sufficient to promote the decondensation and relocation of loci between different sub-regions of each chromosome territories.<sup>160</sup>

### Large-scale chromatin decondensation (melting) events

Large-scale chromatin decondensation was first observed in the form of chromosome puffs that accompany high transcription

levels in polytene chromosomes of fly salivary gland cells.<sup>161,162</sup> A connection between chromatin decondensation and transcription induction was also reported in tandem arrays of the mammary tumor virus promoter and after retinoic-acid-induced *Hoxb* expression.<sup>154,163</sup> More recently, chromatin decondensation events have been detected genome-wide at hundreds of long genes by genome architecture mapping (GAM).<sup>32</sup> Decondensation or “melting” of long genes is highly cell type specific and, in neurons, tends to occur when genes are most expressed and with highest chromatin accessibility levels. Extensive decondensation of long and highly expressed genes has also been explored by FISH in terminally differentiated human cells.<sup>160</sup> Both studies suggest that chromatin decondensation may result from high RNA Pol II occupancy and facilitate access of the transcriptional machinery. Long melting genes also tend to be sensitive to topoisomerase I inhibition,<sup>32,164</sup> which suggests that their high expression levels are particularly dependent on resolution of DNA supercoiling. Chromosome recondensation can accompany gene repression, which likely contributes to a reduced probability of subsequent activation by modulating access and binding of nuclear factors and enzymatic activities.<sup>165,166</sup>


### Higher-order transcriptional assemblies involving TFs and super-enhancers

TFs hold key roles in the dynamic interplay between 3D genome organization and gene expression control.<sup>167</sup> Many TFs undergo weak multivalent interactions through their IDRs, producing condensates that can have characteristics of liquid-liquid phase separation (LLPS),<sup>168</sup> and have been proposed to concentrate general components of the transcriptional machinery, such as Mediator and RNA Pol II.<sup>169</sup> Multivalent interactions of TFs and co-factors may help stabilize them onto DNA for efficient gene activation.<sup>170</sup> TF-mediated microenvironments often occur at, and are possibly driven by, groups of CREs known as super-enhancers<sup>171,172</sup> and can be modulated by RNAs.<sup>173</sup> Higher-order chromatin associations between super-enhancer elements and highly transcribed genomic segments spanning distances of tens of megabases have also been observed,<sup>39,174</sup> but the roles and the preferred partners of such long-range associations remain unexplored.

### THE PAST, PRESENT, AND FUTURE AVENUES IN GENOME RESEARCH

In this final section, we elaborate on ongoing efforts to address some of the remaining “missing links” needed to gain a comprehensive understanding of the general processes governing genome structure and function at the molecular level.

Imaging techniques pioneered the field of genome biology for decades, providing invaluable insights, such as detailed preferential genome positioning in the 3D space of the nucleus, and uncovering cell-to-cell variability of genome conformations within cell populations. New-age methods begin to increase the imaging throughput to provide parallel measurements of a larger number of genomic regions, RNAs, and proteins that can be measured at once. In parallel, the rapid development of sequencing-based approaches has revolutionized the 4D



**Figure 5. Comparison of modern techniques to study genome organization**

Methods compared with respect to specific features (from top clockwise): required biological input material, preservation of tissue context information, depth of sub-nuclear spatial resolution, throughput relative to entire genomic sequence, potential for parallel measurements of DNA, RNAs, and proteins (or other biological phenomarkers), breadth of captured 3D genome structures (incl. pairwise and multiway chromatin contacts, radial positioning, and association with nuclear microenvironments; for more details, see Kempfer and Pombo<sup>176</sup>).

nucleome field.<sup>175</sup> The entire genome can now be interrogated across its unidimensional linear sequence with respect to gene expression, histone modifications, chromatin binding, accessibility, and association with specific nuclear landmarks and in three dimensions in terms of multiway chromatin interactions with the rest of the genome and positioning within the nuclear landscape.<sup>176</sup> Most of the emerging sequencing approaches are now adaptable to application in single cells, giving new insights into dynamic and cell-type-specific events. A major remaining challenge in 3D genome research lies in creating, whenever possible, robust frameworks to integrate data from different experimental approaches and resolve contradictions (Figure 5).

Genome-wide interactions of a gene measured by ligation-based techniques are routinely integrated with histone modification data in sequence space, to infer the candidate target genes of putative CREs and vice versa. However, cell-to-cell variability inevitably causes interpretation issues. For example, a specific E-P contact may be detected in a given cell population in which the gene is globally active; yet, it is not trivial whether the E-P contact occurs in cells where the enhancer and/or gene are active, or in other cells where one or both are silent.

Other limitations come with the integration of different types of data (e.g., imaging and sequencing-based), where shared “reference points” may be sparse or even absent. Computational modeling using physics theory and/or machine learning aspire to fill the gap between 3D genome organization and 1D features along the linear sequence by estimating how linear genome features are folded in 3D,<sup>58,177,178</sup> by inferring the coherent binding patterns of chromatin factors along the linear

sequence from chromatin organization data,<sup>179,180</sup> or by deriving models that jointly consider linear features and 3D conformation.<sup>181</sup> Further work is needed to directly test predictions made through modeling approaches beyond correlations with *in vivo* data.

To capture the complex relationship between genome structure and gene regulation, multimodal technologies are needed for simultaneous localization and quantification of DNA interactions, RNAs and proteins, including post-translational modifications (PTMs), in single cells ideally with spatial resolution in real tissues or biopsies. Exciting developments are ongoing in multimodal ligation-based methods (e.g., methyl-Hi-C and, more recently, combining Hi-C with RNA sequencing [RNA-seq]) to enable measurements of pairwise chromatin contacts in parallel with DNA methylation (DNAm) or RNA abundance.<sup>182,183</sup> Sequential FISH has recently mapped genome organization, multiple RNAs, and proteins using successive rounds of probe hybridization and imaging in mouse embryonic stem cells and in brain cells.<sup>57,184</sup> Split-pool recognition of interactions by tag extension (SPRITE) and GAM are also well suited to incorporate measurements of both RNA and protein modalities. SPRITE is a ligation-free technique that relies on splitting, barcoding, and pooling cross-linked chromatin clusters a number of times, followed by their sequencing to measure various aspects of genome organization.<sup>174</sup> SPRITE is applicable in single cells,<sup>185</sup> and can incorporate simultaneous RNA measurements within each chromatin cluster.<sup>186</sup> GAM is a single-cell assay based on sequencing genomic DNA extracted from ultrathin nuclear cryosections from structurally preserved cells, while retaining tissue



spatial information,<sup>32,39</sup> and is compatible by design with parallel measurements of RNAs<sup>187</sup> and proteins. Multimodal genome structure data will be especially valuable to train models of nuclear architecture and/or gene regulation to increase their predictive power, with applications in early prognostics of complex non-coding genetic disorders.

The development of novel integrative and multi-omics approaches also promises invaluable insights on the contributions of TFs to 3D genome structure and gene expression programs, their dynamics during development and alteration in disease. The actions of TFs, and how they cooperate to modulate chromatin architecture and achieve productive communication of enhancers with their target genes, remain largely unknown. Studies on TF function in the context of the spatially organized nucleus represent an exciting direction for the future of research in genome biology, especially how changes in the abundance of developmental TFs act on 3D genome structure throughout lineage commitment.<sup>188</sup>

Almost 70 years ago, Francis Crick laid one of the biggest milestones in molecular biology by proposing a simple set of rules that define the transfer of sequence information between DNA, RNA, and protein, at a time when fundamental concepts such as the genetic code had not yet been described. The advent of single-cell imaging and unbiased epigenomics has taught us that gene expression is regulated by proteins, RNAs, DNA sequences, and chromatin structure acting together in the 3D space of the nucleus with dynamic properties at the single-cell level. Understanding the cooperativity and interdependency between all different mechanisms and factors remains an important task for the next decades to deepen our knowledge on cellular decision-making and homeostasis and to gain insight into human disease inception and response to therapies to forge tomorrow's medicine.

#### ACKNOWLEDGMENTS

A.P. acknowledges support from the Helmholtz Association (Germany), National Institutes of Health Common Fund 4D Nucleome Program grant 1UM1HG011585, the Deutsche Forschungsgemeinschaft (DFG; German Research Foundation) under Germany's Excellence Strategy—EXC-2049 390688087, and Priority Program SPP2202 (PO 861/4-1).

#### DECLARATION OF INTERESTS

A.P. holds a patent on “Genome Architecture Mapping”: Pombo, A., Edwards, P.A.W., Nicodemi, M., Scialdone, A., and Beagrie, R.A. Patent PCT/EP2015/079413 (2015).

#### REFERENCES

- Crick, F. (1956). Ideas on protein synthesis. <https://profiles.nlm.nih.gov/101584582X65>.
- Branco, M.R., and Pombo, A. (2006). Intermingling of chromosome territories in interphase suggests role in translocations and transcription-dependent associations. *PLoS Biol.* *4*, e138.
- Singh, A.K., and Mueller-Planitz, F. (2021). Nucleosome positioning and spacing: from mechanism to function. *J. Mol. Biol.* *433*, 166847.
- Lander, E.S., Linton, L.M., Birren, B., Nusbaum, C., Zody, M.C., Baldwin, J., Devon, K., Dewar, K., Doyle, M., FitzHugh, W., et al. (2001). Initial sequencing and analysis of the human genome. *Nature* *409*, 860–921.
- Hatje, K., Mühlhausen, S., Simm, D., and Kollmar, M. (2019). The protein-coding human genome: annotating high-hanging fruits. *BioEssays* *41*, 1900066.
- Clark, M.B., Amaral, P.P., Schlesinger, F.J., Dinger, M.E., Taft, R.J., Rinn, J.L., Ponting, C.P., Stadler, P.F., Morris, K.V., Morillon, A., et al. (2011). The reality of pervasive transcription. discussion e1001102. *PLoS Biol.* *9*, e1000625.
- Dinger, M.E., Amaral, P.P., Mercer, T.R., and Mattick, J.S. (2009). Pervasive transcription of the eukaryotic genome: functional indices and conceptual implications. *Brief. Funct. Genomic. Proteomic.* *8*, 407–423.
- Perteau, M., Shumate, A., Perteau, G., Varabyou, A., Breitwieser, F.P., Chang, Y.C., Madugundu, A.K., Pandey, A., and Salzberg, S.L. (2018). CHES: a new human gene catalog curated from thousands of large-scale RNA sequencing experiments reveals extensive transcriptional noise. *Genome Biol.* *19*, 208.
- Khaitovich, P., Kelso, J., Franz, H., Visagie, J., Giger, T., Joerchel, S., Petzold, E., Green, R.E., Lachmann, M., and Pääbo, S. (2006). Functionality of intergenic transcription: an evolutionary comparison. *PLoS Genet.* *2*, e171.
- Kapranov, P., Cheng, J., Dike, S., Nix, D.A., Dutttagupta, R., Willingham, A.T., Stadler, P.F., Hertel, J., Hackermüller, J., Hofacker, I.L., et al. (2007). RNA maps reveal new RNA classes and a possible function for pervasive transcription. *Science* *316*, 1484–1488.
- Hon, C.C., and Carninci, P. (2020). Expanded ENCODE delivers invaluable genomic encyclopedia. *Nature* *583*, 685–686.
- Meuleman, W., Muratov, A., Rynes, E., Halow, J., Lee, K., Bates, D., Diegel, M., Dunn, D., Neri, F., Teodosiadis, A., et al. (2020). Index and biological spectrum of human DNase I hypersensitive sites. *Nature* *584*, 244–251.
- Partridge, E.C., Chhetri, S.B., Prokop, J.W., Ramaker, R.C., Jansen, C.S., Goh, S.T., Mackiewicz, M., Newberry, K.M., Brandsmeier, L.A., Meadows, S.K., et al. (2020). Occupancy maps of 208 chromatin-associated proteins in one human cell type. *Nature* *583*, 720–728.
- Vierstra, J., Lazar, J., Sandstrom, R., Halow, J., Lee, K., Bates, D., Diegel, M., Dunn, D., Neri, F., Haugen, E., et al. (2020). Global reference mapping of human transcription factor footprints. *Nature* *583*, 729–736.
- ENCODE Project Consortium, Moore, J.E., Purcaro, M.J., Pratt, H.E., Epstein, C.B., Shores, N., Adrian, J., Kawli, T., Davis, C.A., Dobin, A., et al. (2020). Expanded encyclopaedias of DNA elements in the human and mouse genomes. *Nature* *583*, 699–710.
- Doolittle, W.F. (2018). We simply cannot go on being so vague about ‘function.’ *Genome Biol.* *19*, 223.
- ENCODE Project Consortium, Snyder, M.P., Gingeras, T.R., Moore, J.E., Weng, Z., Gerstein, M.B., Ren, B., Hardison, R.C., Stamatoyannopoulos, J.A., Graveley, B.R., et al. (2020). Perspectives on ENCODE. *Nature* *583*, 693–698.
- Dixon, J.R., Selvaraj, S., Yue, F., Kim, A., Li, Y., Shen, Y., Hu, M., Liu, J.S., and Ren, B. (2012). Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* *485*, 376–380.
- Nora, E.P., Lajoie, B.R., Schulz, E.G., Giorgetti, L., Okamoto, I., Servant, N., Piolot, T., van Berkum, N.L., Meisig, J., Sedat, J., et al. (2012). Spatial partitioning of the regulatory landscape of the X-inactivation centre. *Nature* *485*, 381–385.
- Sexton, T., Yaffe, E., Kenigsberg, E., Bantignies, F., Leblanc, B., Hoichman, M., Parrinello, H., Tanay, A., and Cavalli, G. (2012). Three-dimensional folding and functional organization principles of the Drosophila genome. *Cell* *148*, 458–472.
- Dostie, J., Richmond, T.A., Arnaout, R.A., Selzer, R.R., Lee, W.L., Honan, T.A., Rubio, E.D., Krumm, A., Lamb, J., Nusbaum, C., et al. (2006). Chromosome conformation capture carbon copy (5C): A massively parallel solution for mapping interactions between genomic elements. *Genome Res.* *16*, 1299–1309.
- Lieberman-Aiden, E., van Berkum, N.L., Williams, L., Imakaev, M., Ragoczy, T., Telling, A., Amit, I., Lajoie, B.R., Sabo, P.J., Dorschner,

- M.O., et al. (2009). Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science* 326, 289–293.
23. Sun, J.H., Zhou, L., Emerson, D.J., Phyo, S.A., Titus, K.R., Gong, W., Gilgenast, T.G., Beagan, J.A., Davidson, B.L., Tassone, F., et al. (2018). Disease-associated short tandem repeats co-localize with chromatin domain boundaries. *Cell* 175, 224–238.e15.
  24. Nora, E.P., Goloborodko, A., Valton, A.-L., Gibcus, J.H., Uebersohn, A., Abdennur, N., Dekker, J., Mirny, L.A., and Bruneau, B.G. (2017). Targeted degradation of CTCF decouples local insulation of chromosome domains from genomic compartmentalization. *Cell* 169, 930–944.e22.
  25. Schwarzer, W., Abdennur, N., Goloborodko, A., Pekowska, A., Fudenberg, G., Loe-Mie, Y., Fonseca, N.A., Huber, W., Haering, C.H., Mirny, L., et al. (2017). Two independent modes of chromatin organization revealed by cohesin removal. *Nature* 551, 51–56.
  26. Davidson, I.F., Bauer, B., Goetz, D., Tang, W., Wutz, G., and Peters, J.M. (2019). DNA loop extrusion by human cohesin. *Science* 366, 1338–1345.
  27. Rao, S.S.P., Huntley, M.H., Durand, N.C., Stamenova, E.K., Bochkov, I.D., Robinson, J.T., Sanborn, A.L., Machol, I., Omer, A.D., Lander, E.S., et al. (2014). A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell* 159, 1665–1680.
  28. Lazar, N.H., Nevenon, K.A., O’Connell, B., McCann, C., O’Neill, R.J., Green, R.E., Meyer, T.J., Okhovat, M., and Carbone, L. (2018). Epigenetic maintenance of topological domains in the highly rearranged gibbon genome. *Genome Res.* 28, 983–997.
  29. Vietri Rudan, M., Barrington, C., Henderson, S., Ernst, C., Odom, D.T., Tanay, A., and Hadjir, S. (2015). Comparative Hi-C reveals that CTCF underlies evolution of chromosomal domain architecture. *Cell Rep.* 10, 1297–1309.
  30. Loof, G., Szabó, D., Garg, V., Kukalev, A., Zea-Redondo, L., Kempfer, R., Sparks, T.M., Zhang, Y., Thieme, C.J., Carvalho, S., et al. (2022). 3D genome topologies distinguish pluripotent epiblast and primitive endoderm cells in the mouse blastocyst. Preprint at bioRxiv.
  31. McArthur, E., and Capra, J.A. (2021). Topologically associating domain boundaries that are stable across diverse cell types are evolutionarily constrained and enriched for heritability. *Am. J. Hum. Genet.* 108, 269–283.
  32. Winick-Ng, W., Kukalev, A., Harabula, I., Zea-Redondo, L., Szabó, D., Meijer, M., Serebreni, L., Zhang, Y., Bianco, S., Chiariello, A.M., et al. (2021). Cell-type specialization is encoded by specific chromatin topologies. *Nature* 599, 684–691.
  33. Bintu, B., Mateo, L.J., Su, J.-H., Sinnott-Armstrong, N.A., Parker, M., Kinrot, S., Yamaya, K., Boettiger, A.N., and Zhuang, X. (2018). Super-resolution chromatin tracing reveals domains and cooperative interactions in single cells. *Science* 362, eaau1783.
  34. Kolovos, P., Knoch, T.A., Grosveld, F.G., Cook, P.R., and Papanonis, A. (2012). Enhancers and silencers: an integrated and simple model for their function. *Epigenetics Chromatin* 5, 1.
  35. Beagan, J.A., and Phillips-Cremins, J.E. (2020). On the existence and functionality of topologically associating domains. *Nat. Genet.* 52, 8–16.
  36. Lupiáñez, D.G., Kraft, K., Heinrich, V., Krawitz, P., Brancati, F., Klopocki, E., Horn, D., Kayserili, H., Opitz, J.M., Laxova, R., et al. (2015). Disruptions of topological chromatin domains cause pathogenic rewiring of gene-enhancer interactions. *Cell* 161, 1012–1025.
  37. Rodríguez-Carballo, E., Lopez-Delisle, L., Zhan, Y., Fabre, P.-J., Beccari, L., El-Idrissi, I., Huynh, T.H.N., Ozadam, H., Dekker, J., and Duboule, D. (2017). The HoxD cluster is a dynamic and resilient TAD boundary controlling the segregation of antagonistic regulatory landscapes. *Genes Dev.* 31, 2264–2281.
  38. Willemin, A., Lopez-Delisle, L., Bolt, C.C., Gadolini, M.-L., Duboule, D., and Rodríguez-Carballo, E. (2021). Induction of a chromatin boundary in vivo upon insertion of a TAD border. *PLoS Genet.* 17, e1009691.
  39. Beagrie, R.A., Scialdone, A., Schueler, M., Kraemer, D.C.A., Chotalia, M., Xie, S.Q., Barbieri, M., de Santiago, I., Lavitas, L.M., Branco, M.R., et al. (2017). Complex multi-enhancer contacts captured by genome architecture mapping. *Nature* 543, 519–524.
  40. Chakraborty, S., Kopitchinski, N., Zuo, Z., Eraso, A., Awasthi, P., Chari, R., Mitra, A., Tobias, I.C., Moorthy, S.D., Dale, R.K., et al. (2023). Enhancer-promoter interactions can bypass CTCF-mediated boundaries and contribute to phenotypic robustness. *Nat. Genet.* 55, 280–290.
  41. Hsieh, T.-H.S., Cattoglio, C., Slobodyanyuk, E., Hansen, A.S., Darzacq, X., and Tjian, R. (2022). Enhancer-promoter interactions and transcription are largely maintained upon acute loss of CTCF, cohesin, WAPL or YY1. *Nat. Genet.* 54, 1919–1932.
  42. Kubo, N., Ishii, H., Xiong, X., Bianco, S., Meitinger, F., Hu, R., Hocker, J.D., Conte, M., Gorkin, D., Yu, M., et al. (2021). Promoter-proximal CTCF binding promotes distal enhancer-dependent gene activation. *Nat. Struct. Mol. Biol.* 28, 152–161.
  43. Thiecke, M.J., Wutz, G., Muhar, M., Tang, W., Bevan, S., Malysheva, V., Stocsits, R., Neumann, T., Zuber, J., Fraser, P., et al. (2020). Cohesin-dependent and -independent mechanisms mediate chromosomal contacts between promoters and enhancers. *Cell Rep.* 32, 107929.
  44. Wurmser, A., and Basu, S. (2022). Enhancer-promoter communication: it’s not just about contact. *Front. Mol. Biosci.* 9, 867303.
  45. Andrey, G., Schöpflin, R., Jerković, I., Heinrich, V., Ibrahim, D.M., Paliou, C., Hochradel, M., Timmermann, B., Haas, S., Vingron, M., et al. (2017). Characterization of hundreds of regulatory landscapes in developing limbs reveals two regimes of chromatin folding. *Genome Res.* 27, 223–233.
  46. Paliou, C., Guckelberger, P., Schöpflin, R., Heinrich, V., Esposito, A., Chiariello, A.M., Bianco, S., Annunziatella, C., Helmuth, J., Haas, S., et al. (2019). Preformed chromatin topology assists transcriptional robustness of Shh during limb development. *Proc. Natl. Acad. Sci. USA* 116, 12390–12399.
  47. Kragestein, B.K., Spielmann, M., Paliou, C., Heinrich, V., Schöpflin, R., Esposito, A., Annunziatella, C., Bianco, S., Chiariello, A.M., Jerković, I., et al. (2018). Dynamic 3D chromatin architecture contributes to enhancer specificity and limb morphogenesis. *Nat. Genet.* 50, 1463–1473.
  48. Robson, M.I., Ringel, A.R., and Mundlos, S. (2019). Regulatory landscaping: how enhancer-promoter communication is sculpted in 3D. *Mol. Cell* 74, 1110–1122.
  49. Alexander, J.M., Guan, J., Li, B., Maliskova, L., Song, M., Shen, Y., Huang, B., Lomvardas, S., and Weiner, O.D. (2019). Live-cell imaging reveals enhancer-dependent Sox2 transcription in the absence of enhancer proximity. *eLife* 8, e41769.
  50. Benabdallah, N.S., Williamson, I., Illingworth, R.S., Kane, L., Boyle, S., SenGupta, D., Grimes, G.R., Therizols, P., and Bickmore, W.A. (2019). Decreased enhancer-promoter proximity accompanying enhancer activation. *Mol. Cell* 76, 473–484.e7.
  51. Bhat, P., Honson, D., and Guttman, M. (2021). Nuclear compartmentalization as a mechanism of quantitative control of gene expression. *Nat. Rev. Mol. Cell Biol.* 22, 653–670.
  52. Faber, G.P., Nadav-Eliyahu, S., and Shav-Tal, Y. (2022). Nuclear speckles – a driving force in gene expression. *J. Cell Sci.* 135, jcs259594.
  53. Kuhn, T.M., and Capelson, M. (2019). Nuclear pore proteins in regulation of chromatin state. *Cells* 8, 1414.
  54. Morrison, O., and Thakur, J. (2021). Molecular complexes at euchromatin, heterochromatin and centromeric chromatin. *Int. J. Mol. Sci.* 22, 6922.
  55. Brown, J.M., Green, J., das Neves, R.P., Wallace, H.A.C., Smith, A.J.H., Hughes, J., Gray, N., Taylor, S., Wood, W.G., Higgs, D.R., et al. (2008). Association between active genes occurs at nuclear speckles and is modulated by chromatin environment. *J. Cell Biol.* 182, 1083–1097.
  56. Shopland, L.S., Johnson, C.V., Byron, M., McNeil, J., and Lawrence, J.B. (2003). Clustering of multiple specific genes and gene-rich R-bands around SC-35 domains: evidence for local euchromatic neighborhoods. *J. Cell Biol.* 162, 981–990.

57. Takei, Y., Yun, J., Zheng, S., Ollikainen, N., Pierson, N., White, J., Shah, S., Thomassie, J., Suo, S., Eng, C.-H.L., et al. (2021). Integrated spatial genomics reveals global architecture of single nuclei. *Nature* **590**, 344–350.
58. Barbieri, M., Xie, S.Q., Torlai Triglia, E., Chiariello, A.M., Bianco, S., de Santiago, I., Branco, M.R., Rueda, D., Nicodemi, M., and Pombo, A. (2017). Active and poised promoter states drive folding of the extended HoxB locus in mouse embryonic stem cells. *Nat. Struct. Mol. Biol.* **24**, 515–524.
59. Kimura, H., and Sato, Y. (2022). Imaging transcription elongation dynamics by new technologies unveils the organization of initiation and elongation in transcription factories. *Curr. Opin. Cell Biol.* **74**, 71–79.
60. Martin, S., and Pombo, A. (2003). Transcription factories: quantitative studies of nanostructures in the mammalian nucleus. *Chromosome Res.* **11**, 461–470.
61. Osborne, C.S., Chakalova, L., Brown, K.E., Carter, D., Horton, A., Debrand, E., Goyenechea, B., Mitchell, J.A., Lopes, S., Reik, W., et al. (2004). Active genes dynamically colocalize to shared sites of ongoing transcription. *Nat. Genet.* **36**, 1065–1071.
62. Ragoczy, T., Bender, M.A., Telling, A., Byron, R., and Groudine, M. (2006). The locus control region is required for association of the murine  $\beta$ -globin locus with engaged transcription factories during erythroid maturation. *Genes Dev.* **20**, 1447–1457.
63. Bizhanova, A., and Kaufman, P.D. (2021). Close to the edge: heterochromatin at the nucleolar and nuclear peripheries. *Biochim. Biophys. Acta Gene Regul. Mech.* **1864**, 194666.
64. Pombo, A., and Dillon, N. (2015). Three-dimensional genome architecture: players and mechanisms. *Nat. Rev. Mol. Cell Biol.* **16**, 245–257.
65. Montavon, T., Shukeir, N., Erikson, G., Engist, B., Onishi-Seebacher, M., Ryan, D., Musa, Y., Mittler, G., Meyer, A.G., Genoud, C., et al. (2021). Complete loss of H3K9 methylation dissolves mouse heterochromatin organization. *Nat. Commun.* **12**, 4359.
66. Saksouk, N., Simboeck, E., and Déjardin, J. (2015). Constitutive heterochromatin formation and transcription in mammals. *Epigenetics Chromatin* **8**, 3.
67. Maeda, R., and Tachibana, M. (2022). HP1 maintains protein stability of H3K9 methyltransferases and demethylases. *EMBO Rep.* **23**, e35381.
68. Nielsen, A.L., Oulad-Abdeghani, M., Ortiz, J.A., Remboutsika, E., Chambon, P., and Losson, R. (2001). Heterochromatin formation in mammalian cells: interaction between histones and HP1 proteins. *Mol. Cell* **7**, 729–739.
69. Haws, S.A., Simandi, Z., Barnett, R.J., and Phillips-Cremins, J.E. (2022). 3D genome, on repeat: higher-order folding principles of the heterochromatinized repetitive genome. *Cell* **185**, 2690–2707.
70. Nikolakaki, E., Mylonis, I., and Giannakouros, T. (2017). Lamin B receptor: interplay between structure, function and localization. *Cells* **6**, 28.
71. Ye, Q., Callebaut, I., Pezhman, A., Courvalin, J.-C., and Worman, H.J. (1997). Domain-specific interactions of human HP1-type chromodomain proteins and inner nuclear membrane protein LBR. *J. Biol. Chem.* **272**, 14983–14989.
72. Hiragami-Hamada, K., Shinmyozu, K., Hamada, D., Tatsu, Y., Uegaki, K., Fujiwara, S., and Nakayama, J. (2011). N-terminal phosphorylation of HP1 $\alpha$  promotes its chromatin binding. *Mol. Cell Biol.* **31**, 1186–1200.
73. Nishibuchi, G., Machida, S., Osakabe, A., Murakoshi, H., Hiragami-Hamada, K., Nakagawa, R., Fischle, W., Nishimura, Y., Kurumizaka, H., Tagami, H., et al. (2014). N-terminal phosphorylation of HP1 $\alpha$  increases its nucleosome-binding specificity. *Nucleic Acids Res.* **42**, 12498–12511.
74. Maison, C., Bailly, D., Peters, A.H.F.M., Quivy, J.-P., Roche, D., Taddei, A., Lachner, M., Jenuwein, T., and Almouzni, G. (2002). Higher-order structure in pericentric heterochromatin involves a distinct pattern of histone modification and an RNA component. *Nat. Genet.* **30**, 329–334.
75. Muchardt, C., Guillemé, M., Seeler, J.-S., Trouche, D., Dejean, A., and Yaniv, M. (2002). Coordinated methyl and RNA binding is required for heterochromatin localization of mammalian HP1 $\alpha$ . *EMBO Rep.* **3**, 975–981.
76. Biscotti, M.A., Canapa, A., Forconi, M., Olmo, E., and Barucca, M. (2015). Transcription of tandemly repetitive DNA: functional roles. *Chromosome Res.* **23**, 463–477.
77. Maison, C., Bailly, D., Roche, D., Montes de Oca, R.M., Probst, A.V., Vassias, I., Dingli, F., Lombard, B., Loew, D., Quivy, J.-P., et al. (2011). SUMOylation promotes de novo targeting of HP1 $\alpha$  to pericentric heterochromatin. *Nat. Genet.* **43**, 220–227.
78. Roach, R.J., Garavis, M., González, C., Jameson, G.B., Filichev, V.V., and Hale, T.K. (2020). Heterochromatin protein 1 $\alpha$  interacts with parallel RNA and DNA G-quadruplexes. *Nucleic Acids Res.* **48**, 682–693.
79. Larson, A.G., Elnatan, D., Keenen, M.M., Trnka, M.J., Johnston, J.B., Burlingame, A.L., Agard, D.A., Redding, S., and Narlikar, G.J. (2017). Liquid droplet formation by HP1 $\alpha$  suggests a role for phase separation in heterochromatin. *Nature* **547**, 236–240.
80. Wang, L., Gao, Y., Zheng, X., Liu, C., Dong, S., Li, R., Zhang, G., Wei, Y., Qu, H., Li, Y., et al. (2019). Histone modifications regulate chromatin compartmentalization by contributing to a phase separation mechanism. *Mol. Cell* **76**, 646–659.e6.
81. Guthmann, M., Burton, A., and Torres-Padilla, M.-E. (2019). Expression and phase separation potential of heterochromatin proteins during early mouse development. *EMBO Rep.* **20**, e47952.
82. Erdel, F., Rademacher, A., Vlijm, R., Tünnermann, J., Frank, L., Weinmann, R., Schweigert, E., Yserentant, K., Hummert, J., Bauer, C., et al. (2020). Mouse heterochromatin adopts digital compaction states without showing hallmarks of HP1-driven liquid-liquid phase separation. *Mol. Cell* **78**, 236–249.e7.
83. Barbieri, M., Chotalia, M., Fraser, J., Lavitas, L.-M., Dostie, J., Pombo, A., and Nicodemi, M. (2012). Complexity of chromatin folding is captured by the strings and binders switch model. *Proc. Natl. Acad. Sci. USA* **109**, 16173–16178.
84. Erdel, F., and Rippe, K. (2018). Formation of chromatin subcompartments by phase separation. *Biophys. J.* **114**, 2262–2270.
85. Rodríguez-Carballo, E., Lopez-Delisle, L., Yakushiji-Kaminatsui, N., Ullate-Agote, A., and Duboule, D. (2019). Impact of genome architecture on the functional activation and repression of Hox regulatory landscapes. *BMC Biol.* **17**, 55.
86. Eeftens, J.M., Kapoor, M., Michieletto, D., and Brangwynne, C.P. (2021). Polycomb condensates can promote epigenetic marks but are not required for sustained chromatin compaction. *Nat. Commun.* **12**, 5888.
87. Tatavosian, R., Kent, S., Brown, K., Yao, T., Duc, H.N., Huynh, T.N., Zhen, C.Y., Ma, B., Wang, H., and Ren, X. (2019). Nuclear condensates of the Polycomb protein chromobox 2 (CBX2) assemble through phase separation. *J. Biol. Chem.* **294**, 14511–1463.
88. Chittock, E.C., Latwiel, S., Miller, T.C.R., and Müller, C.W. (2017). Molecular architecture of polycomb repressive complexes. *Biochem. Soc. Trans.* **45**, 193–205.
89. Zhen, C.Y., Tatavosian, R., Huynh, T.N., Duc, H.N., Das, R., Kokotovic, M., Grimm, J.B., Lavis, L.D., Lee, J., Mejia, F.J., et al. (2016). Live-cell single-molecule tracking reveals co-recognition of H3K27me3 and DNA targets polycomb Cbx7-PRC1 to chromatin. *eLife* **5**, e17667.
90. Kalb, R., Latwiel, S., Baymaz, H.I., Jansen, P.W.T.C., Müller, C.W., Vermeulen, M., and Müller, J. (2014). Histone H2A monoubiquitination promotes histone H3 methylation in Polycomb repression. *Nat. Struct. Mol. Biol.* **21**, 569–571.
91. Blackledge, N.P., and Klose, R.J. (2021). The molecular principles of gene regulation by Polycomb repressive complexes. *Nat. Rev. Mol. Cell Biol.* **22**, 815–833.
92. Brookes, E., de Santiago, I., Hebenstreit, D., Morris, K.J., Carroll, T., Xie, S.Q., Stock, J.K., Heidemann, M., Eick, D., Nozaki, N., et al. (2012). Polycomb associates genome-wide with a specific RNA polymerase II variant, and regulates metabolic genes in ESCs. *Cell Stem Cell* **10**, 157–170.

93. Ferrai, C., Torlai Triglia, E., Risner-Janiczek, J.R., Rito, T., Rackham, O.J., de Santiago, I., Kukalev, A., Nicodemi, M., Akalin, A., Li, M., et al. (2017). RNA polymerase II primes Polycomb-repressed developmental genes throughout terminal neuronal differentiation. *Mol. Syst. Biol.* **13**, 946.
94. Tee, W.W., Shen, S.S., Oksuz, O., Narendra, V., and Reinberg, D. (2014). ERK1/2 activity promotes chromatin features and RNAPII phosphorylation at developmental promoters in mouse ESCs. *Cell* **156**, 678–690.
95. Kanhere, A., Viiri, K., Araújo, C.C., Rasaiyaah, J., Bouwman, R.D., Whyte, W.A., Pereira, C.F., Brookes, E., Walker, K., Bell, G.W., et al. (2010). Short RNAs are transcribed from repressed polycomb target genes and interact with polycomb repressive Complex-2. *Mol. Cell* **38**, 675–688.
96. Skourti-Stathaki, K., Torlai Triglia, E.T., Warburton, M., Voigt, P., Bird, A., and Pombo, A. (2019). R-loops enhance polycomb repression at a subset of developmental regulator genes. *Mol. Cell* **73**, 930–945.e4.
97. Statello, L., Guo, C.-J., Chen, L.-L., and Huarte, M. (2021). Gene regulation by long non-coding RNAs and its biological functions. *Nat. Rev. Mol. Cell Biol.* **22**, 96–118.
98. Long, Y., Hwang, T., Gooding, A.R., Goodrich, K.J., Rinn, J.L., and Cech, T.R. (2020). RNA is essential for PRC2 chromatin occupancy and function in human pluripotent stem cells. *Nat. Genet.* **52**, 931–938.
99. Ray, M.K., Wiskow, O., King, M.J., Ismail, N., Ergun, A., Wang, Y., Plys, A.J., Davis, C.P., Kathrein, K., Sadreyev, R., et al. (2016). CAT7 and cat7l long non-coding RNAs tune polycomb repressive Complex 1 function during human and zebrafish development. *J. Biol. Chem.* **291**, 19558–19572.
100. Schertzer, M.D., Bracerros, K.C.A., Starmer, J., Cherney, R.E., Lee, D.M., Salazar, G., Justice, M., Bischoff, S.R., Cowley, D.O., Ariel, P., et al. (2019). lncRNA-induced spread of Polycomb controlled by genome architecture, RNA abundance, and CpG island DNA. *Mol. Cell* **75**, 523–537.e10.
101. Plys, A.J., Davis, C.P., Kim, J., Rizki, G., Keenen, M.M., Marr, S.K., and Kingston, R.E. (2019). Phase separation of Polycomb-repressive complex 1 is governed by a charged disordered region of CBX2. *Genes Dev.* **33**, 799–813.
102. Grau, D.J., Chapman, B.A., Garlick, J.D., Borowsky, M., Francis, N.J., and Kingston, R.E. (2011). Compaction of chromatin by diverse Polycomb group proteins requires localized regions of high charge. *Genes Dev.* **25**, 2210–2221.
103. Lau, M.S., Schwartz, M.G., Kundu, S., Savol, A.J., Wang, P.I., Marr, S.K., Grau, D.J., Schorderet, P., Sadreyev, R.I., Tabin, C.J., et al. (2017). Mutation of a nucleosome compaction region disrupts Polycomb-mediated axial patterning. *Science* **355**, 1081–1084.
104. Isono, K., Endo, T.A., Ku, M., Yamada, D., Suzuki, R., Sharif, J., Ishikura, T., Toyoda, T., Bernstein, B.E., and Koseki, H. (2013). SAM domain polymerization links subnuclear clustering of PRC1 to gene silencing. *Dev. Cell* **26**, 565–577.
105. Seif, E., Kang, J.J., Sasseville, C., Senkovich, O., Kaltashov, A., Boulter, E.L., Kapur, I., Kim, C.A., and Francis, N.J. (2020). Phase separation by the polyhomeotic sterile alpha motif compartmentalizes Polycomb Group proteins and enhances their activity. *Nat. Commun.* **11**, 5609.
106. van Steensel, B., Delrow, J., and Henikoff, S. (2001). Chromatin profiling using targeted DNA adenine methyltransferase. *Nat. Genet.* **27**, 304–308.
107. Guelen, L., Pagie, L., Brassat, E., Meuleman, W., Faza, M.B., Talhout, W., Eussen, B.H., de Klein, A., Wessels, L., de Laat, W., et al. (2008). Domain organization of human chromosomes revealed by mapping of nuclear lamina interactions. *Nature* **453**, 948–951.
108. van Steensel, B., and Belmont, A.S. (2017). Lamina-associated domains: links with chromosome architecture, heterochromatin, and gene repression. *Cell* **169**, 780–791.
109. Harr, J.C., Luperchio, T.R., Wong, X., Cohen, E., Wheelan, S.J., and Reddy, K.L. (2015). Directed targeting of chromatin to the nuclear lamina is mediated by chromatin state and A-type lamins. *J. Cell Biol.* **208**, 33–52.
110. See, K., Kiseleva, A.A., Smith, C.L., Liu, F., Li, J., Poleshko, A., and Epstein, J.A. (2020). Histone methyltransferase activity programs nuclear peripheral genome positioning. *Dev. Biol.* **466**, 90–98.
111. Vertii, A., Ou, J., Yu, J., Yan, A., Pagès, H., Liu, H., Zhu, L.J., and Kaufman, P.D. (2019). Two contrasting classes of nucleolus-associated domains in mouse fibroblast heterochromatin. *Genome Res.* **29**, 1235–1249.
112. Bizhanova, A., Yan, A., Yu, J., Zhu, L.J., and Kaufman, P.D. (2020). Distinct features of nucleolus-associated domains in mouse embryonic stem cells. *Chromosoma* **129**, 121–139.
113. van Koningsbruggen, S., Gierliński, M., Schofield, P., Martin, D., Barton, G.J., Ariyurek, Y., den Dunnen, J.T., and Lamond, A.I. (2010). High-resolution whole-genome sequencing reveals that specific chromatin domains from most human chromosomes associate with nucleoli. *Mol. Biol. Cell* **21**, 3735–3748.
114. Peng, T., Hou, Y., Meng, H., Cao, Y., Wang, X., Jia, L., Chen, Q., Zheng, Y., Sun, Y., Chen, H., et al. (2023). Mapping nucleolus-associated chromatin interactions using nucleolus Hi-C reveals pattern of heterochromatin interactions. *Nat. Commun.* **14**, 350.
115. Gupta, S., and Santoro, R. (2020). Regulation and roles of the nucleolus in embryonic stem cells: from ribosome biogenesis to genome organization. *Stem Cell Rep.* **15**, 1206–1219.
116. Kind, J., Pagie, L., de Vries, S.S., Nahidiazar, L., Dey, S.S., Bienko, M., Zhan, Y., Lajoie, B., de Graaf, C.A., Amendola, M., et al. (2015). Genome-wide maps of nuclear lamina interactions in single human cells. *Cell* **163**, 134–147.
117. Meuleman, W., Peric-Hupkes, D., Kind, J., Beaudry, J.B., Pagie, L., Kellis, M., Reinders, M., Wessels, L., and van Steensel, B. (2013). Constitutive nuclear lamina–genome interactions are highly conserved and associated with A/T-rich sequence. *Genome Res.* **23**, 270–280.
118. Peric-Hupkes, D., Meuleman, W., Pagie, L., Bruggeman, S.W.M., Solovei, I., Brugman, W., Gräf, S., Flicek, P., Kerkhoven, R.M., Lohuizen, M.V., et al. (2010). Molecular maps of the reorganization of genome-nuclear lamina interactions during differentiation. *Mol. Cell* **38**, 603–613.
119. Borsos, M., Perricone, S.M., Schauer, T., Pontabry, J., de Luca, K.L., de Vries, S.S., Ruiz-Morales, E.R., Torres-Padilla, M.-E., and Kind, J. (2019). Genome–lamina interactions are established de novo in the early mouse embryo. *Nature* **569**, 729–733.
120. Robson, M.I., de las Heras, J.I., Czapiewski, R., Lê Thành, P., Booth, D.G., Kelly, D.A., Webb, S., Kerr, A.R.W., and Schirmer, E.C. (2016). Tissue-specific gene repositioning by muscle nuclear membrane proteins enhances repression of critical developmental genes during myogenesis. *Mol. Cell* **62**, 834–847.
121. Shah, P.P., Keough, K.C., Gjoni, K., Santini, G.T., Abdill, R.J., Wickramasinghe, N.M., Dundes, C.E., Karnay, A., Chen, A., Salomon, R.E.A., et al. (2023). An atlas of lamina-associated chromatin across twelve human cell types reveals an intermediate chromatin subtype. *Genome Biol.* **24**, 16.
122. Reddy, K.L., Zullo, J.M., Bertolino, E., and Singh, H. (2008). Transcriptional repression mediated by repositioning of genes to the nuclear lamina. *Nature* **452**, 243–247.
123. Finlan, L.E., Sproul, D., Thomson, I., Boyle, S., Kerr, E., Perry, P., Ylstra, B., Chubb, J.R., and Bickmore, W.A. (2008). Recruitment to the nuclear periphery can alter expression of genes in human cells. *PLoS Genet.* **4**, e1000039.
124. Kumaran, R.I., and Spector, D.L. (2008). A genetic locus targeted to the nuclear periphery in living cells maintains its transcriptional competence. *J. Cell Biol.* **180**, 51–65.
125. Leemans, C., van der Zwalm, M.C.H., Brueckner, L., Comoglio, F., van Schaik, T., Pagie, L., van Arensbergen, J., and van Steensel, B. (2019). Promoter-intrinsic and local chromatin features determine gene repression in LADs. *Cell* **177**, 852–864.e14.
126. Brueckner, L., Zhao, P.A., van Schaik, T., Leemans, C., Sima, J., Peric-Hupkes, D., Gilbert, D.M., and van Steensel, B. (2020). Local rewiring

- of genome–nuclear lamina interactions by transcription. *EMBO J.* 39, e103159.
127. Takizawa, T., Gudla, P.R., Guo, L., Lockett, S., and Misteli, T. (2008). Allele-specific nuclear positioning of the monoallelically expressed astrocyte marker GFAP. *Genes Dev.* 22, 489–498.
  128. Williams, R.R.E., Azuara, V., Perry, P., Sauer, S., Dvorkina, M., Jørgensen, H., Roix, J., McQueen, P., Misteli, T., Merckenschlager, M., et al. (2006). Neural induction promotes large-scale chromatin reorganization of the *Mash1* locus. *J. Cell Sci.* 119, 132–140.
  129. Kim, S.H., McQueen, P.G., Lichtman, M.K., Shevach, E.M., Parada, L.A., and Misteli, T. (2004). Spatial genome organization during T-cell differentiation. *Cytogenet. Genome Res.* 105, 292–301.
  130. Kosak, S.T., Skok, J.A., Medina, K.L., Riblet, R., Le Beau, M.M., Fisher, A.G., and Singh, H. (2002). Subnuclear compartmentalization of immunoglobulin loci during lymphocyte development. *Science* 296, 158–162.
  131. Walczak, A., Szczepankiewicz, A.A., Ruszczycki, B., Magalska, A., Zamylnska, K., Dzwonek, J., Wilczek, E., Zybura-Broda, K., Rylski, M., Malinowska, M., et al. (2013). Novel higher-order epigenetic regulation of the *BDNF* gene upon seizures. *J. Neurosci.* 33, 2507–2511.
  132. Küpper, K., Kölbl, A., Biener, D., Dittrich, S., von Hase, J., Thormeyer, T., Fiegler, H., Carter, N.P., Speicher, M.R., Cremer, T., et al. (2007). Radial chromatin positioning is shaped by local gene density, not by gene expression. *Chromosoma* 116, 285–306.
  133. Chen, Y., Zhang, Y., Wang, Y., Zhang, L., Brinkman, E.K., Adam, S.A., Goldman, R., van Steensel, B., Ma, J., and Belmont, A.S. (2018). Mapping 3D genome organization relative to nuclear compartments using TSA-seq as a cytological ruler. *J. Cell Biol.* 217, 4025–4048.
  134. Zhang, L., Zhang, Y., Chen, Y., Gholamalamdari, O., Wang, Y., Ma, J., and Belmont, A.S. (2021). TSA-seq reveals a largely conserved genome organization relative to nuclear speckles with small position changes tightly correlated with gene expression changes. *Genome Res.* 31, 251–264.
  135. Boveri, T. (1909). Die Blastomerenkerne von *Ascaris megalcephala* und die Theorie der Chromosomenindividualität. *Arch. Zellforsch.* 3, 181–268.
  136. Rabl, C. (1885). Über Zelltheilung. *Morphol. Jahrb.* 10, 214–330.
  137. Bolzer, A., Kreth, G., Solovei, I., Koehler, D., Saracoglu, K., Fauth, C., Müller, S., Eils, R., Cremer, C., Speicher, M.R., et al. (2005). Three-dimensional maps of all chromosomes in human male fibroblast nuclei and prometaphase rosettes. *PLoS Biol.* 3, e157.
  138. Cremer, T., and Cremer, M. (2010). Chromosome territories. *Cold Spring Harb. Perspect. Biol.* 2, a003889.
  139. Cremer, T., Cremer, C., Baumann, H., Luedtke, E.K., Sperling, K., Teuber, V., and Zorn, C. (1982). Rabl's model of the interphase chromosome arrangement tested in Chinese hamster cells by premature chromosome condensation and laser-UV-microbeam experiments. *Hum. Genet.* 60, 46–56.
  140. Boyle, S., Gilchrist, S., Bridger, J.M., Mahy, N.L., Ellis, J.A., and Bickmore, W.A. (2001). The spatial organization of human chromosomes within the nuclei of normal and emerin-mutant cells. *Hum. Mol. Genet.* 10, 211–219.
  141. Cremer, M., Küpper, K., Wagler, B., Wizelman, L., von Hase, J., Weiland, Y., Kreja, L., Diebold, J., Speicher, M.R., and Cremer, T. (2003). Inheritance of gene density–related higher order chromatin arrangements in normal and tumor cell nuclei. *J. Cell Biol.* 162, 809–820.
  142. Wiblin, A.E., Cui, W., Clark, A.J., and Bickmore, W.A. (2005). Distinctive nuclear organization of centromeres and regions involved in pluripotency in human embryonic stem cells. *J. Cell Sci.* 118, 3861–3868.
  143. Croft, J.A., Bridger, J.M., Boyle, S., Perry, P., Teague, P., and Bickmore, W.A. (1999). Differences in the localization and morphology of chromosomes in the human nucleus. *J. Cell Biol.* 145, 1119–1131.
  144. Federico, C., Scavo, C., Cantarella, C.D., Motta, S., Saccone, S., and Bernardi, G. (2006). Gene-rich and gene-poor chromosomal regions have different locations in the interphase nuclei of cold-blooded vertebrates. *Chromosoma* 115, 123–128.
  145. Goetze, S., Mateos-Langerak, J., Gierman, H.J., de Leeuw, W., Giromus, O., Indemans, M.H.G., Koster, J., Ondrej, V., Versteeg, R., and van Driel, R. (2007). The three-dimensional structure of human interphase chromosomes is related to the transcriptome map. *Mol. Cell. Biol.* 27, 4475–4487.
  146. Tan, L., Xing, D., Chang, C.H., Li, H., and Xie, X.S. (2018). Three-dimensional genome structures of single diploid human cells. *Science* 361, 924–928.
  147. Branco, M.R., Branco, T., Ramirez, F., and Pombo, A. (2008). Changes in chromosome organization during PHA-activation of resting human lymphocytes measured by cryo-FISH. *Chromosome Res.* 16, 413–426.
  148. Ferrai, C., Xie, S.Q., Luraghi, P., Munari, D., Ramirez, F., Branco, M.R., Pombo, A., and Crippa, M.P. (2010). Poised transcription factories prime silent uPA gene prior to activation. *PLoS Biol.* 8, e1000270.
  149. Shah, S., Takei, Y., Zhou, W., Lubeck, E., Yun, J., Eng, C.-H.L., Koulou, N., Cronin, C., Karp, C., Liaw, E.J., et al. (2018). Dynamics and spatial genomics of the nascent transcriptome by intron seqFISH. *Cell* 174, 363–376.e16.
  150. Parada, L.A., and Misteli, T. (2002). Chromosome positioning in the interphase nucleus. *Trends Cell Biol.* 12, 425–432.
  151. Loviglio, M.N., Leleu, M., Männik, K., Passeggeri, M., Giannuzzi, G., van der Werf, I., Waszak, S.M., Zazhytska, M., Roberts-Caldeira, I., Gheldof, N., et al. (2017). Chromosomal contacts connect loci associated with autism, BMI and head circumference phenotypes. *Mol. Psychiatry* 22, 836–849.
  152. Hepperger, C., Mannes, A., Merz, J., Peters, J., and Dietzel, S. (2008). Three-dimensional positioning of genes in mouse cell nuclei. *Chromosoma* 117, 535–551.
  153. Volpi, E.V., Chevret, E., Jones, T., Vatcheva, R., Williamson, J., Beck, S., Campbell, R.D., Goldsworthy, M., Powis, S.H., Ragoussis, J., et al. (2000). Large-scale chromatin organization of the major histocompatibility complex and other regions of human chromosome 6 and its response to interferon in interphase nuclei. *J. Cell Sci.* 113, 1565–1576.
  154. Chambeyron, S., and Bickmore, W.A. (2004). Chromatin decondensation and nuclear reorganization of the *HoxB* locus upon induction of transcription. *Genes Dev.* 18, 1119–1130.
  155. Williams, R.R.E., Broad, S., Sheer, D., and Ragoussis, J. (2002). Subchromosomal positioning of the epidermal differentiation complex (EDC) in keratinocyte and lymphoblast interphase nuclei. *Exp. Cell Res.* 272, 163–175.
  156. Yadav, M., Jalan, M., and Srivastava, M. (2022). Enhancer dependent repositioning of *TCRb* locus with respect to the chromosome territory. *J. Mol. Biol.* 434, 167509.
  157. Morey, C., Da Silva, N.R., Perry, P., and Bickmore, W.A. (2007). Nuclear reorganization and chromatin decondensation are conserved, but distinct, mechanisms linked to *Hox* gene activation. *Development* 134, 909–919.
  158. Noordermeer, D., Branco, M.R., Splinter, E., Klous, P., IJcken, W.V., Swagemakers, S., Koutsourakis, M., van der Spek, P., Pombo, A., and de Laat, W. (2008). Transcription and chromatin organization of a house-keeping gene cluster containing an integrated  $\beta$ -globin locus control region. *PLoS Genetics* 4, e1000016.
  159. Mahy, N.L., Perry, P.E., and Bickmore, W.A. (2002). Gene density and transcription influence the localization of chromatin outside of chromosome territories detectable by FISH. *J. Cell Biol.* 159, 753–763.
  160. Leidescher, S., Ribisel, J., Ullrich, S., Feodorova, Y., Hildebrand, E., Galitsyna, A., Bultmann, S., Link, S., Thanisch, K., Mulholland, C., et al. (2022). Spatial organization of transcribed eukaryotic genes. *Nat. Cell Biol.* 24, 327–339.
  161. Daneholt, B. (1975). Transcription in polytene chromosomes. *Cell* 4, 1–9.

162. Zuckerkandl, E. (1976). Gene control in eukaryotes and the c-value paradox “excess” DNA as an impediment to transcription of coding sequences. *J. Mol. Evol.* **9**, 73–104.
163. Müller, W.G., Walker, D., Hager, G.L., and McNally, J.G. (2001). Large-scale chromatin decondensation and recondensation regulated by transcription from a natural promoter. *J. Cell Biol.* **154**, 33–48.
164. King, I.F., Yandava, C.N., Mabb, A.M., Hsiao, J.S., Huang, H.-S., Pearson, B.L., Calabrese, J.M., Starmer, J., Parker, J.S., Magnuson, T., et al. (2013). Topoisomerases facilitate transcription of long genes linked to autism. *Nature* **501**, 58–62.
165. Ashburner, M. (1990). Puffs, genes, and hormones revisited. *Cell* **61**, 1–3.
166. Martin, R.M., and Cardoso, M.C. (2010). Chromatin condensation modulates access and binding of nuclear proteins. *FASEB J.* **24**, 1066–1072.
167. Kim, S., and Shendure, J. (2019). Mechanisms of interplay between transcription factors and the 3D genome. *Mol. Cell* **76**, 306–319.
168. Boija, A., Klein, I.A., Sabari, B.R., Dall’Agnese, A., Coffey, E.L., Zamudio, A.V., Li, C.H., Shrinivas, K., Manteiga, J.C., Hannett, N.M., et al. (2018). Transcription factors activate genes through the phase-separation capacity of their activation domains. *Cell* **175**, 1842–1855.e16.
169. Cho, W.-K., Spille, J.-H., Hecht, M., Lee, C., Li, C., Grube, V., and Cisse, I.I. (2018). Mediator and RNA polymerase II clusters associate in transcription-dependent condensates. *Science* **361**, 412–415.
170. Trojanowski, J., Frank, L., Rademacher, A., Mücke, N., Grigaitis, P., and Rippe, K. (2022). Transcription activation is enhanced by multivalent interactions independent of phase separation. *Mol. Cell* **82**, 1878–1893.e10.
171. Sabari, B.R., Dall’Agnese, A., Boija, A., Klein, I.A., Coffey, E.L., Shrinivas, K., Abraham, B.J., Hannett, N.M., Zamudio, A.V., Manteiga, J.C., et al. (2018). Coactivator condensation at super-enhancers links phase separation and gene control. *Science* **361**, eaar3958.
172. Whyte, W.A., Orlando, D.A., Hnisz, D., Abraham, B.J., Lin, C.Y., Kagey, M.H., Rahl, P.B., Lee, T.I., and Young, R.A. (2013). Master transcription factors and mediator establish super-enhancers at key cell identity genes. *Cell* **153**, 307–319.
173. Henninger, J.E., Oksuz, O., Shrinivas, K., Sagi, I., LeRoy, G., Zheng, M.M., Andrews, J.O., Zamudio, A.V., Lazaris, C., Hannett, N.M., et al. (2021). RNA-mediated feedback control of transcriptional condensates. *Cell* **184**, 207–225.e24.
174. Quinodoz, S.A., Ollikainen, N., Tabak, B., Palla, A., Schmidt, J.M., Detmar, E., Lai, M.M., Shishkin, A.A., Bhat, P., Takei, Y., et al. (2018). Higher-order inter-chromosomal hubs shape 3D genome organization in the nucleus. *Cell* **174**, 744–757.e24.
175. Dekker, J., Alber, F., Aufmkolk, S., Beliveau, B.J., Bruneau, B.G., Belmont, A.S., Bintu, L., Boettiger, A., Calandrelli, R., Distèche, C.M., et al. (2023). Spatial and temporal organization of the genome: current state and future aims of the 4D nucleome project. *Mol. Cell* **83**, 2624–2640.
176. Kempfer, R., and Pombo, A. (2020). Methods for mapping 3D chromosome architecture. *Nat. Rev. Genet.* **21**, 207–226.
177. Olmeda, F., Lohoff, T., Clark, S.J., Benson, L., Krüger, F., Reik, W., and Rulands, S. (2021). Inference of emergent spatio-temporal processes from single-cell sequencing reveals feedback between *de novo* DNA methylation and chromatin condensates. Preprint at bioRxiv.
178. Tan, J., Shenker-Tauris, N., Rodriguez-Hernaez, J., Wang, E., Sakellariopoulos, T., Boccalatte, F., Thandapani, P., Skok, J., Aifantis, I., Fenyö, D., et al. (2023). Cell-type-specific prediction of 3D chromatin organization enables high-throughput in silico genetic screening. *Nat. Biotechnol.* **41**, 1140–1150.
179. Bianco, S., Lupiáñez, D.G., Chiariello, A.M., Annunziatella, C., Kraft, K., Schöpflin, R., Wittler, L., Andrey, G., Vingron, M., Pombo, A., et al. (2018). Polymer physics predicts the effects of structural variants on chromatin architecture. *Nat. Genet.* **50**, 662–667.
180. Conte, M., Fiorillo, L., Bianco, S., Chiariello, A.M., Esposito, A., Musella, F., Flora, F., Abraham, A., and Nicodemi, M. (2022). A polymer physics model to dissect genome organization in healthy and pathological phenotypes. In *Hi-C Data Analysis: Methods and Protocols* Methods in Molecular Biology, S. Bicciato and F. Ferrari, eds. (Springer), pp. 307–316.
181. Coli, D., Orlandini, E., Michieletto, D., and Marenduzzo, D. (2019). Magnetic polymer models for epigenetics-driven chromosome folding. *Phys. Rev. E* **100**, 052410.
182. Li, G., Liu, Y., Zhang, Y., Kubo, N., Yu, M., Fang, R., Kellis, M., and Ren, B. (2019). Joint profiling of DNA methylation and chromatin architecture in single cells. *Nat. Methods* **16**, 991–993.
183. Liu, Z., Chen, Y., Xia, Q., Liu, M., Xu, H., Chi, Y., Deng, Y., and Xing, D. (2023). Linking genome structures to functions by simultaneous single-cell Hi-C and RNA-seq. *Science* **380**, 1070–1076.
184. Takei, Y., Zheng, S., Yun, J., Shah, S., Pierson, N., White, J., Schindler, S., Tischbirek, C.H., Yuan, G.-C., and Cai, L. (2021). Single-cell nuclear architecture across cell types in the mouse brain. *Science* **374**, 586–594.
185. Arrastia, M.V., Jachowicz, J.W., Ollikainen, N., Curtis, M.S., Lai, C., Quinodoz, S.A., Selck, D.A., Ismagilov, R.F., and Guttman, M. (2022). Single-cell measurement of higher-order 3D genome organization with scSPRITE. *Nat. Biotechnol.* **40**, 64–73.
186. Quinodoz, S.A., Jachowicz, J.W., Bhat, P., Ollikainen, N., Banerjee, A.K., Goronzy, I.N., Blanco, M.R., Chovanec, P., Chow, A., Markaki, Y., et al. (2021). RNA promotes the formation of spatial compartments in the nucleus. *Cell* **184**, 5775–5790.e30.
187. Pombo, A., and Sparks, T.M. (2020). Method for nucleic acid detection by oligo hybridization and PCR-based amplification. Patent application EP 20,203,357.7, PCT/EP2021/079393.
188. Stadhouders, R., Vidal, E., Serra, F., Di Stefano, B., Le Dily, F., Quilez, J., Gomez, A., Collombet, S., Berenguer, C., Cuartero, Y., et al. (2018). Transcription factors orchestrate dynamic interplay between genome topology and gene regulation during cell reprogramming. *Nat. Genet.* **50**, 238–249.