

RESEARCH PAPER

UV light exposure versus vitamin D supplementation: A comparison of health benefits and vitamin D metabolism in a pig model

Julia Kühn^{a,*}, Corinna Brandsch^a, Anja C. Bailer^{a,b}, Mikis Kiourtzidis^{a,b}, Frank Hirche^a, Chia-Yu Chen^{c,d,e,f}, Lajos Markó^{c,d,e,f}, Theda U.P. Bartolomaeus^{c,d,e,f}, Ulrike Löber^{c,d,e,f}, Samira Michel^a, Monika Wensch-Dorendorf^a, Sofia K. Forslund-Startceva^{c,d,e,f,g}, Gabriele I. Stangl^{a,b}

^a Institute of Agricultural and Nutritional Sciences, Martin Luther University Halle-Wittenberg, Halle, Saale, Germany

^b Competence Cluster for Nutrition and Cardiovascular Health (nutriCARD), Halle-Jena-Leipzig, Germany

^c Experimental and Clinical Research Center, Max Delbrück Center for Molecular Medicine and Charité-Universitätsmedizin Berlin, Berlin, Germany

^d Charité-Universitätsmedizin Berlin, Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany

^e Max Delbrück Center for Molecular Medicine in the Helmholtz Association (MDC), Berlin, Germany

^f DZHK, German Centre for Cardiovascular Research, Berlin, Germany

^g Structural and Computational Biology Unit, EMBL, Heidelberg, Germany

Received 27 February 2024; received in revised form 12 August 2024; accepted 16 August 2024

Abstract

There is limited data on the effect of UV light exposure versus orally ingested vitamin D₃ on vitamin D metabolism and health. A 4-week study with 16 pigs (as a model for human physiology) was conducted. The pigs were either supplemented with 20 µg/d vitamin D₃ or exposed to UV light for 19 min/d to standardize plasma 25-hydroxyvitamin D₃ levels. Important differences were higher levels of stored vitamin D₃ in skin and subcutaneous fat, higher plasma concentrations of 3-epi-25-hydroxyvitamin D₃ and increases of cutaneous lumisterol₃ in UV-exposed pigs compared to supplemented pigs. UV light exposure compared to vitamin D₃ supplementation resulted in lower hepatic cholesterol, higher circulating plasma nitrite, a marker of the blood pressure-lowering nitric oxide, and a reduction in the release of pro- and anti-inflammatory cytokines from stimulated peripheral blood mononuclear cells. However, plasma metabolome and stool microbiome analyses did not reveal any differences between the two groups. To conclude, the current data show important health relevant differences between oral vitamin D₃ supplementation and UV light exposure. The findings may also partly explain the different vitamin D effects on health parameters obtained from association and intervention studies.

© 2024 The Authors. Published by Elsevier Inc.

This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

Keywords: Vitamin D₃; Lumisterol₃; 3-epi-25-hydroxyvitamin D₃; Cardiovascular risk factors; Immune response; Microbiome.

1. Introduction

Vitamin D plays an essential role in regulating calcium homeostasis and bone health. In recent years, there has been growing interest in the potential non-skeletal health benefits of vitamin D, such as on the immune and cardiovascular system or the intestinal microbiome. Thus, several observational and interventional studies were conducted to investigate the role of vitamin D in disease prevention and therapy. Interestingly, observational studies often found associations between low plasma levels of 25-hydroxyvitamin D (25(OH)D) and increased disease risk such as hy-

perension [1–3], while intervention studies using vitamin D supplements do not seem to lower this risk [4–6].

Vitamin D is known as the “sunshine vitamin” as it is produced in the skin in response to ultraviolet (UV) light exposure, but it can enter the body by oral routes through vitamin D-containing foods or supplements as well. With some exceptions, endogenous vitamin D seems to contribute predominantly to the vitamin D supply in humans [7]. This means that the primary source of vitamin D in individuals included in observational studies comes from UV exposure, while oral vitamin D is the exclusive source of vitamin D in intervention studies. It has been assumed for a long time that orally consumed vitamin D is physiologically equivalent to vitamin D synthesized in the skin, but there are differences in the transport and metabolic fate of dietary and skin-derived vitamin D [8]. Moreover, it is a well-known fact that large quantities of orally ingested vitamin D can have toxic effects,

* Corresponding author at: Julia Kühn, Martin Luther University Halle-Wittenberg, Von-Danckelmann-Platz 2, Halle Saale, Germany, 06120
E-mail address: julia.kuehn@landw.uni-halle.de (J. Kühn).

whereas overdosing through endogenously produced vitamin D is prevented by degradation of excess metabolites in the skin.

Apart from potential differences in the metabolism of oral and endogenously produced vitamin D, sunlight or UV light can also affect health outcomes independent of vitamin D, such as lowering blood pressure or modulating the immune cells [9,10]. It is therefore likely that epidemiological data investigating the association between vitamin D and health outcomes might be biased by UV light exposure. Consequently, it can be assumed that the observed correlation between blood pressure and vitamin D in association studies [1,3,11] is attributable to sunlight or UV light. It may also explain the absent effects of oral vitamin D on blood pressure in intervention studies [4,12,13]. Whether the conflicting results on vitamin D and health parameters between association and intervention studies are due to vitamin D-independent effects of UV light or differences in the efficacy to improve vitamin D status is largely unknown.

The vitamin D status is usually assessed by the measure of 25(OH)D in plasma or serum and does not provide any information on the origin of vitamin D - endogenously synthesized or orally supplied. There is currently no biomarker that allows distinguishing UV light-produced from oral vitamin D. It is further unknown whether UV light affects vitamin D metabolism differently than oral vitamin D.

Due to the need for data on the effects of UV light compared to oral vitamin D, the current study was conducted to investigate the metabolic routes and health implications of dietary and UV light-produced vitamin D in a pig model. The obtained data may help explain the contradictory data from observational and interventional studies on vitamin D and health. To this end, pigs received either dietary vitamin D or were exposed to UV light, and vitamin D metabolism, blood pressure regulators, immunological parameters, microbial stool composition, and several plasma metabolites were assessed. To avoid divergences in the effects between these two interventions that are simply caused by differences in the improvement of vitamin D status, we first conducted a pre-study to identify the UV light exposure time per day that leads to similar 25(OH)D levels as the intake of 20 µg/d vitamin D.

2. Experimental section

To investigate possible differences in the metabolic fate and health effects of oral versus endogenously produced vitamin D, two studies (a pre-study to find the UV light exposure time that is equivalent to oral vitamin D intake and the main study) using pigs as a model were conducted. The study protocol and the animal husbandry were approved by the local council of Saxony-Anhalt (Landesverwaltungsamt, Halle (Saale), Germany; approval number: 42502-2-1527 MLU). Care and handling of the pigs were in accordance with the German animal welfare regulations (Tierschutzgesetz, version of 18 May 2006 and last revised on 1 January 2019 and Tierschutz-Versuchstierverordnung, version of 1 August 2013 and last revised on 8 September 2015). The studies are described in accordance with the ARRIVE guidelines.

The pre- and the main study were performed using female weaned piglets [Piétraîne x (Large White x Landrace)] with an initial age of 6 weeks at baseline and a mean body weight of 12.2 ± 1.73 kg. The studies were conducted in an animal facility that was controlled for temperature and lighting, with a UV-free lighting program from 06:00 am to 06:00 pm and a color temperature of 4,000 K and 80 lux. All pigs had free access to pellet feed and water from nipple drinkers. The individual pen compartments in the facility consisted of concrete slatted floor and plastic floor grids. Before starting the pre-study and the main study, the pigs were vitamin D depleted for 2 weeks by feeding a vita-

min D free pre-starter diet (supplementary Table 1). During the experimental periods, the pigs were daily treated with either 20 µg orally supplemented vitamin D₃ dissolved in 2 ml soybean oil or UV light (complemented by 2 ml of vitamin D-free soybean oil as sham treatment). The soybean oil was administered with an oral dispenser. The basal diet that was fed during the experimental periods contained no analytically detectable vitamin D₃. For the UV light treatment, UV lamps (Exo Terra Reptile UVB 200, Reptilienkosmos, Viersen, Germany) were mounted 95 cm above the floor. The mean emitted UVB intensity (measured on the shoulder height of the pigs) was 17.7 µW/cm². The UV spectrum of the lamps is shown in supplementary Fig. 1. An automated timer was used to control the on/off switch of the UV lamps.

2.1. Experimental design of the pre-study: Identification of the suitable UV light exposure

The pre-study was conducted to ascertain the UV light exposure time, which leads to the same plasma 25(OH)D levels in pigs as feeding 20 µg vitamin D per day. The standardization of 25(OH)D levels was intended to avoid differences in vitamin D metabolism and health parameters between the two groups caused by differences in vitamin D status. Therefore, a total of 20 pigs were randomly assigned to six groups (three to four pigs per group) and were either fed 20 µg/d vitamin D₃, or daily exposed to UV light for 1 min (1.1 mJ/cm²), 5 min (5.3 mJ/cm²), 10 min (10.6 mJ/cm²), 20 min (21.2 mJ/cm²) and 30 min (31.9 mJ/cm²). The animals were housed in groups of 10 pigs per pen and colored ear tags identified the corresponding intervention group. For the daily UV light exposure procedure, the pigs were separated group-wise and brought to a cabin that was equipped with UV lamps. Blood was drawn weekly to monitor the 25(OH)D₃ levels of all pigs over a period of 4 weeks. To this end, blood was collected in heparinized monovettes (Sarstedt, Nümbrecht, Germany) and then centrifuged at 2,000 g for 10 min to obtain plasma. Analyses were conducted to reveal whether there is a linear or quadratic relationship between the 25(OH)D₃ levels in the pigs and the UV light exposure time.

2.2. Experimental design of the main study: investigating physiological differences between oral vitamin D and UV light exposure

The main study was conducted to compare the tissue levels of vitamin D metabolites and health parameters in pigs treated with oral vitamin D₃ or UV light. The UV radiation conditions for the main intervention study were deduced from the pre-study. A total of 16 pigs with an initial age of 4 weeks and a median body weight of 7.45 kg (7.10 kg–8.13 kg) were randomly divided in two groups and housed in two identical pen compartments (8.32 m²). One group was supplemented daily with 20 µg vitamin D₃ and the other group was daily exposed to UV light for 19 min (20.2 mJ/cm²) to produce endogenous vitamin D₃. To this end, the compartment of the UV light-exposed group was equipped with four UV lamps. The experimental period lasted four weeks.

During the intervention period, the pigs were weighted weekly. Blood samples were also taken weekly by venepuncture from the *vena jugularis*. Blood collected in heparinized monovettes (Sarstedt) was then centrifuged at 2,000 g for 10 min to obtain plasma. Five pigs per group were randomly selected to isolate peripheral blood mononuclear cells (PBMC) at baseline (week 0) and at the end of the intervention period (week 4) to analyze cytokine expression. Stool samples were collected at week 0 and week 4 to analyze microbial composition, bile acids and short-chain fatty

acids. The stool samples were collected from the individuals during defecation, stored immediately at -20°C and transferred to -80°C on the next day. At the end of the study, the pigs were food deprived for 12 h, anesthetized by an intramuscular injection of xylazine (Serumwerk, Bernburg, Germany) plus ketamine (Ursotamin, Serumwerk) and a continuous isoflurane/oxygen gas flow before being euthanized by an intracardiac injection of pentobarbital sodium (Release, WDT, Garbsen, Germany) to gather tissue samples. All tissue samples were immediately snap frozen in liquid nitrogen and stored at -80°C until further analyses.

2.3. Isolation and treatment of the PBMC

PBMC were obtained from the blood of 10 randomly selected pigs (five per group) at baseline and after 4 weeks of intervention to analyze the expression of cytokines. The heparinized blood was layered on Histopaque®-1077 (Sigma-Aldrich, Taufkirchen, Germany) to isolate the PBMC by density gradient centrifugation, according to the manufacturer's protocol [14]. The resulting cell pellet was re-suspended in RPMI 1640 medium (Life Technologies, Darmstadt, Germany), supplemented with 5% fetal bovine serum. For the stimulation, PBMC were seeded in 24-multiwell plates (about 4×10^6 cells per well) and incubated with lipopolysaccharide (LPS, 1 $\mu\text{g}/\text{ml}$ RPMI 1640 medium) for 20 h at 37°C and 5% CO_2 . The cells were harvested by centrifugation at 900 g for 5 min. The supernatant was centrifuged a second time (13,000 g at 4°C) to remove cell debris and was stored at -20°C until analyses of the secreted cytokines. PeqGOLD TriFast™ (Peqlab, Erlangen, Germany) was added to the cell pellets and the tubes were stored at -80°C until analysis of mRNA abundance of cytokines by using PCR and analysis of the protein amount by using a NanoDrop Spectrophotometer (Thermo Fisher Scientific Inc., Schwerte, Germany).

2.4. Analysis of vitamin D metabolites

The concentrations of 25(OH) D_3 , vitamin D_3 , 24,25-dihydroxyvitamin D_3 (24,25(OH) $_2\text{D}_3$) and lumisterol $_3$ were analyzed by high performance liquid chromatography-tandem mass spectrometry (HPLC-MS/MS). Before the identification and quantification of these vitamin D metabolites, the plasma samples were spiked with deuterated internal standards (Sigma-Aldrich). The plasma was then hydrolyzed as described elsewhere [15] and transferred to extraction columns (Extrelut NT1, Merck KGaA, Darmstadt, Deutschland). The metabolites were extracted with *n*-hexane/*tert*-butyl methyl ether and the dried eluates were derivatized with 4-phenyl-1,2,4-triazoline-3,5-dione [15]. The plasma samples were injected to the HPLC-MS/MS (1260 Series, Agilent Technologies; QTRAP 5500, Sciex, Darmstadt, Germany) that was equipped with a Poroshell column (EC-C18, 50×4.6 mm 2 , 2.7 μm). For the chromatographic separation, the column temperature was set to 40°C and the mobile phases consisted of (A) acetonitrile and (B) a mixture of acetonitrile/water (1/1, v/v) with 5 mM ammonium formate and 0.1% formic acid. The following gradient was used: 0.0 min, 0% A, 600 $\mu\text{l}/\text{min}$; 2.1 min, 0% A, 600 $\mu\text{l}/\text{min}$; 4.0 min, 23.5% A, 600 $\mu\text{l}/\text{min}$; 6.0 min, 40% A, 600 $\mu\text{l}/\text{min}$; 8.0 min, 60% A, 600 $\mu\text{l}/\text{min}$; 10.0 min, 80% A, 600 $\mu\text{l}/\text{min}$; 11.0 min, 100% A, 600 $\mu\text{l}/\text{min}$; 16.0 min, 100% A, 600 $\mu\text{l}/\text{min}$; 18.0 min, 100% A, 1000 $\mu\text{l}/\text{min}$; 20.0 min, 100% A, 1000 $\mu\text{l}/\text{min}$; 21.0 min, 0% A, 1000 $\mu\text{l}/\text{min}$; 24.0 min, 0% A, 800 $\mu\text{l}/\text{min}$, 25.0 min, 0% A, 600 $\mu\text{l}/\text{min}$. Mass spectrometric conditions, aspects of lumisterol analysis and quantification procedures are described elsewhere [16,17].

The concentration of 3-epi-25(OH) D_3 was assessed in plasma using the MassChrom® kit for HPLC-MS/MS with atmospheric pressure chemical ionization (Chromsystems, Gräfelfing, Germany).

In plasma samples, the limit of quantifications (LOQ) were: vitamin D_3 , 0.26 nmol/l; 24,25(OH) $_2\text{D}_3$, 1.44 nmol/l; 3-epi-25(OH) D_3 , 2.5 nmol/l, lumisterol $_3$, 4 nmol/l.

Tissue samples for the determination of vitamin D_3 , 25(OH) D_3 and lumisterol $_3$ were prepared as described previously [15,18]. The chromatographic separation was performed on a Hypersil ODS C18 column (5 μm , 2.0×150 mm 2 , VDS Optilab, Berlin, Germany) [19]. The LOQ for 25(OH) D_3 in tissue samples was 7 ng/g.

Precision of the HPLC-MS/MS methods was assessed using the MassCheck® controls (Chromsystems) and pooled liver samples. The coefficients of variation for intra- and inter-day precision were all lower than 10%.

The concentration of 1,25-dihydroxyvitamin D_3 (1,25(OH) $_2\text{D}_3$) in the plasma was analysed by a commercial enzyme-linked immunoassay (Immunodiagnostic Systems, Frankfurt am Main, Germany) according to the manufacturer's protocol. The intra-day precision was 5.0%.

2.5. Analysis of the relative mRNA abundance

The relative mRNA abundance of genes involved in vitamin D metabolism and tissue distribution as well as immune response and blood pressure regulation was quantified by real-time RT-PCR in different tissues and PBMC. Total RNA was isolated with the peqGOLD TriFast™ according to the kit's manual. The concentration of total RNA in the sample was determined using a NanoDrop Spectrophotometer (Thermo Fisher Scientific Inc.) and the integrity of the RNA was confirmed by agarose gel electrophoresis. The synthesis of cDNA was performed using M-MLV Reverse Transcriptase (Promega, Madison, WI, USA). Subsequently, real-time RT-PCR was carried out as described elsewhere [20] using the Rotorgene 6,000 system and the Rotorgene software version 1.7 (Corbett Research, Mortlake, Australia). The relative mRNA abundance was calculated by the method of Pfaffl [21] using two appropriate reference genes. The primers used for the analyses are summarized in Table 1.

2.6. Analysis of minerals, lipids, nitrite, and folate

The concentrations of calcium, phosphorus, triglycerides and cholesterol were quantified in the plasma of pigs with test kits from DiaSys (Diagnostic Systems GmbH, Holzheim, Germany). The concentration of cholesterol in bile and liver was determined with the same test kit from DiaSys. Prior to quantification, lipid extracts of bile and liver were prepared as previously described [15,22]. The concentration of nitrite in the plasma, used as a biomarker of nitric oxide production, was determined by a colorimetric assay from Sigma-Aldrich. All procedures were in accordance with the manufacturer's protocols. The intra-day precisions were as follows: calcium, 4.4%; phosphorus, 13%; triglycerides, 10%; cholesterol, 5.6%; nitrite, 7.9%. The quantification of folate species in plasma was conducted by Bevital (Bergen, Norway) using LC-MS.

2.7. Analysis of the microbiome

The microbiome composition of feces and duodenal chyme was analyzed. To this end, total DNA was extracted using the Zymo-BIOMICS DNA Miniprep Kit (Zymo Research, Freiburg, Germany) and as described in details elsewhere [23]. 16S sequencing was performed by LGC Genomics (Berlin, Germany) using the Klindworth V3-V4 primers [24] and an Illumina MiSeq sequencer.

LotuS 1.62 [25] was employed to process the 16S rRNA amplicons from both stool and chyme microbiomes, utilizing the SILVA [26], Greengenes [27], and HITdb databases, thereby generating tables depicting microbiome abundance at all taxonomic levels. Sub-

Table 1
Primer of target and reference genes used for the analysis of mRNA abundance in tissues and PBMC of pigs.

Gene	Obtained from	Product size [bp]
<i>ABCG5</i>	Eurofins Genomics	282
<i>ACTB</i> ¹	Eurofins Genomics	204
<i>CD36</i>	Eurofins Genomics	103
<i>CYP2D25</i>	Sigma-Aldrich ²	104
<i>CYP27A1</i>	Sigma-Aldrich ²	107
<i>CYP27B1</i>	Sigma-Aldrich ²	170
<i>EDN1</i>	Sigma-Aldrich ²	145
<i>EDNRA</i>	Sigma-Aldrich ²	149
<i>EDNRB</i>	Sigma-Aldrich ²	189
<i>GAPDH</i> ¹	Eurofins Genomics	446
<i>IL6</i>	Sigma-Aldrich ²	125
<i>IL10</i>	Eurofins Genomics	446
<i>NOS2</i>	Sigma-Aldrich ²	191
<i>NOS3</i>	Sigma-Aldrich ²	174
<i>NPC1L1</i>	Eurofins Genomics	201
<i>RPS9</i> ¹	Eurofins Genomics	327
<i>SCARB1</i>	Eurofins Genomics	171
<i>TNF</i>	Sigma-Aldrich ²	144

Abbreviations: *ABCG5*, ATP binding cassette subfamily G member 5; *ACTB*, actin beta; *CD36*, CD36 molecule; *CYP2D25*, cytochrome P450 family 2 subfamily D member 25; *CYP27A1*, cytochrome P450 family 27 subfamily A member 1; *CYP27B1*, cytochrome P450 family 27 subfamily B member 1; *EDN1*, endothelin 1; *EDNRA*, endothelin receptor type A; *EDNRB*, endothelin receptor type B; *GAPDH*, glyceraldehyde-3-phosphate dehydrogenase; *IL6*, interleukin 6; *IL10*, interleukin 10; *NOS2*, nitric oxide synthase 2; *NOS3*, nitric oxide synthase 3; *NPC1L1*, NPC1 like intracellular cholesterol transporter 1; *RPS9*, ribosomal protein S9; *SCARB1*, scavenger receptor class B member 1; *TNF*, tumor necrosis factor.

¹ Reference gene.

² Pre-designed primer pair.

sequently, the abundance tables underwent rarefaction using RTK [28], and alpha diversities were calculated by the same tool.

2.8. Analysis of endothelin-1 in the renal artery

The concentration of endothelin-1 was determined in tissue homogenates of the porcine renal artery, using a commercial enzyme-linked immunosorbent assay (Cusabio, Texas, USA). The preparation of tissue homogenates and the assay procedure were following the manufacturer's protocol. The protein concentration of the tissue homogenates was assayed by the Bradford method [29].

2.9. Analysis of cytokine concentration

The concentrations of tumor necrosis factor alpha (TNF-alpha), interleukin 6 (IL6) and interleukin 10 (IL10) were determined in the supernatant of the PBMC after stimulation by use of enzyme-linked immunoassays (Bio-Techne GmbH, Wiesbaden, Germany). All procedures were in accordance with the manufacturer's protocol. Intra-day precisions were 2.4% for TNF-alpha, 4.2% for IL6, and 3.9% for IL10. The determined cytokine concentrations were related to the protein amount of the corresponding cells.

2.10. Analysis of bile acids and short-chain fatty acids (SCFA)

The concentration of bile acids was determined in bile and freeze-dried feces samples by MS-Omics (Vedbaek, Denmark) us-

ing LC-MS [30]. The analysis of SCFA in fecal water samples was carried out with gas chromatography-mass spectrometry by MS-Omics. For the fecal water extraction, feces samples were mixed with phosphate-buffered saline (1/3, w/v), vortexed and centrifuged at 16,000 g for 30 min. The supernatant was filtered through 0.2 µm centrifuge filters at 15,000 g for 2 min. The fecal water was stored at -20°C and shipped to MS-Omics on dry ice.

2.11. Metabolomic analysis in plasma

Metabolomic analysis of plasma samples was carried out by MS-Omics (Vedbaek) using a Thermo Scientific Vanquish LC coupled to Thermo Q Exactive HF MS. An electrospray ionization interface was used as ionization source. Analysis was performed in negative and positive ionization mode. Peak areas were extracted using Compound Discoverer 2.0 (Thermo Scientific). Identification of compounds was performed at four annotation levels; level 1: identification by retention times (compared against in-house authentic standards), accurate mass (with an accepted deviation of 3 ppm), and MS/MS spectra; level 2a: identification by retention times (compared against in-house authentic standards), accurate mass (with an accepted deviation of 3 ppm); level 2b: identification by accurate mass (with an accepted deviation of 3 ppm), and MS/MS spectra; level 3: identification by accurate mass alone (with an accepted deviation of 3 ppm).

2.12. Statistical analyses

Data from plasma and feces samples with repeated measurements were analyzed using the Statistical Analysis System (SAS Institute Inc., Cary, NC, USA). Prior to the *mixed-model procedure* (*PROC MIX*), the data were tested for normal distribution or log-normal distribution. The treatment (group), time of intervention and their interaction (group x time) were considered as fixed effects and the animals were considered random. For significant interaction effects, the Tukey-Kramer post-hoc was conducted. For all data from tissue samples (no repeated measurements) and plasma 3-epi-25(OH)D₃, the two groups were compared by the non-parametric Mann-Whitney *U* test (GraphPad Prism version 9; GraphPad Software, Boston, MA, USA). Values that were below the LOQ were included as LOQ/2 in the statistical analyses, if more than 25% of the data within one group were above the LOQ. Otherwise, no statistical analysis was conducted for this parameter.

For statistical analysis of microbiome, Wilcoxon rank-sum and Spearman correlation tests were carried out using the base R environment. Multivariate analysis, employing the Adonis test based on distance matrices, was conducted using the Vegan [31] R package. Univariate tests were executed utilizing the glimmTMB [32] package, where the abundance of each microbe was fitted through the interaction term of plasma vitamin D concentration and treatment, with the subject included as a random factor. A significance threshold for q-values, adjusted via the Benjamini-Hochberg method, was set at 0.05. Heatmaps were generated using the gplots package. Finally, the repeated measurement correlation test was performed with the Rmcorr [33] package.

For the statistical analysis of metabolome data, the log₂ values of both groups were subjected to a student's t-test with Benjamini-Hochberg correction.

Significant differences were assumed for $P < .05$. If not otherwise stated, data are presented as median and interquartile ranges.




Fig. 1. Plasma concentration of 25-hydroxyvitamin D₃ (25(OH)D₃) in pigs with a daily UV exposure of 0 min (dietary vitamin D), 1 min, 5 min, 10 min, 20 min or 30 min. Pigs exposed to UV light did not receive oral vitamin D₃. Data are reported as mean values \pm standard deviation.

3. Results

3.1. Determining the UV light exposure time in the pre-study

The pre-study aimed to assess the daily UV light exposure time which leads to similar plasma 25(OH)D levels as an oral administration of 20 μ g/d vitamin D₃ (Fig. 1). At baseline, the 25(OH)D₃ plasma concentration (given as mean \pm standard deviation) of all pigs was 10.6 \pm 2.3 nmol/l. The highest increase of plasma 25(OH)D₃ was observed within the first week of treatment in all groups (Fig. 1). After week 1, orally supplemented pigs showed plasma 25(OH)D₃ concentrations that corresponded to those of groups exposed to UV light for 20 min/d and 30 min/d, respectively. After 2 weeks of treatment, orally supplemented pigs had plasma levels of 25(OH)D₃ similar to the groups exposed to UV light for 10 min/d and 20 min/d, respectively (Fig. 1). To determine what daily UV light exposure time over the entire treatment period of 4 weeks represents best the 25(OH)D₃ concentrations obtained by dietary vitamin D supply, a model comparison analysis was conducted. Based on the Akaike information criterion values and F-test, a linear association between UV light exposure time and plasma 25(OH)D₃ was found in week 1, 2, 3 and 4 of the experiment. By using the weekly assessed regression equations, the daily UV light exposure time that resulted in plasma 25(OH)D₃ levels similar to those of daily supplemented pigs was calculated. Based on this calculation, a UV light exposure time of 19 min/d was considered to be sufficient to produce 25(OH)D₃ plasma levels similar to those analyzed in the group fed 20 μ g/d vitamin D₃ (supplementary Fig. 2). This UV light exposure time was used for the main study.

3.2. Main study

3.2.1. Vitamin D metabolism in pigs treated with oral vitamin D₃ or UV light exposure

All pigs were vitamin D deficient at week 0. The plasma concentration of 25(OH)D₃ (measured as the sum of non-epimerized 25(OH)D₃ and 3-epi-25(OH)D₃) increased during the study in both groups ($P < .001$, Fig. 2A) with the highest rise within the first week of intervention. As intended, plasma 25(OH)D₃ concentrations were not different between both groups throughout the experimental period (Fig. 2B). Interestingly, the concentration of 3-




Fig. 2. Plasma vitamin D metabolite concentration in pigs over four weeks of intervention. (a) 25-hydroxyvitamin D₃ (25(OH)D₃), reflecting the sum of 3-epimerized and non-epimerized 25(OH)D₃. (b) 3-epi-25-hydroxyvitamin D₃ (3-epi-25(OH)D₃). (c) Vitamin D₃. (d) 24,25-dihydroxyvitamin D₃ (24,25(OH)₂D₃). Data are reported as median and interquartile range ($n=8$). * Significant difference between the groups ($* P < .05$, $** P < .01$, $*** P < .001$, (b) Mann-Whitey U test, (c) Tukey-Kramer test). # Median values below the limit of quantification (3-epi-25(OH)D₃: 2.5 nmol/l, vitamin D₃: 0.26 nmol/l, 24,25(OH)₂D₃: 1.44 nmol/l).




Fig. 3. Plasma concentration of (a) 1,25-dihydroxyvitamin D₃ (1,25(OH)₂D₃) and (b) parathyroid hormone (PTH) in pigs that were supplemented with 20 µg/d vitamin D₃ or treated with UV light for 19 min/d during four weeks of intervention. Data are reported as median and interquartile range ($n=8$). Not significant (ns).

epi-25(OH)D₃ increased in response to UV light exposure, but remained below the LOQ (2.5 nmol/l) in the supplemented group during the whole study (Fig. 2B). The concentration of plasma vitamin D₃ was below the LOQ (0.26 nmol/l) at baseline and strongly increased within the first week of vitamin D supply in both groups (Fig. 2C). Interestingly, the treatment affected the plasma concentration of vitamin D₃, whereby the UV light-exposed group had higher levels than the supplemented group (Fig. 2B, $P < .05$). The most marked difference between the two groups was found at the end of the study. The concentration of 3-epi-25(OH)D₃ increased in

response to UV light exposure, but remained below the LOQ in the supplemented group during the whole study. However, the plasma concentrations of 24,25(OH)₂D₃ (Fig. 2D and 1) and 25(OH)₂D₃ (Fig. 3A) increased during the experimental period ($P < .001$), but were not affected by the treatment. The plasma concentration of parathyroid hormone (PTH) did not change with the treatment and time (Fig. 3B). Lumisterol₃, a photoproduct of the UV irradiation, was not quantifiable in the plasma of pigs (LOQ: 4 nmol/l).

Vitamin D metabolites were then quantified in the skin of the dorsal back, where the highest UV light exposure and vitamin D synthesis was expected. Remarkably, the quantities of vitamin D₃ and lumisterol₃ were nearly 50 times higher in the UV light-exposed group than in the supplemented group ($P < .001$, Fig. 4). Additionally, higher quantities of vitamin D₃ were also seen in kidney ($P = .010$), subcutaneous fat ($P < .001$), and bile ($P < .05$) of UV light-exposed that in the supplemented pigs (Fig. 5). However, analysis of fecal vitamin D₃ revealed lower values in the UV light-exposed group compared to the supplemented group ($P < .001$, Fig. 5). There were no significant differences in the quantities of vitamin D₃ in liver, visceral fat, trapezius muscle and intestinal mucosa (Fig. 5). The concentration of 25(OH)D₃ in liver (dietary vitamin D: 2.57 ng/g [1.99-4.35 ng/g], UV light exposure: 2.54 ng/g [2.11-2.71 ng/g]) and kidney (dietary vitamin D: 2.98 ng/g [1.43-3.70 ng/g], UV light exposure: 2.63 ng/g [2.19-4.53 ng/g]) were not different. The tissue concentrations of 25(OH)D₃ were below the LOQ (7 ng/g) in subcutaneous fat, visceral fat, muscle, intestinal mucosa, bile and faeces.

Next, the mRNA abundance of cytochrome P450 (CYP) genes that encode the porcine-specific vitamin D hydroxylase CYP2D25, the mitochondrial CYP27A1 and the 1 α -hydroxylase CYP27B1 was assessed (Fig 6). Dorsal skin of UV light-exposed pigs showed a higher mRNA abundance of CYP2D25 ($P < .05$) and a trend toward higher mRNA abundance of CYP27A1 ($P < .1$) than that of supplemented pigs, whereas in liver, intestinal mucosa and kidney the mRNA abundance of the vitamin D hydroxylases did not differ.

For intestinal and hepatic lipid transporter, the only difference in mRNA abundance was observed for the hepatic sterol transporter NPC1 like intracellular cholesterol transporter 1 (NPC1L1) which was higher in the UV light-exposed group than in the supplemented group ($P < .05$, Table 2). In contrast, intestinal NPC1L1, and ATP binding cassette subfamily G member 5 (ABCG5), CD36 molecule (CD36) and scavenger receptor class B member 1 (SCARB1) in the intestinal mucosa and liver did not differ between the two groups (Table 2).




Fig. 4. Concentration of (a) vitamin D₃ and (b) lumisterol₃ in the skin of pigs that were supplemented with 20 µg/d vitamin D₃ or treated with UV light for 19 min/d after four weeks of intervention. ($n=8$) *** Significant difference between the groups (Mann-Whitney U test, $P < .001$).




Fig. 5. Concentration of vitamin D₃ in samples of pigs that were supplemented with 20 µg/d vitamin D₃ or treated with UV light for 19 min/d after four weeks of intervention. (a) liver, (b) kidney, (c) subcutaneous fat, (d) visceral fat, (e) trapezius muscle, (f) intestinal mucosa, (g) bile, (h) feces. (n=8) * Significant difference between the groups (Mann-Whitney U test, * P < .05, ** P < .01, *** P < .001).




Fig. 6. Relative mRNA abundance of vitamin D hydroxylases in tissues of pigs that were supplemented with 20 $\mu\text{g}/\text{d}$ vitamin D_3 or treated with UV light for 19 min/d after four weeks of intervention. (a) Cytochrome P450 family 27 subfamily A member 1 (*CYP27A1*), (b) cytochrome P450 family 2 subfamily D member 25 (*CYP2D25*), (c) cytochrome P450 family 27 subfamily B member 1 (*CYP27B1*). Reference genes: glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) and ribosomal protein S9 (*RPS9*), ($n=8$). * Significant difference between the groups (Mann-Whitney *U* test, $P < .05$). # Trend toward significant difference between the groups (Mann-Whitney *U* test, $P < .1$).

Table 2

Relative mRNA abundance of transporters expressed in small intestine mucosa and liver of pigs after four weeks of intervention.

	Dietary vitamin D_3		UV exposure		<i>P</i> value
	Median	IQR	Median	IQR	
Intestinal mucosa					
<i>ABCG5</i>	0.84	0.60–1.22	0.78	0.64–0.87	Ns
<i>CD36</i>	0.93	0.78–1.20	1.63	0.87–2.51	< .1
<i>NPC1L1</i>	0.87	0.49–1.25	1.54	0.58–2.07	Ns
<i>SCARB1</i>	0.94	0.45–1.53	2.05	0.51–3.05	Ns
Liver					
<i>ABCG5</i>	1.06	0.63–1.24	1.16	0.82–1.25	Ns
<i>NPC1L1</i>	1.07	0.46–1.44	2.01	1.36–2.66	< .05
<i>SCARB1</i>	1.00	0.74–1.30	1.44	0.82–1.77	Ns

Reference genes: glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) and ribosomal protein S9 (*RPS9*).

Abbreviations: ABCG5, ATP binding cassette subfamily G member 5; CD36, CD36 molecule; IQR, interquartile range; Ns, not significant; NPC1L1, NPC1 like intracellular cholesterol transporter 1; SCARB1, scavenger receptor class B member 1. Groups were compared by the Mann-Whitney *U* test ($n=8$).

3.2.2. Concentration of minerals, lipids, and folate

The plasma concentrations of calcium and phosphorus, cholesterol, triglycerides (Table 3) and folate (Supplementary Table 2) did not differ between the two groups. Interestingly, the hepatic cholesterol level was moderately lower in UV light-exposed pigs than in supplemented pigs ($P < .1$), although biliary cholesterol was not different between the two groups (Table 3).

3.2.3. Microbiome composition, bile acids and short-chain fatty acids

In the examination of the stool microbiome, no statistically significant differences (Wilcoxon rank-sum test $P \geq .05$) in alpha diversities (including richness, evenness, Shannon diversity, and Chao1 index) were discerned neither among various treatments nor across distinct time points. Employing multivariate analysis via the Adonis test, with the time variable representing variations in vitamin D concentration and pig growth, it was found that the time variable significantly influenced microbial composition ($P < .05$), while the treatment factor did not exhibit any significant impact. Although significant alterations in microbial populations were observed across different time points, indicative of fluctuations in vitamin D levels, our investigation did not yield substantive evidence linking the sources of vitamin D to discernible effects on the stool microbiome, whether through multivariate or univariate assessments. On the other hand, when scrutinizing the chyme microbiome, no statistically significant variations in alpha diversities (encompassing richness, evenness, Shannon diversity, and the Chao1 index) were detected among treatments. Employing a multivariate analysis (Adonis test) at the species level, the treatment factor did not exert a discernible influence on microbial composition. Likewise, no evidence was found to suggest that the sources of vitamin D significantly impacted the chyme microbiome, whether through multivariate or univariate analytical approaches.

Correlation analyses conducted between the stool and chyme microbiomes have unveiled a multitude of statistically significant correlations existing at the species, genus, and family taxonomic levels (Supplementary Fig. 3–5). This evidence underscores a robust and close association between the stool and chyme microbiome. Furthermore, when analyzing the correlation between the stool microbiome and plasma metabolite concentrations, a positive correlation was identified between the abundance of *Mailhella massiliensis* and creatine (Supplementary Fig. 6).

Table 3
Concentrations of minerals and lipids in pigs during four weeks of intervention.

	Dietary vitamin D ₃		UV exposure		P values		
	Median	IQR	Median	IQR	Group	Time	Group x time
Plasma calcium (mmol/l)							
Week 0	2.55	2.47–2.67	2.57	2.49–2.82	Ns	Ns	Ns
Week 2	2.66	2.39–2.75	2.76	2.65–2.82			
Week 4	2.54	2.34–2.73	2.67	2.45–2.85			
Plasma phosphorus (mmol/l)							
Week 0	2.76	2.46–2.96	2.96	2.52–3.45	Ns	<i>P</i> < .001	Ns
Week 2	3.24	3.00–3.60	3.32	3.13–3.55			
Week 4	4.03	3.84–4.31	4.20	3.97–4.49			
Plasma cholesterol (mmol/l)							
Week 0	2.17	2.08–2.29	2.22	1.91–2.38	Ns	<i>P</i> < .01	Ns
Week 2	2.54	2.26–2.88	2.75	2.36–3.40			
Week 4	2.77	2.63–2.88	2.57	2.45–3.02			
Plasma triglycerides (mmol/l)							
Week 0	0.42	0.35–0.56	0.42	0.29–0.57	Ns	<i>P</i> < .01	Ns
Week 2	0.51	0.42–0.61	0.50	0.43–0.58			
Week 4	0.57	0.34–0.90	0.56	0.51–0.93			
Hepatic cholesterol (mg/g)							
Week 4	2.56	2.28–2.73	2.31	1.79–2.41	Ns ¹		
Biliary cholesterol (mg/g)							
Week 4	3.96	3.12–4.46	3.32	2.99–3.63	Ns ¹		

Abbreviations: IQR, Interquartile range; Ns, not significant. Plasma concentrations with repeated measures were statistically analysed by the *mixed-model procedure*.

¹ Tissue concentrations without repeated measures were analysed by the Mann-Whitney *U* test (*n*=8).

Additionally, bile acids in feces and bile, as well as short-chain fatty acids in feces did not differ significantly between the two groups of pigs (Supplementary Table 3–5).

3.2.4. Regulators of blood pressure

The concentration of plasma nitrite, a marker of nitric oxide production, decreased during the study period (*P* < .001), whereby the decrease was less pronounced in UV light-exposed pigs (Fig. 7), indicating a vasodilative response to UV radiation. However, the mRNA abundance of nitric oxide synthase 2 (*NOS2*) and nitric oxide synthase 3 (*NOS3*) in the renal artery, skin and subcutaneous fat was not differently affected by the treatments (Fig. 7). To investigate factors involved in vasoconstriction, the relative mRNA abundance of endothelin-1 (*EDN1*), endothelin receptor type A (*EDNRA*) and endothelin receptor type B (*EDNRB*) was analyzed in the renal artery. A trend toward lower *EDN1* mRNA abundance was observed in pigs exposed to UV light compared to vitamin D₃-supplemented pigs (*P* < .1), whereas the protein expression of endothelin-1 and the relative mRNA abundance of *EDNRA* and *EDNRB* did not differ between the two groups (Fig. 8).

3.2.5. Cytokine expression in PBMC

PBMC isolated from both groups and stimulated with LPS showed a lower relative mRNA abundance and protein release of TNF-alpha and IL10 in pigs exposed to UV light compared to pigs supplemented with vitamin D₃ (*P* < .05, Table 4). No differences between the groups were observed for the mRNA abundance and the concentration of released IL6 (Table 4).

3.2.6. Plasma metabolome

An untargeted metabolome analysis was conducted in plasma samples of UV light-exposed pigs and pigs with dietary vitamin D supplementation and 153 compounds with an annotation level 1, 2a or 2b were identified (Supplementary Table 6). Four compounds were increased in UV light-exposed pigs with a high significant

Table 4

Cytokine expression in stimulated peripheral blood mononuclear cells (PBMC) of pigs after four weeks of intervention.

	Dietary vitamin D ₃		UV exposure		P value
	Median	IQR	Median	IQR	
Relative mRNA abundance ¹					
<i>IL6</i>	1.05	0.61–1.38	0.49	0.13–1.26	Ns
<i>IL10</i>	1.03	0.80–1.19	0.46	0.38–0.79	< .05
<i>TNF</i>	0.99	0.91–1.10	0.42	0.30–0.67	< .05
Cytokine concentration [pg/mg protein]					
<i>IL6</i>	283	184–482	226	81–355	Ns
<i>IL10</i>	140	121–404	83	65–100	< .05
<i>TNF-alpha</i>	3236	2261–3892	863	457–1276	< .05

Abbreviations: IL6, Interleukin 6; IL10, interleukin 10; IQR, interquartile range; ns, not significant; TNF, tumor necrosis factor. Groups were compared by the Mann-Whitney *U* test (*n*=5).

¹ Reference genes: actin beta (*ACTB*) and glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*).

P value < 0.001: hexadecanamide, trans-anethole, 1-dodecyl-2-pyrrolidinone and (2E,4E)-N-(2-methylpropyl)deca-2,4-dienamide. Two compounds were increased in the UV light-exposed group with a significant *P* value < 0.05: 4-methylumbelliferone hydrate and monobutyl phthalate. However, none of these metabolites could be assigned a physiological function.

4. Discussion

The current study compared the effects of UV light exposure versus vitamin D₃ supplementation on vitamin D metabolites, and factors indicating health or disease risk. The most marked findings of this study were differences in the profile and tissue distribution of vitamin D₃ metabolites, factors associated with blood pressure




Fig. 7. Plasma nitrite concentration and relative mRNA abundance of nitric oxide synthases in pigs that were supplemented with 20 $\mu\text{g/d}$ vitamin D_3 or treated with UV light for 19 min/d. (a) Plasma nitrite reflects the concentration of nitric oxide. Data are reported as median and interquartile range. (b) Inducible nitric oxide synthase (*NOS2*), (c) endothelial nitric oxide synthase (*NOS3*). Data of (b) and (c) were measured after four weeks of treatment. ($n=8$). ** Significant difference between the groups ($P < .01$, Tukey-Kramer test). Not significant (ns).

regulation, and the immune response. Many other factors with relevance for health, including plasma lipids, gut microbiome, short-chain fatty acids, folic acid status, and other metabolites, did not show differences between the two treatments. An important finding of this study was that UV light-exposed pigs had more vitamin D_3 in plasma, skin, kidney, and subcutaneous fat tissue than supplemented pigs, although the plasma 25(OH) D_3 levels were similar in both groups. We assume that the higher vitamin D_3 stores in skin, and subcutaneous fat, which constitute a large part of the whole body fat [34,35], of UV light-exposed pigs are physiologically important as they may contribute to counteract declining plasma 25(OH) D_3 levels in times of insufficient vitamin D_3 synthesis. The importance of vitamin D stores in adipose tissue has already been described in a human study that found high vitamin D stores associated with a smaller decline in 25(OH) D during winter periods [36]. It is important to note that the increase of cutaneous vitamin D_3 in UV light-exposed pigs was accompanied by a higher mRNA expression of *CYP2D25*, a vital vitamin D_3 hydroxylase. An upregulated cutaneous expression of vitamin D hydroxylases after UV light treatment has also been found by others [37,38], and is probably a response to higher substrate availability. However, recent research has uncovered alternative pathways of vitamin D_3 metabolism, particularly involving the enzyme *CYP11A1* [39], which has been upregulated in human skin samples

exposed to UV light [40]. It must therefore be assumed that UV light-exposed pigs in the current study had produced some alternative vitamin D_3 metabolites that exhibit biological activities that differ from the classical vitamin D metabolites. Interestingly, UV light primarily increases vitamin D_3 in the subcutaneous fat rather than in visceral fat, which is probably because vitamin D is stored mainly close to the site of production. The current data might also confirm previous findings demonstrating that circulating 25(OH) D_3 does not necessarily reflect the tissue stores of vitamin D_3 in the body [41].

When UV radiation penetrates the skin, 7-dehydrocholesterol is converted to pre-vitamin D_3 , which undergoes a temperature-dependent isomerization to vitamin D_3 . Pre-vitamin D_3 can also isomerize to photochemical by-products of UV light exposure such as lumisterol $_3$ [42]. Remarkably high quantities of lumisterol $_3$ is what we found in the skin of UV light-exposed pigs in contrast to supplemented pigs. We assume that this finding is also very important with respect to the fact that lumisterol $_3$ has been identified as a molecule with beneficial effects in keratinocytes such as proliferation and antioxidative responses [43].

In addition, we found another quite important difference in plasma 3-epi-25(OH) D_3 , that largely increased in UV light-exposed pigs over the time. To date, little is known about the precise function and health implication of C3-epimerized vitamin D forms, and




Fig. 8. Expression of endothelin-1 and endothelin receptors in the renal artery of pigs that were supplemented with 20 $\mu\text{g}/\text{d}$ vitamin D_3 or treated with UV light for 19 min/d after four weeks of intervention. (a) relative mRNA abundance of endothelin-1 (*EDN1*) and endothelin receptor A (*EDNRA*) and B (*EDNRB*), (b) Protein expression of endothelin-1. ($n=8$) # Trend toward significant difference between the groups (Mann-Whitney *U* test, $P = .1$).

also the gene encoding for the epimerization enzyme has not yet been identified [44]. It is suggested that 3-epi-1,25(OH) $_2\text{D}_3$, which has been shown to be synthesized in the endoplasmic reticulum of hepatocytes, bone cells and keratinocytes [45], can stimulate the expression of vitamin D regulated genes by binding to and activating the vitamin D receptor, but less effective than the non-epimeric calcitriol [45]. Data on external factors that stimulate 3-epi-25(OH) D_3 formation are scarce. There are only two studies that identified higher epimeric vitamin D levels in oral supplemented compared to UV light-exposed mice [46], whereas in another study an increase of 3-epi-25(OH) D_3 was observed in mice exposed to LED lamps emitting light with wavelengths in the range of 275–900 nm compared to mice kept in the dark [47]. The current study has contributed to identify UV light exposure as an environmental factor that stimulates the production of 3-epi-25(OH) D_3 .

One of the most apparent differences between dietary vitamin D supply and endogenous synthesis is the way vitamin D enters the body. The uptake of vitamin D from food sources is a complex process that involves both passive diffusion and active transport mechanisms that are facilitated by intestinal lipid transporters, such as NPC1L1, located at the apical site of the enterocyte [16,18]. The expression of these lipid transporters was measured, in order to test if the expression is modified by oral vitamin D_3 intake, but observed no differences between the supplemented and the UV light-exposed pigs. However, NPC1L1 is a sterol transporter that is not only located in the gut, but also in the liver, where it mediates the hepatic re-uptake of cholesterol to prevent excessive biliary loss [48]. Telford et al. [49] observed an increase of hepatic NPC1L1 in response to tissue cholesterol depletion in pigs. Interestingly, our study found moderately lower hepatic cholesterol levels and a higher mRNA abundance of hepatic *NPC1L1* in UV light-exposed pigs than in supplemented pigs. We hypothesize that the upregulation of hepatic *NPC1L1* was caused by the lower levels of liver cholesterol and that UV light-exposed pigs might benefit from an enhanced biliary re-uptake of vitamin D_3 as a secondary effect.

The role of vitamin D and UV light in blood pressure regulation is still a topic of scientific debate [10,50]. Vitamin D has been shown to affect the production of nitric oxide, the most potent endothelial vasodilator, and is therefore notably involved in the modulation of vascular tone and resistance [51]. Due to its very short lifespan, measuring nitric oxide in biological samples is challenging, and nitrite has become an essential surrogate biomarker for nitric oxide production [52,53]. The higher plasma levels of nitrite observed in the UV light-exposed pigs are probably indicative of

a higher nitric oxide release from the skin into the circulation, although the gene expression of *NOS2* and *NOS3* did not change by the UV light exposure. This phenomenon has already been described in the context of UV light exposure [54,55]. An antagonist of nitric oxide is endothelin-1, the most potent vasoconstrictor that mediates its effects through the activation of endothelin receptors on vascular smooth muscle cells [56]. Although it is suggested that both vitamin D and UV light can stimulate endothelin expression and/or response *in vitro* and *in vivo* [57–60], we found a moderate down-regulation of endothelin-1 in UV light-exposed pigs in comparison to supplemented pigs. On the other hand, nitric oxide has been described to be capable of decreasing circulating endothelin-1 levels [61–63], which may explain the reduced expression of *EDN1* in the renal artery of UV light-exposed pigs. However, a limitation of the study is that blood pressure was not directly measured in pigs due to their high stress level induced by restraining.

Vitamin D and UV light are known to influence immune response. In the present study, UV light exposure was associated with lower expression and release of the proinflammatory TNF-alpha and the anti-inflammatory IL10 in stimulated PBMC in comparison to the vitamin D_3 supplementation. This finding fits well with the commonly known immune suppressive impact of UV radiation [9,64], and data from humans exposed to artificial or natural UV light whose isolated and *ex vivo* stimulated PBMC showed a reduced capacity to secrete the cytokines TNF-alpha and IL10 [65,66]. In the present study, the reduced release of TNF-alpha and IL10 might indicate an overall suppression or attenuation of the innate immune response, which might be beneficial in controlling inflammation but could also impair the ability to fight infections effectively.

When comparing both sources of vitamin D supply, one should keep in mind that UV light exposure is associated with numerous systemic effects that arise independently of vitamin D [40,67,68]. To better point out health outcomes which are derived from vitamin D and are not the consequence of UV light exposure, it would be helpful to identify sensitive and reliable biomarkers of UV light exposure. Unfortunately, the untargeted plasma metabolome analysis in the current study did not provide promising metabolites.

To summarize, the current study demonstrates some health relevant differences between oral vitamin D_3 supplementation and UV light exposure. UV light exposure in contrast to vitamin D_3 supplementation resulted in higher levels of stored vitamin D_3 in skin and subcutaneous fat, which might counteract a decline in vitamin

D levels in times of insufficient vitamin D synthesis. UV light exposure further contributed to an increase in the production of the physiologically active lumisterol₃ and 3-epi-25(OH)D₃. And finally, it was shown that UV light exposure in contrast to supplemented vitamin D₃ can modify regulators of blood pressure and the immune response. These differences are important with respect to recommendations concerning the best option to improve people's vitamin D status and may also explain discrepancies in the results from observational studies and intervention studies.

Declaration of competing interest

The authors declare that there are no conflicts of interests.

CRedit authorship contribution statement

Julia Kühn: Writing – original draft, Visualization, Project administration, Methodology, Investigation, Formal analysis, Conceptualization. **Corinna Brandsch:** Writing – review & editing, Visualization, Investigation, Formal analysis. **Anja C. Bailer:** Writing – review & editing, Investigation, Conceptualization. **Mikis Kiourtzidis:** Writing – review & editing, Investigation. **Frank Hirche:** Writing – review & editing, Investigation. **Chia-Yu Chen:** Writing – review & editing, Writing – original draft, Visualization, Investigation, Formal analysis. **Lajos Markó:** Writing – review & editing, Conceptualization. **Theda U.P. Bartolomaeus:** Investigation. **Ulrike Löber:** Investigation. **Samira Michel:** Writing – review & editing, Investigation. **Monika Wensch-Dorendorf:** Writing – original draft, Formal analysis. **Sofia K. Forslund-Startceva:** Writing – review & editing, Supervision, Resources. **Gabriele I. Stangl:** Writing – original draft, Supervision, Project administration, Funding acquisition, Conceptualization.

Data availability statement

The data that support the findings of this study are available on reasonable request from the corresponding author. Bacterial 16S rRNA reads were deposited in the NCBI Sequence Read Archive (SRA) database (BioProject ID: PRJNA1062174).

Acknowledgements

This work was funded by the Federal Ministry of Education and Research (BMBF, grant no. 1EA1808C). We thank H. Giese and C. Leibelt for technical assistance. We further thank Dr. G. Woitow, K. Müller, S. Reich and E. Dietrich for their expertise and assistance with the care and handling of the pigs.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.jnutbio.2024.109746](https://doi.org/10.1016/j.jnutbio.2024.109746).

References

- Forman JP, Giovannucci E, Holmes MD, Bischoff-Ferrari HA, Tworoger SS, Willett WC, et al. Plasma 25-hydroxyvitamin D levels and risk of incident hypertension. *Hypertension* 2007;49(5):1063–9. doi:10.1161/HYPERTENSIONAHA.107.087288.
- Tamez H, Thadhani RI. Vitamin D and hypertension: an update and review. *Curr Opin Nephrol Hypertens* 2012;21(5):492–9. doi:10.1097/MNH.0b013e3283557bf0.
- Parikh S, Guo D-H, Pollock NK, Petty K, Bhagatwala J, Gutin B, et al. Circulating 25-hydroxyvitamin D concentrations are correlated with cardiometabolic risk among American black and white adolescents living in a year-round sunny climate. *Diabet Care* 2012;35(5):1133–8. doi:10.2337/dc11-1944.
- Seibert E, Lehmann U, Riedel A, Ulrich C, Hirche F, Brandsch C, et al. Vitamin D3 supplementation does not modify cardiovascular risk profile of adults with inadequate vitamin D status. *Eur J Nutr* 2017;56(2):621–34. doi:10.1007/s00394-015-1106-8.
- Qi D, Nie X, Cai J. The effect of vitamin D supplementation on hypertension in non-CKD populations: a systemic review and meta-analysis. *Int J Cardiol* 2017;227:177–86. doi:10.1016/j.ijcard.2016.11.040.
- Zhang D, Cheng C, Wang Y, Sun H, Yu S, Xue Y, et al. Effect of vitamin D on blood pressure and hypertension in the general population: an update meta-analysis of cohort studies and randomized controlled trials. *Prev Chronic Dis* 2020;17:E03. doi:10.5888/pcd17.190307.
- Webb AR, Pilbeam C, Hanafin N, Holick MF. An evaluation of the relative contributions of exposure to sunlight and of diet to the circulating concentrations of 25-hydroxyvitamin D in an elderly nursing home population in Boston. *Am J Clin Nutr* 1990;51(6):1075–81. doi:10.1093/ajcn/51.6.1075.
- Fraser DR. Physiological significance of vitamin D produced in skin compared with oral vitamin D. *J Nutr Sci* 2022;11:e13. doi:10.1017/jns.2022.11.
- Ullrich SE. Mechanisms underlying UV-induced immune suppression. *Mutat Res* 2005;571(1-2):185–205. doi:10.1016/j.mrfmmm.2004.06.059.
- Scragg R, Rahman J, Thornley S. Association of sun and UV exposure with blood pressure and cardiovascular disease: a systematic review. *J Steroid Biochem Mol Biol* 2019;187:68–75. doi:10.1016/j.jsbmb.2018.11.002.
- He JL, Scragg RK. Vitamin D, parathyroid hormone, and blood pressure in the National Health and Nutrition Examination Surveys. *Am J Hypertens* 2011;24(8):911–17. doi:10.1038/ajh.2011.73.
- Larsen T, Mose FH, Bech JN, Hansen AB, Pedersen EB. Effect of cholecalciferol supplementation during winter months in patients with hypertension: a randomized, placebo-controlled trial. *Am J Hypertens* 2012;25(11):215–22. doi:10.1038/ajh.2012.111.
- McMullan CJ, Borgi L, Curhan GC, Fisher N, Forman JP. The effect of vitamin D on renin-angiotensin system activation and blood pressure: a randomized control trial. *J Hypertens* 2017;35(4):822–9. doi:10.1097/HJH.0000000000001220.
- Gräber T, Kluge H, Granica S, Horn G, Brandsch C, Stangl GI. Studies on the health impact of agrimonia provera in piglets. *BMC Vet Res* 2014;10:210. doi:10.1186/s12917-014-0210-y.
- Kühn J, Hirche F, Geissler S, Stangl GI. Oral intake of 7-dehydrocholesterol increases vitamin D(3) concentrations in the liver and kidney. *J Steroid Biochem Mol Biol* 2016;164:199–204. doi:10.1016/j.jsbmb.2015.12.017.
- Kiourtzidis M, Kühn J, Schutkowski A, Baur AC, Hirche F, Stangl GI. Inhibition of niemann-pick C1-like protein 1 by ezetimibe reduces uptake of deuterium-labeled vitamin D in mice. *J Steroid Biochem Mol Biol* 2020;197:105504. doi:10.1016/j.jsbmb.2019.105504.
- Kotwan J, Kühn J, Baur AC, Stangl GI. Oral intake of lumisterol affects the metabolism of vitamin D. *Mol Nutr Food Res* 2021;65(14):e2001165. doi:10.1002/mnfr.202001165.
- Kiourtzidis M, Kühn J, Brandsch C, Stangl GI. Vitamin D status of mice deficient in scavenger receptor class B Type 1, cluster determinant 36 and ATP-binding cassette proteins G5/G8. *Nutrients* 2020;12(8):2169. doi:10.3390/nu12082169.
- Kühn J, Wassermann C, Ebschke S, Schutkowski A, Thamm K, Wensch-Dorendorf M, et al. Feasibility of artificial light regimes to increase the vitamin D content in indoor-laid eggs. *Poult Sci* 2019;98(10):5177–87. doi:10.3382/ps/pez234.
- Radtke J, Geissler S, Schutkowski A, Brandsch C, Kluge H, Duranti MM, et al. Lupin protein isolate versus casein modifies cholesterol excretion and mRNA expression of intestinal sterol transporters in a pig model. *Nutr Metab (Lond)* 2014;11(1):9. doi:10.1186/1743-7075-11-9.
- Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 2001;29(9):e45.
- Hara A, Radin NS. Lipid extraction of tissues with a low-toxicity solvent. *Anal Biochem* 1978;90(1):420–6.
- Bartolomaeus TUP, Birkner T, Bartolomaeus H, Löber U, Avery EG, Mähler A, et al. Quantifying technical confounders in microbiome studies. *Cardiovasc Res* 2021;117(3):863–75. doi:10.1093/cvr/cvaa128.
- Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, et al. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res* 2013;41(1):e1. doi:10.1093/nar/gks808.
- Hildebrand F, Tadeo R, Voigt AY, Bork P, Raes J. LotuS: an efficient and user-friendly OTU processing pipeline. *Microbiome* 2014;2(1):30. doi:10.1186/2049-2618-2-30.
- Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res* 2013;41(Database issue):D590–6. doi:10.1093/nar/gks1219.
- McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, Probst A, et al. An improved greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME J* 2012;6(3):610–18. doi:10.1038/ismej.2011.139.
- Saary P, Forslund K, Bork P, Hildebrand F. RTK: efficient rarefaction analysis of large datasets. *Bioinformatics* 2017;33(16):2594–5. doi:10.1093/bioinformatics/btx206.
- Kruger NJ. The Bradford method for protein quantitation. *Meth Mol Biol* 1994;32:9–15. doi:10.1385/0-89603-268-X:9.
- Kühn J, Brandsch C, Kiourtzidis M, Nier A, Bieler S, Matthäus B, et al. Microalgae-derived sterols do not reduce the bioavailability of oral vita-

- min D(3) in mice. *Int J Vitam Nutr Res* 2022;93(6):507–17. doi:10.1024/0300-9831/a000766.
- [31] Dixon P. VEGAN, a package of R functions for community ecology. *J Veg Sci* 2003;14(6):927–30. doi:10.1111/j.1654-1103.2003.tb02228.x.
- [32] Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, et al. Modeling zero-inflated count data with glmmTMB; 2017. Accessed September 6, 2024 <https://doi.org/10.1101/132753>.
- [33] Bakdash JZ, Marusich LR. Repeated measures correlation. *Front Psychol* 2017;8:456. doi:10.3389/fpsyg.2017.00456.
- [34] Kolstad K. Fat deposition and distribution measured by computer tomography in three genetic groups of pigs. *Livest Prod Sci* 2001;67(3):281–92. doi:10.1016/S0301-6226(00)00195-0.
- [35] Richmond RJ, Berg RT. Fat distribution in swine as influenced by liveweight, breed, sex and ration. *Can J Anim Sci* 1971;51(2):523–31. doi:10.4141/cjas71-068.
- [36] Martinaityte I, Kamycheva E, Didriksen A, Jakobsen J, Jorde R. Vitamin D stored in fat tissue during a 5-year intervention affects serum 25-hydroxyvitamin D levels the following year. *J Clin Endocrinol Metab* 2017;102(10):3731–8. doi:10.1210/jc.2017-01187.
- [37] Shin MH, Lee Y, Kim M-K, Lee DH, Chung JH. UV increases skin-derived 1 α ,25-dihydroxyvitamin D3 production, leading to MMP-1 expression by altering the balance of vitamin D and cholesterol synthesis from 7-dehydrocholesterol. *J Steroid Biochem Mol Biol* 2019;195:105449. doi:10.1016/j.jsbmb.2019.105449.
- [38] Krause R, Roth HJ, Kaase H, Stange R, Holick MF. Vitamin D status in chronic kidney disease - UVB irradiation is superior to oral supplementation. *Anti-cancer Res* 2016;36(3):1397–401.
- [39] Slominski AT, Tuckey RC, Jetten AM, Holick MF. Recent advances in vitamin D biology: something new under the sun. *J Invest Dermatol* 2023;143(12):2340–2. doi:10.1016/j.jid.2023.07.003.
- [40] Skobowiat C, Dowdy JC, Sayre RM, Tuckey RC, Slominski A. Cutaneous hypothalamic-pituitary-adrenal axis homolog: regulation by ultraviolet radiation. *Am J Physiol Endocrinol Metab* 2011;301(3):E484–93. doi:10.1152/ajpendo.00217.2011.
- [41] Kiourtzidis M, Kühn J, Brandsch C, Baur A-C, Wensch-Dorendorf M, Stangl GI. Markers indicating body vitamin D stores and responses of liver and adipose tissues to changes in vitamin D intake in male mice. *Nutrients* 2020;12(5):1391. doi:10.3390/nu12051391.
- [42] Holick MF, MacLaughlin JA, Doppelt SH. Regulation of cutaneous previtamin D3 photosynthesis in man: skin pigment is not an essential regulator. *Science* 1981;211(4482):590–3.
- [43] Slominski AT, Kim T-K, Hobrath JV, Janjetovic Z, Oak ASW, Postlethwaite A, et al. Characterization of a new pathway that activates lumisterol in vivo to biologically active hydroxylumisterols. *Sci Rep* 2017;7(1):11434. doi:10.1038/s41598-017-10202-7.
- [44] Tuckey RC, Cheng CYS, Slominski AT. The serum vitamin D metabolome: what we know and what is still to discover. *J Steroid Biochem Mol Biol* 2019;186:4–21. doi:10.1016/j.jsbmb.2018.09.003.
- [45] Al-Zohily B, Al-Menhali A, Gariballa S, Haq A, Shah I. Epimers of vitamin D: a review. *Int J Mol Sci* 2020;21(2):470. doi:10.3390/ijms21020470.
- [46] Ghaly S, Bliuc D, Center JR, Clarke MW, Jones AP, Trend S, et al. Vitamin D C3-epimer levels are proportionally higher with oral vitamin D supplementation compared to ultraviolet irradiation of skin in mice but not humans. *J Steroid Biochem Mol Biol* 2019;186:110–16. doi:10.1016/j.jsbmb.2018.10.002.
- [47] Sohail A, Al Menhali A, Hisaindee S, Shah I. An LC-MS/MS method for analysis of vitamin D metabolites and C3 epimers in mice serum: oral supplementation compared to UV irradiation. *Molecules* 2021;26(17):5182. doi:10.3390/molecules26175182.
- [48] Betters JL, Yu L. NPC1L1 and cholesterol transport. *FEBS Lett* 2010;584(13):2740–7. doi:10.1016/j.febslet.2010.03.030.
- [49] Telford DE, Sutherland BG, Edwards JY, Andrews JD, Barrett PHR, Huff MW. The molecular mechanisms underlying the reduction of LDL apoB-100 by ezetimibe plus simvastatin. *J Lipid Res* 2007;48(3):699–708. doi:10.1194/jlr.M600439-JLR200.
- [50] Tamez H, Kalim S, Thadhani RI. Does vitamin D modulate blood pressure? *Curr Opin Nephrol Hypertens* 2013;22(2):204–9. doi:10.1097/MNH.0b013e32835d919b.
- [51] Kim D-H, Meza CA, Clarke H, Kim J-S, Hickner RC. Vitamin D and endothelial function. *Nutrients* 2020;12(2):575. doi:10.3390/nu12020575.
- [52] Kleinbongard P, Dejam A, Lauer T, Rassaf T, Schindler A, Picker O, et al. Plasma nitrite reflects constitutive nitric oxide synthase activity in mammals. *Free Radic Biol Med* 2003;35(7):790–6. doi:10.1016/S0891-5849(03)00406-4.
- [53] Kelm M, Preik-Steinhoff H, Preik M, Strauer BE. Serum nitrite sensitively reflects endothelial NO formation in human forearm vasculature: evidence for biochemical assessment of the endothelial L-arginine-NO pathway. *Cardiovasc Res* 1999;41(3):765–72. doi:10.1016/S0008-6363(98)00259-4.
- [54] Opländer C, Volkmar CM, Paunel-Görgülü A, van Faassen EE, Heiss C, Kelm M, et al. Whole body UVA irradiation lowers systemic blood pressure by release of nitric oxide from intracutaneous photolabile nitric oxide derivatives. *Circ Res* 2009;105(10):1031–40. doi:10.1161/CIRCRESAHA.109.207019.
- [55] Liu D, Fernandez BO, Hamilton A, Lang NN, Gallagher JMC, Newby DE, et al. UVA irradiation of human skin vasodilates arterial vasculature and lowers blood pressure independently of nitric oxide synthase. *J Invest Dermatol* 2014;134(7):1839–46. doi:10.1038/jid.2014.27.
- [56] Davenport AP, Hyndman KA, Dhaun N, Southan C, Kohan DE, Pollock JS, et al. Endothelin. *Pharmacol Rev* 2016;68(2):357–418. doi:10.1124/pr.115.011833.
- [57] Tsuboi R, Sato C, Oshita Y, Hama H, Sakurai T, Goto K, et al. Ultraviolet B irradiation increases endothelin-1 and endothelin receptor expression in cultured human keratinocytes. *FEBS Lett* 1995;371(2):188–90. doi:10.1016/0014-5793(95)00912-s.
- [58] Metz M, Lammell V, Gibbs BF, Maurer M. Inflammatory murine skin responses to UV-B light are partially dependent on endothelin-1 and mast cells. *Am J Pathol* 2006;169(3):815–22. doi:10.2353/ajpath.2006.060037.
- [59] Martínez-Miguel P, Valdivielso JM, Medrano-Andrés D, Román-García P, Cano-Peñalver JL, Rodríguez-Puyol M, et al. The active form of vitamin D, calcitriol, induces a complex dual upregulation of endothelin and nitric oxide in cultured endothelial cells. *Am J Physiol Endocrinol Metab* 2014;307(12):E1085–96. doi:10.1152/ajpendo.00156.2014.
- [60] Absi M, Ward DT. Increased endothelin-1 responsiveness in human coronary artery smooth muscle cells exposed to 1,25-dihydroxyvitamin D(3). *Am J Physiol Cell Physiol* 2013;304(7):C666–72. doi:10.1152/ajpcell.00349.2012.
- [61] Smith APL, Demoncheaux EAG, Higenbottam TW. Nitric oxide gas decreases endothelin-1 mRNA in cultured pulmonary artery endothelial cells. *Nitric Oxide* 2002;6(2):153–9. doi:10.1006/niox.2001.0400.
- [62] Raoch V, Rodríguez-Pascual F, López-Martínez V, Medrano-Andrés D, Rodríguez-Puyol M, Lamas S, et al. Nitric oxide decreases the expression of endothelin-converting enzyme-1 through mRNA destabilization. *Arterioscler Thromb Vasc Biol* 2011;31(11):2577–85. doi:10.1161/ATVBAHA.111.232025.
- [63] Ahlborg G, Lundberg JM. Nitric oxide-endothelin-1 interaction in humans. *J Appl Physiol* (1985) 1997;82(5):1593–600. doi:10.1152/jappl.1997.82.5.1593.
- [64] Ponda MP, Liang Y, Kim J, Hutt R, Dowd K, Gilleaudeau P, et al. A randomized clinical trial in vitamin D-deficient adults comparing replenishment with oral vitamin D3 with narrow-band UV type B light: effects on cholesterol and the transcriptional profiles of skin and blood. *Am J Clin Nutr* 2017;105(5):1230–8. doi:10.3945/ajcn.116.150367.
- [65] Søyland E, Heier I, Rodríguez-Gallego C, Mollnes TE, Johansen F-E, Holven KB, et al. Sun exposure induces rapid immunological changes in skin and peripheral blood in patients with psoriasis. *Br J Dermatol* 2011;164(2):344–55. doi:10.1111/j.1365-2133.2010.10149.x.
- [66] Milliken SVI, Wassall H, Lewis BJ, Logie J, Barker RN, Macdonald H, et al. Effects of ultraviolet light on human serum 25-hydroxyvitamin D and systemic immune function. *J Allergy Clin Immunol* 2012;129(6):1554–61. doi:10.1016/j.jaci.2012.03.001.
- [67] Slominski AT, Zmijewski MA, Plonka PM, Szafarski JP, Paus R. How UV light touches the brain and endocrine system through skin, and why. *Endocrinology* 2018;159(5):1992–2007. doi:10.1210/en.2017-03230.
- [68] Skobowiat C, Postlethwaite AE, Slominski AT. Skin exposure to ultraviolet B rapidly activates systemic neuroendocrine and immunosuppressive responses. *Photochem Photobiol* 2017;93(4):1008–15. doi:10.1111/php.12642.