

A phase I study of MAGE-A1-targeted T1367 T-cell receptor-based cell therapy in patients with advanced multiple myeloma

Authors

Josefine Krüger,¹ Matthias Obenaus,¹ Igor Wolfgang Blau,² Dana Hoser,³ Martin Vaegler,⁴ Hana Rauschenbach,⁴ Ioannis Anagnostopoulos,⁵ Korinna Jöhrens,⁵ Vivian Scheuplein,⁶ Elisa Kieback,⁶ Judith Böhme,⁵ Ann-Christin von Brünneck,⁵ Jan Krönke,^{1,7} Antonia Busse,^{1,6,7} Gerald Willimsky,^{3,7,8} Thomas Blankenstein,⁶ Antonio Pezzutto,^{1,6} Ulrich Keller^{1,6,7} and Axel Nogai¹

¹Department of Hematology, Oncology and Cancer Immunology, Campus Benjamin Franklin, Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin; ²Department of Hematology, Oncology and Cancer Immunology, Campus Virchow Klinikum, Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin; ³Institute of Immunology, Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, Berlin Institute of Health, Berlin; ⁴Experimental and Clinical Research Center, Zellkulturlabor für Klinische Prüfung ZKP, Charité-

Universitätsmedizin Berlin, Campus Berlin Buch, Berlin; ⁵Institute of Pathology, Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin; ⁶Max-Delbrück-Center for Molecular Medicine, Berlin; ⁷German Cancer Consortium (DKTK), partner site Berlin, a partnership between German Cancer Research Center (DKFZ) and Charité-Universitätsmedizin Berlin, Berlin and ⁸German Cancer Research Center (DKFZ), Heidelberg, Germany

Correspondence:

A. NOGAI - nogai@onkologie-tiergarten.de

U. KELLER - ulrich.keller@charite.de

<https://doi.org/10.3324/haematol.2024.286124>

Received: June 28, 2024.

Accepted: September 2, 2024.

Early view: September 12, 2024.

©2025 Ferrata Storti Foundation

Published under a CC BY-NC license 

A phase I study of MAGE-A1-targeted T1367 T-cell receptor-based cell therapy in patients with advanced multiple myeloma

Josefine Krüger¹, Matthias Obenaus¹, Igor Wolfgang Blau², Dana Hoser³, Martin Vaegler⁴, Hana Rauschenbach⁴, Ioannis Anagnostopoulos⁵, Korinna Jöhrens⁵, Vivian Scheuplein⁶, Elisa Kieback⁶, Judith Böhme⁵, Ann-Christin von Brünneck⁵, Jan Krönke^{1,7}, Antonia Busse^{1,6,7}, Gerald Willimsky^{3,7,8}, Thomas Blankenstein⁶, Antonio Pezzutto^{1,6}, Ulrich Keller^{1,6,7}, Axel Noga¹

¹Department of Hematology, Oncology and Cancer Immunology, Campus Benjamin Franklin, Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, 12203 Berlin, Germany

²Department of Hematology, Oncology and Cancer Immunology, Campus Virchow Klinikum, Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, 13353 Berlin, Germany

³Institute of Immunology, Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, Berlin Institute of Health, Berlin, Germany

⁴Experimental and Clinical Research Center, Zellkulturlabor für Klinische Prüfung ZKP, Charité-Universitätsmedizin Berlin, Campus Berlin Buch, 13125 Berlin, Germany

⁵Institute of Pathology, Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, 10117 Berlin

⁶Max-Delbrück-Center for Molecular Medicine, 13125 Berlin, Germany

⁷German Cancer Consortium (DKTK), partner site Berlin, a partnership between German Cancer Research Center (DKFZ) and Charité-Universitätsmedizin Berlin, 12203 Berlin, Germany

⁸German Cancer Research Center (DKFZ), 69120 Heidelberg, Germany

Supplementary Data:

Supplementary Tables

SUPPLEMENTARY TABLES

Supplemental Table 1. Patient characteristics of the diagnostic MAGE-A1 expression study.

	n (%)
all patients	213 (100)
female	75 (35)
Disease stage	
first diagnosis	92 (43)
relapse	114 (54)
sequentially primary and relapse	7 (3)
Occurrence of EMD at any time of disease	84 (39)
EM-B	66 (31)
EM-E	18 (9)
FiSH results available	131 (62)
standard risk	59 (28)
high risk, defined as...	72 (34)
Gain(1q21) or amplification(1q21)	55 (26)
del(17p)	18 (8)
t(4;14)	21 (10)
t(14;16)	7 (3)

EMD: extramedullary disease

EM-B: bone-related extramedullary myeloma

EM-E: extraosseous extramedullary myeloma

Supplemental Table 2. Phase 1 study of TCR-1367 T cells, patient characteristics.

Patient ID	004	006
Sex	female	male
Age	57	62
EMD	no	yes (EM-B)
MAGE-A1+ myeloma cells in BM at screening [%]	80%	90%
Cytogenetics/FISH	standard risk	high risk (+1q21)
Lines of previous therapies	4	7
Refractory status		
IMiDs	yes	yes
PI	yes	yes
any anti-CD38 antibody	yes	yes
any anti-BCMA directed therapy	no	yes

EMD: extramedullary disease

BM: bone marrow

IMiDs: immunomodulatory drugs

PI: proteasome inhibitors

BCMA: B-cell maturation antigen

Supplemental Table 3. Phase 1 study of TCR-1367 T cells, adverse events.

Patient 004			
Term	Grade	TEAE?	Causality to T-cell therapy
Nausea	2	no	Not related
Pyrexia	1	no	Not related
Pain in extremity	1	yes	Possible
Arthralgia	1	yes	Possible
Cancer pain	1	yes	Possible
Sinus tachycardia	1	yes	Possible
Pyrexia	1	yes	Possible
Pyrexia	1	yes	Possible
Chest pain	1	yes	Possible
Neutropenia	4	yes	Possible
Anemia	2	yes	Possible
Anemia	2	yes	Possible
White blood cell count decreased	2	yes	Not related
White blood cell count decreased	4	yes	Possible

Patient 006			
Term	Grade	TEAE?	Causality to T-cell therapy
Anemia	3	no	Not related
Platelet count decreased	4	no	Not related
Urinary tract infection	2	no	Not related
White blood cell count decreased	3	no	Not related
Nausea	2	no	Not related
Lymphocyte count decreased	4	no	Not related
Dry mouth	1	no	Not related
Depression	2	yes	Not related

Petechiae	1	yes	Not related
Neoplasm progression	1	yes	Not related
Neoplasm progression	3	yes	Not related
Mouth hemorrhage	1	yes	Not related
Monoparesis	3	yes	Not related

TEAE: treatment-emergent adverse events