

The crosstalk between immune activation and metabolism in heart failure. A scientific statement of the Heart Failure Association of the ESC

Gabriele Fragasso¹, Davide Stolfo^{2,3}, Markus S. Anker⁴, Antoni Bayes-Genis⁵, Ovidiu Chioncel⁶, Stephane Heymans⁷, Pardeep S. Jhund⁸, Basil S. Lewis⁹, Gary D. Lopaschuk¹⁰, Lars H. Lund¹¹, Arantxa Gonzalez¹², Matteo Pagnesi¹³, Gabriele Giacomo Schiattarella^{14,15,16,17}, Carlo Gabriele Tocchetti¹⁸, Peter van der Meer¹⁹, Sophie Von Linthout¹⁶, Sven Wassmann²⁰, B. Daan Westenbrink¹⁹, Marco Metra¹³, Giuseppe M.C. Rosano^{21,22}, and Gianluigi Savarese^{3*}

¹Heart Failure Unit, Department of Cardiology, IRCCS Scientific Institute San Raffaele, Milan, Italy; ²Cardiology, Cardiothoracic Department, Azienda Sanitaria Universitaria Friuli Centrale (ASUFC), Udine, Italy; ³Department of Medicine Clinical Science and Education, Södersjukhuset Karolinska Institutet, Stockholm, Sweden; ⁴Deutsches Herzzentrum der Charité, Department of Cardiology, Angiology and Intensive Care Medicine CBF, DZHK, BCRT, FU & HU, Berlin, Germany; ⁵Department of Cardiology, Germans Trias University Hospital, CIBERCV, Badalona, Spain; ⁶Emergency Institute for Cardiovascular Diseases 'Prof. C.C. Iliescu', University of Medicine 'Carol Davila', Bucharest, Romania; ⁷Department of Cardiology, Cardiovascular Research Institute Maastricht, University of Maastricht & Maastricht University Medical Centre, Maastricht, The Netherlands; ⁸BHF Glasgow Cardiovascular Research Centre, University of Glasgow, Glasgow, UK; ⁹Lady Davis Carmel Medical Center and the Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel; ¹⁰Cardiovascular Research Centre, University of Alberta, Edmonton, AB, Canada; ¹¹Division of Cardiology, Department of Medicine, Karolinska Institutet, Stockholm, Sweden; ¹²Program of Cardiovascular Disease, CIMA Universidad de Navarra, Department of Cardiology, Clinica Universidad de Navarra and CIBERCV, Pamplona, Spain; ¹³Institute of Cardiology, ASST Spedali Civili di Brescia, Department of Medical and Surgical Specialties, Radiological Sciences, and Public Health, University of Brescia, Brescia, Italy; ¹⁴Division of Cardiology, Department of Advanced Biomedical Sciences, Federico II University, Naples, Italy; ¹⁵Deutsches Herzzentrum der Charité (DHZC), Charité-Universitätsmedizin Berlin, Berlin, Germany; ¹⁶DZHK (German Centre for Cardiovascular Research), partner site Berlin, Berlin, Germany; ¹⁷Translational Approaches in Heart Failure and Cardiometabolic Disease, Max Delbrück Center for Molecular Medicine in the Helmholtz Association (MDC), Berlin, Germany; ¹⁸Department of Translational Medical Sciences (DISMET), Center for Basic and Clinical Immunology Research (CISI), Interdepartmental Center of Clinical and Translational Sciences (CIRCET), Interdepartmental Hypertension Research Center (CIRIAPA), Federico II University, Naples, Italy; ¹⁹Department of Cardiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands; ²⁰Cardiology Pasing, Munich, and Faculty of Medicine, University of the Saarland, Homburg, Germany; ²¹Department of Human Sciences and Promotion of Quality of Life, San Raffaele Open University, Rome, Italy; and ²²IRCCS San Raffaele, Italy, Milano, Italy

Received 26 September 2024; revised 29 January 2025; accepted 8 May 2025

A better understanding of additional mechanisms of heart failure (HF) progression may allow a different and more complete phenotyping of the disease and identification of novel therapeutic targets. Persistent latent myocardial inflammation/immune activation in HF may represent an attempt to restore tissue homeostasis in the failing heart, where cardiomyocytes and immune cells undergo metabolic reprogramming, which allows them to deal with decreased availability of nutrients and oxygen. This status can trigger a metabolic crosstalk between immune cells and cardiomyocytes which, depending on the outcome, can either perpetuate the maladaptive remodelling of the heart, or determine an adaptive response. Therefore, the interplay between immune activation and metabolism is gaining recognition as a potential therapeutic framework. On these premises, future studies addressing novel HF treatments should attempt to evaluate the potential therapeutic role of direct metabolic and immunological crosstalk modulation. The aim of the present scientific statement from the Heart Failure Association of the ESC is to summarize the current evidence for the connection between inflammatory and immune activation and metabolic adaptation in the onset and progression of HF, in order to promote future strategies for the development of targeted-disease preventive and therapeutic measures.

Keywords

Aging • Heart failure • Immunometabolism • Inflammation • Maladaptive • Myocardial Metabolism • Therapy

*Corresponding author. Department of Clinical Science and Education, Södersjukhuset, Karolinska Institutet, Sjukhusbacken 10, 118 83 Stockholm, Sweden. Email: gianluigi.savarese@ki.se

Introduction

The pathophysiological pathways that initiate and perpetuate heart failure (HF) are complex. Multiple maladaptive and adaptive compensatory mechanisms exist, which span from the subcellular level to organ-to-organ interactions. The contemporary treatments for HF aim at counteracting the haemodynamic impairment and the pathological neurohormonal hyperactivation that characterize the evolution of the syndrome. Nevertheless, increasing evidence shows that other mechanisms, including immunoinflammatory activation and metabolic adaptation, are active part of disease progression, are tightly interconnected to each other, and play a central role in the pathophysiology of HF. Systemic metabolism and the immune system respond in a coordinated manner to stress, such as infection, cancer, or other organ and tissue injuries. Haemodynamic impairment is a trigger for cardiac inflammation and changes in cellular metabolism. Immune cells also play important non-immune functions, including neural development, and modulation of cardiovascular function and metabolism. Experimental and clinical studies on HF pathophysiology have highlighted the role of these factors in the onset and progression of HF, but their comprehension is still partial and a large gap remains in the understanding of their dynamic interplay. The concept of immunometabolism may advance our knowledge of the complex interaction between energetic metabolism pathways and immune cell response.¹

The aim of the present scientific statement from the Heart Failure Association (HFA) of the European Society of Cardiology (ESC) is to summarize current evidence on the connection between inflammatory, immune activation and metabolic adaptation in the development and progression of HF, in order to outline future strategies for the identification of targeted-disease preventive and therapeutic measures.

Metabolic and immune adaptation in heart failure: pathophysiological concepts

Cardiac metabolic changes in heart failure

The heart requires a large amount of adenosine triphosphate (ATP) to sustain contractile function, which is obtained from mitochondrial oxidative phosphorylation and glycolysis. The healthy heart is 'metabolically flexible', and readily shifts between fatty acids, lactate, glucose, ketones, and amino acids as fuel sources to maintain ATP production.² In contrast, the failing heart loses its metabolic flexibility and can become energy deficient due to a decrease in ATP production,^{3–5} primarily due to a reduced mitochondrial oxidative capacity and impaired mitochondrial creatine kinase energy metabolism.^{6–11} These altered mitochondrial dynamics are important contributors to the overall reduction in mitochondrial oxidative capacity and therefore ATP production in HF (Figure 1).

Decreased mitochondrial oxidative metabolism in HF leads to the induction of glycolysis in an attempt to compensate

for the decrease in mitochondrial ATP production.^{3–5} However, this increase in glycolysis is inefficient and not accompanied by enhanced mitochondrial oxidation of pyruvate derived from glycolysis.^{12–22} Inhibition of cardiac mitochondrial pyruvate uptake results in the development of HF in mice.^{23,24} The reduction in pyruvate utilization may also contribute to adverse left ventricular remodelling.^{15,23,25–27}


Heart failure is also associated with alterations in cardiac fatty acid oxidation. It is reported that reductions in mitochondrial fatty acid oxidation occur in HF, but data are not consistent across different studies.^{23,26,28–39} Myocardial fatty acid oxidation also increases in response to conditions such as type 2 diabetes (T2D), obesity and insulin resistance.^{40–43} Despite the close association, a causal role for increased fatty acid oxidation rates in cardiac dysfunction under obesity or diabetic conditions is less certain. Since cardiac lipid uptake is increased in these models, an imbalance between lipid uptake and oxidation likely contributes to lipotoxicity.

Cardiac ketone oxidation is also altered in the failing hearts. The increase in ketone metabolism is thought to be an adaptive process in HF,⁴⁴ and may provide the 'starving' failing heart with an extra source of ATP.

In summary, the failing heart is energy deficient, primarily due to a decrease in mitochondrial oxidative capacity, an increased glycolysis uncoupled with glucose oxidation, and either a decrease or no change in fatty acid oxidation. In contrast, if there is increased fatty acid availability, as occurs in obesity, an increase in cardiac fatty acid oxidation may occur. These energy metabolic changes result in the failing heart becoming less efficient.

Immunological changes in heart failure

The primary role of an inflammatory response is to resolve the source of the disturbance, thereby allowing the involved tissues to ultimately restore their function. In the setting of chronic HF, while some inflammation may be required to repair a short-term injury and therefore be protective, a persistent myocardial and systemic inflammatory state takes place and can contribute to disease progression.⁴⁵ The enduring expression of pro-inflammatory cytokines and ongoing inflammation that have been demonstrated in the failing heart indicate a state of ongoing chronic inflammation that is intermediate between normality and acute inflammation.⁴⁶ These pro-inflammatory cytokines contribute to the pathogenesis and progression of HF primarily by left ventricular remodelling and dysfunction, and secondarily by interacting with the neurohormonal system and cardiac metabolism.⁴⁷ Activation of neurohormonal systems in HF further contributes to maintain systemic and cardiac inflammation.⁴⁸ Both the innate and adaptive immune systems have a pro-inflammatory role in HF. The immune response-triggered inflammation mechanism is called immune or sterile inflammation. This intermediate state has also been termed para-inflammation, and does not require overt tissue injury or infection for it to be sustained, but instead represents a sustained inflammatory response aimed at restoring tissue functionality.⁴⁹ Since HF is generally characterized by decreased availability of nutrients and oxygen, cardiomyocytes and non-cardiomyocytes, including fibroblasts



and immune cells, undergo metabolic reprogramming in order to adapt to this condition.⁵⁰ Competition for nutrients and increased production of signalling metabolites initiate a metabolic crosstalk between immune cells and cardiomyocytes, which is proposed to tip the balance between resolution of inflammation versus adverse cardiac remodelling.⁵¹ With glucose metabolism and ATP generation being deeply involved in immune cell activation, proliferation, trafficking and effector functions, reprogramming of the metabolic status of immune cells leads to changes in their functional properties and supports phenotype switching between different subtypes^{52–56} (Figure 2).

Diet and HF-associated gut microbiota further affect immune cell metabolism and foster chronic inflammation and ineffective adaptive immunity. This implies that therapeutic modulation of systemic metabolism can re-establish immune cell homeostasis and modulate adverse cardiac remodelling. Caloric excess shortens lifespan, promotes systemic inflammation and increases the risk of cardiometabolic diseases.⁵⁷ In contrast, caloric restriction and fasting have been shown to be inversely associated with cardiovascular diseases, and there is now an increasing attention to its therapeutic potential.⁵⁸ In HF, and especially in obese patients with

a preserved ejection fraction phenotype, caloric restriction or fasting programmes, alone or in association with exercise training, improved peak oxygen uptake at cardiopulmonary exercise testing and quality of life.^{59,60}

However, prolonged fasting and reduction in caloric intake can have also negative effects, particularly on immunity regulation. During starvation, organs and systems set a sparing energy process with a survival-driven hierarchy. The production of immune cells slows down relatively early in the process of conserving resources, potentially increasing the infective risk. A recent study has shown that during nutrient scarcity the marrow is a safe haven for monocytes and re-feeding prompts mobilization culminating in monocyte apoptosis of chronologically older and transcriptionally distinct monocytes.⁶¹ These shifts could alter the adaptation to external stressors response, outlining that diet, in particular a diet temporal dynamic balance, modulates monocyte lifespan, with consequences on regulating the immune response, inflammation, and tissue repair. An appropriate regulation of immune cells turnover is vital for overall health and immune function, therefore the definition of the optimal balance is necessary to preserve the organism's immunological response.




Figure 2 Immunological changes in heart failure. Myocardial injury can trigger the pathological process leading to heart failure involving multiple mechanisms. Indeed, the classical neurohormonal activation mediated by the circulatory and mechanical stress is paired with immunometabolic changes which flow in a state of persistent inflammation. The process is mediated by the activation of both adaptive and innate immunity and their interaction. The maladaptive remodelling of the heart resulting from the negative effects of each of the involved mechanisms is characterized by several alterations in cellular functions and a pro-fibrotic state. ECM, extracellular matrix; NOD, nucleotide-binding oligomerization domain-like; TLR, Toll-like receptor.

Adaptive role of immunity and interplay between metabolism and inflammation

Although immune activation is typically maladaptive in HF, it is crucial to acknowledge that inflammation is a fundamentally adaptive process (Figure 3). In response to viral myocarditis for instance, inflammation initially serves an adaptive purpose, but excessive immune cell activation can cause a switch from host defence to injury.⁶² In fact, in HF the adaptive immune system is dysregulated, with an exaggerated inflammatory response that contributes to the progression of the disease. Macrophages, neutrophils, T cells, B cells, and humoral immunity all play key roles in driving chronic inflammation, fibrosis, and tissue remodelling. Understanding the complex interactions between these immune cells and pathways is crucial for developing targeted therapies to modulate the immune response and potentially slow the progression of HF.

Macrophages are central to the inflammatory process in HF and play both protective and detrimental roles, depending on their polarization and activation state. In HF, mechanisms of activation are altered, which contributes to chronic inflammation, fibrosis, and adverse cardiac remodelling. M1 macrophages secrete pro-inflammatory cytokines such as tumour necrosis factor- α , interleukin (IL)-1, and IL-6, which exacerbate cardiac inflammation and injury. Increased recruitment and activation of macrophages in the heart tissue may contribute to myocardial inflammation, fibrosis, and remodelling by secreting various cytokines, chemokines, and matrix metalloproteinases.⁶³

Neutrophils are part of the innate immune response and are actively involved during acute exacerbations or after myocardial infarction, infiltrating the heart and releasing pro-inflammatory cytokines, reactive oxygen species (ROS) and enzymes that contribute to tissue damage and amplification of inflammation. In chronic HF, sustained neutrophil activation contributes




Figure 3 Adaptive role of immunity and interplay between metabolism and inflammation in heart failure. Inflammation plays a significant role in the pathophysiology of heart failure, acting both as a potentially adaptive response (blue lines) and as a contributor to disease progression (violet lines). Heart failure is both driven by and exacerbates a state of chronic inflammation, characterized by elevated levels of inflammatory cytokines, which are involved in recruiting immune cells to sites of injury, helping the removal of dead cells and promoting tissue repair and remodelling. Inflammatory signals can also stimulate angiogenesis. Additionally, inflammation contributes to the structural and functional remodelling of the heart in response to injury or stress by promoting adaptive cardiac hypertrophy. Finally, inflammatory cytokines are involved in remodelling the extracellular matrix of the heart by regulation of matrix metalloproteinases. Activated immune cells, such as macrophages and T cells, shift their metabolism towards glycolysis. This metabolic shift supports rapid proliferation and function of immune cells, promoting their activation and prolonging the adaptive inflammatory response. Inflammatory signals can also contribute to energy metabolism modification in cardiac cells to meet the increased energy demands during stress. This includes a shift from fatty acid oxidation to glucose utilization, which is a more efficient metabolic pathway, especially under ischaemic conditions. Inflammatory cytokines can activate the sympathetic nervous system (SNS) and the renin–angiotensin–aldosterone system (RAAS) which help to maintain blood pressure and perfusion of vital organs. Brown lines represent functions with conflicting or unclear evidence. While inflammation serves adaptive purposes, chronic and excessive inflammation can become maladaptive, contributing to the progression of heart failure. Persistent inflammation can lead to continuous cardiac remodelling, myocyte apoptosis and drive systemic effects contributing to cachexia and worsening overall health.

to ongoing inflammation, oxidative stress, and endothelial dysfunction.⁶⁴

T cells have been found to play both beneficial and harmful roles in HF, contributing to both the adaptive immune response and the chronic inflammation. T helper cells, particularly Th1 and Th17 cells, are associated with increased inflammation and tissue damage in HF with the production of cytokines such as interferon- γ and IL-17, which are promoters of enhanced macrophagic response and inflammatory activation, myocardial fibrosis and adverse remodelling.⁶⁵ Regulatory T cells, that typically act as down-modulators of immune activation and inflammation, are down-regulated in HF. There is also evidence suggesting that cytotoxic T cells (CD8⁺) may contribute to myocardial injury by inducing cell death in the heart tissue, exacerbating the inflammatory milieu and HF progression.⁶⁶

B cells in HF may produce autoantibodies leading to autoimmunity. For example, autoantibodies directed against β -adrenergic receptors have been observed in HF patients.⁶⁷ These antibodies can impair receptor function, contributing to cardiac dysfunction. In some forms of HF, particularly in the context of myocarditis or autoimmune diseases, B cells may infiltrate the myocardium and contribute to local inflammation and tissue damage.⁶⁸

Additionally, multiple cardiac cell types, including myocytes, fibroblasts, and vascular cells, actively contribute to inflammation in response to pressure overload, suggesting that cardiac inflammation may also serve initially as an adaptive response to cardiac stress.⁶⁹ For example, the NLRP3 inflammasome is the activated downstream of calcium signalling pathways in response to haemodynamic stress, which implies that inflammation is partly driven by cardiac autonomous mechanisms.⁷⁰ Furthermore, while inhibition of immune cell activation in HF models decreases disease severity and slows down the progression of cardiac hypertrophy and fibrosis, anti-cytokine therapies have not yet provided significant clinical benefits in patients with HF.⁷¹ Nevertheless, since persistent inflammation can lead to continuous cardiac remodelling and myocyte apoptosis and determine systemic effects contributing to cachexia and worsening overall health, further research into therapeutic anti-inflammatory strategies is warranted.

The interplay between inflammation and metabolism is gaining recognition as a potential underlying factor in HF.⁷² Of particular interest are the recent observations relative to the potential harmful role of epicardial adipose tissue via paracrine or vasocrine secretion of pro-inflammatory and pro-fibrotic cytokines, leading

to both local and systemic inflammation.⁷³ In particular, inflamed epicardial adipose tissue may provide a direct source of cardiac inflammation due to its close proximity to the myocardium.⁷⁴ Second, research has revealed that inflammatory cytokines, such as IL-6, can induce pathological cardiomyocyte hypertrophy, which could contribute to the typical metabolic changes observed in HF.⁶⁹ Third, cytokines have the ability to directly target metabolic pathways, as evidenced by the insulin resistance and suppression of the metabolic sensor AMPK that was observed following IL-6 infusion in a HF with preserved ejection fraction (HFpEF) model.⁷⁵ Finally, the increase in ROS production associated with inflammation can lead to mitochondrial damage and dysfunction, further exacerbating metabolic derangements.⁶⁹ A confounding aspect is that the interplay between inflammation and metabolism may differ between HF with reduced ejection fraction (HFrEF) and HFpEF. In general, inflammation is retained to develop during the course of HF as a response to remodelling, haemodynamic impairment and neurohormonal activation in patients with HFrEF, whereas it may be the primary cause of diastolic dysfunction and hypertrophy in HFpEF as a consequence to extra-cardiac conditions such as diabetes, metabolic syndrome, chronic kidney disease.⁷⁶ These findings underscore the critical interplay between inflammation and metabolic dysregulation in HF, supporting the need for more in-depth research into the mechanisms.

Metabolically trained immunity

Long-term epigenetic reprogramming of innate immune cells in response to microbes is known as 'trained immunity', and causes prolonged altered cellular functionality to protect from secondary infections.⁷⁷ Immune cells, especially macrophages, reprogram their metabolic pathways to support an enhanced inflammatory response. This often involves a shift from oxidative phosphorylation to glycolysis, which supports a robust immune response. In addition to metabolic reprogramming, trained immunity involves epigenetic modifications that alter gene expression. 'Trained' immune cells can provide a stronger response to infections and inflammatory injuries. This has positive effects in terms of effective pathogens' clearance but can also be responsible for a hyper-inflammatory response.

In a recent animal study, a sterile trigger of inflammation such as western diet feeding, has been shown to generate increased proliferation and enhanced innate immune response, with the NLRP3 inflammasome appearing to be critical for this response.⁷⁷ Hence, the use of small molecule inhibitors that block the NLRP3 signalling pathway could represent a new therapeutic approach to better modulate the innate immune response following inflammatory triggers and thereby mediate the potentially deleterious effects of trained immunity in chronic inflammatory conditions.

Immunometabolic abnormalities in specific conditions

Ageing

The principal characteristic of cardiac ageing is a decrease of tissue capacity of recovery and regeneration (Figure 4). Along with

increased development of tissue fibrosis, the number of cardiomyocytes and their turnover declines progressively with ageing.^{78,79} As discussed above, the immune system plays an important role in the development of cardiac damage and progression of HF, especially in the elderly.⁸⁰ It is well established that the inflammatory system is chronically over activated in patients with HF, and that there is an association with patient outcome.⁸¹ The immune response is also dynamically remodelled with ageing, determining a phenomenon described as 'immunosenescence'.^{82,83} This phenomenon increases susceptibility to multiple clinical conditions such as infections, autoimmune disorders, and malignancies, and causes a subclinical chronic pro-inflammatory state also known as 'inflamm-aging'.^{84,85} A possible sequence of events could be initiated by cardiac ageing, which is associated with decline in mitochondrial function and accumulation of dysfunctional mitochondria^{86,87} producing and accumulating ROS that, in turn, could lead to accumulation of damaged DNA, proteins, and lipids as well as mitochondrial DNA damage and release.⁸⁸ In older subjects, ROS could increase the release of mitochondrial DNA from the cellular cytosol,⁸⁹ which produces damage-associated molecular patterns, and is considered to be a driver of inflammatory responses.⁹⁰ Inflammatory cells release fibrogenic cytokines and growth factors stimulating the reparative process, characterized by fibroblasts undergoing proliferation to replace lost cardiomyocytes and resulting in remodelling and fibrosis.⁹¹ In this context, preservation of mitochondrial morphology, dynamics, and function might be a therapeutic approach to prevent cardiac ageing. There are several potential drugs and nutraceuticals that could potentially improve age-related mitochondrial dysfunction and, thus, attenuate the process of cardiac 'inflamm-aging'.^{92–95} Interestingly, physical exercise has also been shown to preserve mitochondrial health and metabolism and, as a consequence, to reduce 'inflamm-aging'.^{96,97}

Finally, clonal haematopoiesis of indeterminate potential (CHIP), the clonal expansion of blood stem cells with preleukemic acquired genetic variants, is an age-related condition affecting approximately 10% of individuals aged at least 70 years.⁹⁸ Recent work suggests CHIP is also an independent risk factor for incident HF, with experimental models indicating a role for CHIP in the development of cardiac dysfunction through inflammatory dysregulation and fibrotic remodelling.⁹⁹ These findings may have implications for the prevention and management of HF in the elderly, including development of targeted therapies.

In conclusion, the balance between adaptive and maladaptive immune activation and metabolism in senescence is crucial for determining the overall impact on health and ageing. Understanding this balance can guide therapeutic strategies to promote healthy ageing and prevent age-related diseases.

Obesity

In obesity, metabolic alterations and immune dysregulation are intertwined.^{72,100} Changes in systemic- and tissue-specific metabolism drive immune alterations and vice versa, participating to the pathogenesis of HF in subjects with obesity.^{101,102} The crosstalk between immunity and metabolism in concomitant obesity and HF is bidirectional. The perspective of identifying an

skeletal muscle play a pathophysiological role in T2D-associated HF, but also abnormal cardiac extracellular matrix deposition, metabolic disturbances, oxidative stress and inflammatory pathways may contribute to adverse cardiac remodelling.^{122,123} The interplay between cardiomyocytes and non-myocytes in the heart (fibroblasts, vascular cells, autonomic neurons, and immune cells) is crucial.¹¹⁸ Excessive and sustained local inflammation in response to a persistent stress/trigger such as T2D may be 'maladaptive' and favour cardiac injury. T2D-mediated oxidative stress enhances a pro-inflammatory milieu and promotes the mobilization of leucocytes (neutrophils, monocytes, macrophages and lymphocytes).^{118,121} Rather than favouring the polarization of macrophages towards a reparative M2 phenotype, the secretion of pro-inflammatory cytokines by neutrophils alters the macrophage function and promotes an inflammatory M1-like phenotype, further enhancing macrophage-mediated local inflammation and injury.^{118,124} Abnormalities in T-cell signalling and regulation have also been described, further contributing to cardiac inflammation, hypertrophy and fibrosis.^{118,125–128} Furthermore, cardiac autoimmunity may have a role, especially among patients with type 1 diabetes. In both type 1 and type 2 diabetes, chronic hyperglycaemia causes subclinical myocardial injury with consequent exposure of heart muscle proteins to the immune system, thus a dysregulated adaptive immune system (typical of type 1 diabetes with poor glycaemic control) may overreact, leading to a local pro-inflammatory state and to the development of autoantibodies to cardiac antigens.^{129,130}

Gut microbiome

In the presence of HF, low cardiac output and venous fluid overload contribute to a decrease in intestinal perfusion, mucosal ischaemia, and ultimately in a disrupted intestinal mucosa and increased gut permeability.^{131,132} By this means, gut microbiome-derived metabolites, like trimethylamine-N-oxide (TMAO), can reach the systemic circulation, fuel metabolic inflammation and HF exacerbations.^{133,134} TMAO levels are in fact a strong predictor of clinical outcomes in patients with HF regardless of the underlying aetiology.¹³⁵ The HF-associated gut luminal hypoxia and decrease in mucosal pH further alter the microbiota to pathogenic microbiota and lead to loss of microbial diversity or 'gut dysbiosis', which in turn contributes to the raise in gut permeability and systemic inflammation.^{136,137} On the other hand, different gut microbiome compositions have varying abilities to generate TMAO, and higher levels of circulating TMAO can be attributable to a TMA-producing microbiome harbouring TMA lyases.¹³⁸

Vice versa, there is accumulating evidence showing that under the influence of certain environmental factors and host genetic susceptibility, interactions between the microbiome and the host immune system contribute to cardiovascular disease and HF.¹³⁹ Risk factors of HF, including Western lifestyle with high salt composition, can lead to hypertension and cardiovascular disease involving reduced survival of *Lactobacillus murinus* and increased T_H17 cells.¹⁴⁰ Western diet, characterized by a high intake of fat and a low intake of fiber, decreases the production of short chain fatty acids (SCFAs) by intestinal microbiota, accounting

for the altered regulatory T-cell frequency in mesenteric lymph nodes.¹⁴¹ On the other hand, caloric restriction modulates gut microbiota composition, leading to regulatory T-cell expansion.¹⁴² In obesity, microbiome-mediated tryptophan metabolites modulate white adipose tissue inflammation,¹⁴³ fostering low-grade systemic inflammation and HFpEF.⁷⁶ In metabolic dysfunction, the gut contains increased numbers of (γδ)T cells, macrophages, dendritic cells, NK cells, CD8+ cells (αβ TCR) and Th1 T cells but less T regulatory cells,¹⁴⁴ linked to intestinal barrier dysfunction, and intestinal dysbiosis. Evidence for the specific involvement of gut immune cells in cardiovascular disease and HF comes from integrin β7 (ITGb7) knockout mice, which lack intraepithelial (αβ) and (γδ)T cells, B cells and myeloid cells in the gut.¹⁴⁵ ITGb7, LDLr knockout mice fed with a high cholesterol diet exhibit lower adipose tissue inflammation, reduced numbers of Ly6C^{high} monocytes and smaller aortic lesions.¹⁴⁶ ITGb7 knockout mice are further protected from hypertension, obesity and diabetes and have higher levels of the gut incretin hormone glucagon-like peptide-1 (GLP-1) due to absence of gut immune cells expressing GLP-1 receptor.¹⁴⁶ GLP-1 is increased in HF independent of food intake as an endogenous protective counter-regulatory response.¹⁴⁷ GLP-1 receptor agonists have been shown to decrease hospitalization for HF in T2D patients by 11%.^{148,149} Further, prospective randomized controlled clinical trials in patients with HFpEF and obesity, comparing the GLP-1 receptor agonist semaglutide with placebo, have shown a decrease in body weight with an improvement in quality of life and exercise tolerance altogether with a reduction in C-reactive protein and N-terminal pro-B-type natriuretic peptide (NT-proBNP) plasma concentrations, consistent with a reduction in inflammatory activation and an improvement in diastolic function with the study drug.^{150,151} A pre-specified pooled analysis of these trials has also shown better outcomes with semaglutide versus placebo.¹⁵²

Besides the abovementioned mutual interplay between HF and microbiota/dysbiosis-induced systemic inflammation and (extra)-intestinal immunity, it has been shown that cardiac myosin-specific T_H17 cells can be generated, imprinted in the intestine by a commensal *Bacteroides* species peptide mimic, promoting progression of myocarditis to lethal heart disease.¹⁵³ This is supported by both the successful prevention of lethal disease in mice by antibiotic therapy and the significantly elevated *Bacteroides*-specific CD4⁺ T-cell and B-cell responses observed in myocarditis patients. Taken together, these findings highlight a strong mutual inter-relation between the gut and the heart, involving metabolism and the immune system.

Gender

The differences between men and women in epidemiology, pathophysiology, outcomes and treatment of HF are very broad.¹⁵⁴ The hormonal background is considered to yield a protective effect on the cardiovascular system in young women. In women, after menopause the loss of the hormonal balance leads to several changes which increase the risk of HF, and in particular HFpEF.^{155,156} Metabolic dysregulation and pro-inflammatory state are only two of the multiple mechanisms behind the different pathogenesis and manifestation of HF in the two sexes.

Microvascular dysfunction, together with the hyperactivation of systemic and local inflammation, is a major determinant of HFpEF especially in women, whereas in men the typical atherosclerotic degeneration of the epicardial coronary circulation predominates.¹⁵⁷ Impaired activation of the endothelial nitric oxide pathway and release of pro-angiogenic factors trigger microvascular dysfunction enhanced by the post-menopausal reduction in oestrogens.^{156,158} An additional contribution to microvascular alterations in women comes from the immune-mediated predisposition to endothelial inflammation. Adipokines and inflammatory markers such as leptin and C-reactive protein, both central in the pathogenesis of HFpEF, are indeed expressed to a greater extent in women compared with men, regardless of the post-menopausal period.¹⁵⁹ This is linked with the differences in immune response which is enhanced in women, ensuring prompter recovery from infection and higher protection towards cancer, but the increased systemic inflammatory activation and the stimulation of multiple pro-inflammatory pathways mediated by the stronger immune response indirectly expose women to higher risk of HFpEF.¹⁶⁰ It is of interest and worthy of more deep investigations that sex-specific risk factors (i.e. sex hormones, pregnancy and breast cancer therapy) are strongly linked each other and involved in the pathogenesis of sex-specific diseases or sex-prevalent diseases, such as peripartum cardiomyopathy and stress cardiomyopathy. The common pathophysiological mechanisms behind these diseases involve indeed endothelial inflammation and microvascular dysfunction.^{161,162}

Dysfunctional tissue and systemic metabolism characterizing HF also presents some discrepancies in women and men. In men, increased afterload leads to more pronounced down-regulation of lipid and energy metabolism related genes and this implies less preservation of mitochondrial function compared with women.^{163,164} On the other hand, females show an up-regulation of peroxisome-dependent lipid utilization genes representing an alternative pathway to cover greater energy demand for the myocardium exposed to increased afterload.¹⁶⁵ Oestrogens are important mediators of mitochondrial function, exerting through different mechanisms a protective effect under the influence of detrimental stressors,^{166–168} and also stimulate the activation of antioxidative pathways.

Therapeutic approaches to inflammation and metabolism in heart failure

Therapeutic targets

As outlined above, it is evident that normal cellular homeostasis relies on the crosstalk between the immune system and metabolic regulation, and that several diseases are produced and exacerbated by their dysfunction and crosstalk. Apart from primary inflammatory/metabolic/immune diseases, the plausible biological mechanism in most chronic HF phenotypes is primarily a mechanical incompetence which, in turn, determines metabolic adaptation and inflammatory/immune consequences. The targets

of the presently available drugs for the treatment of HF are mainly maladaptive neurohormonal activation and, consequently, improving cardiac mechanical function, which ultimately regulates also metabolic and immune adaptations. This precise sequence of events is deduced by the unsuccessful results of clinical trials of anti-inflammatory therapies in HF, despite the fact that the chronic inflammatory state associated with progression and complications of HF had identified this dysregulated immunologic state as a potential additional therapeutic target.^{169,170} On the other hand, improvement of metabolic efficiency may limit inflammation by promoting endogenous anti-inflammatory effects, providing strong evidence for the therapeutic potential also in targeting metabolic processes as for controlling immune effector functions and thereby alleviating pathological inflammation,¹⁷¹ again likely consequent to cardiac mechanical incompetence. Nevertheless, intensive⁷¹ research is still active to define the potential therapeutic role of anti-cytokine therapy in HF. Some studies employing anakinra, canakinumab and colchicine have shown a significant reduction in inflammatory biomarkers along with an improvement in exercise performance and quality of life and a reduction in natriuretic peptides.^{172,173} A clinical trial with ziltivekimab is still ongoing.¹⁷⁴

In the real world, evidence-based treatments for HF may contribute to modulate the metabolic and immunological crosstalk in HF.^{52,175} (Table 1). HF induces neurohormonal and immune activation, which directly and indirectly modify myocardial metabolism. The autonomic nervous system contributes to the regulation of cardiac function. Enhanced sympathetic stimulation induced by cardiac mechanical failure increases myocardial energy metabolic requirements. Additionally, sympathetic hyperactivity is associated with increased production of myeloid immune cells in the bone marrow, resulting in higher numbers of circulating myeloid immune cells.¹⁷⁶ Recent studies have shown that microglia, known as brain-resident immune cells, may play an important role in regulating sympathetic nervous system activities and cardiovascular function by releasing cytokines, chemokines, and growth factors.^{176–178} Beta-blockers blunt the cardiovascular response to adrenergic stimulation and, apart from reducing ischaemia and protecting the infarcted myocardium, have also been shown to directly affect myocardial energetics, by reducing circulating levels of free fatty acids (FFA) and, due to substrate competition, inducing a shift of heart energy metabolism towards a greater utilization of carbohydrates^{179–181} (Table 1). The observed increase in cardiac carbohydrate metabolism after beta-blockade probably results from decreased FFA delivery and oxidation, as well as from augmented arterial glutamate availability.^{182–186} Therefore, a higher rate of carbohydrate utilization induced by beta-blockade may result in a greater cardiac energy production at similar levels of oxygen consumption. Interestingly, there are profound differences in the metabolic profile of the non-cardioselective and selective beta-blockers, the former causing significant inhibition of the lipolytic, glycogenolytic and the growth hormone-releasing effects of adrenaline when compared with the cardioselective ones.¹⁸⁷ Additionally, catecholamines play an important role in the regulation of the physiological immune response and, therefore, the efficacy of beta-blockers in HF may also depend by their crucial immunoregulatory role in modifying a dysregulated cytokine

Table 1 Immunometabolic effects of established and potential medical therapies for heart failure

Drug	Metabolic effects
Beta-blockers	<ul style="list-style-type: none"> • Reduction of peripheral lipolysis • Increased carbohydrate utilization • Improved insulin sensitivity
RAAS inhibitors	<ul style="list-style-type: none"> • Immunoregulatory role in modifying a dysregulated cytokine network • Improved glucose homeostasis by increased blood flow in skeletal muscle • More efficient insulin release • Reduction of lipolysis • Anti-inflammatory effect
Partial fatty acid oxidation inhibitors	<ul style="list-style-type: none"> • Enhancement of insulin sensitivity • Increased glucose oxidation • Modulation of late sodium current, thereby reducing the accumulation of intracellular Ca^{++} • Attenuation of macrophage infiltration and pro-inflammatory responses in sepsis-induced myocardial dysfunction
Coenzyme Q10	<ul style="list-style-type: none"> • Inhibition of the mitochondrial permeability transition pore • Inactivation of apoptotic cascades and the oxidative inactivation of key proteins involved in ATP production
SGLT2i	<ul style="list-style-type: none"> • Increased bioavailability and myocardial utilization of ketone bodies • Ketone body-mediated anti-inflammatory effects • Inhibition of epicardial adipose tissue accumulation and inflammation
Statins	<ul style="list-style-type: none"> • Positive influence on pro-inflammatory cytokine production, immune cell migration and T-cell signalling • Inhibition of epicardial adipose tissue accumulation and inflammation
GLP-1 receptor agonists	<ul style="list-style-type: none"> • Mitigation of inflammation in visceral fat adipocytes • Modulation of the immune system
Adipose triglyceride lipase inhibitors	<ul style="list-style-type: none"> • Improved adipose tissue storage capacities • Reduced lipolytic activity
Ketone bodies and short-chain fatty acids	<ul style="list-style-type: none"> • Anti-apoptotic, anti-fibrotic, and anti-inflammatory actions • Alternative fuel sources • Inhibition of NLRP3 inflammasome formation

ATP, adenosine triphosphate; GLP-1, glucagon-like peptide-1; RAAS, renin–angiotensin–aldosterone system; SGLT2i, sodium–glucose cotransporter 2 inhibitor.

network.^{188–191} Indeed, in patients with HF, beta-blockers have been shown to reduce the expression of inflammation, regardless of left ventricular functional response.¹⁹²

The renin–angiotensin–aldosterone system (RAAS) is another system that is hyperactivated in presence of cardiac mechanical dysfunction. Apart from regulating blood pressure, angiotensin II is also an important modulator of cardiac energy metabolism and function. Angiotensin II damages mitochondria in the cardiomyocyte by increasing ROS production,^{193,194} and affects mitochondrial oxidative phosphorylation, including FFA oxidation.^{19,193} There is also evidence that angiotensin II regulates glucose oxidation and that inhibition of angiotensin II may exert beneficial metabolic effects.^{195–198} In addition, by decreasing oxidative metabolism, angiotensin II can reduce ATP production.¹⁹⁹ In this context, apart from blood pressure and fluid balance regulation, angiotensin II antagonism represents an attractive therapeutic approach to target metabolic deregulation in HF. Additionally, as for beta-blockers, RAAS antagonism has been shown to yield a significant reduction in markers of inflammation in different cardiological contexts. Angiotensin-converting enzyme inhibitors

(ACEi), compared with angiotensin receptor blockers, appear to produce a more robust anti-inflammatory effect.²⁰⁰ However, the role of ACEi in inflammation is still debated. Intrinsic ACE has been detected in macrophages and neutrophils, where its overexpression results in enhanced immune responses, independent of angiotensin II. In contrast, ACE activity is also elevated in certain autoimmune diseases and its inhibition benefits patient outcome where inflammatory immune cells are overactive. Further research on the potential role of ACEi in modulating immune response are definitely warranted.²⁰¹

Cardiac metabolism in HF can also be beneficially targeted by specific drugs, by improving mitochondrial membrane stability, mainly dependent on cardiolipin and coenzyme Q10.^{202,203} Coenzyme Q10 is an obligatory component of the respiratory chain in mitochondria and plays an essential role in ATP formation in most tissues, including the heart. In addition, Q10 has antioxidant properties and protects circulating low-density lipoprotein particles from oxidation. Its inhibition of the mitochondrial permeability transition pore prevents the activation of apoptotic cascades and the oxidative inactivation of key proteins involved in




Figure 5 Targeting other sources of inflammation. Statins exert several beneficial pleiotropic effects in the vasculature and cardiac tissue. Obese visceral adipose tissue and epicardial adipose tissue promote pro-inflammatory and lipid-associated pathological effects as well as pro-fibrotic actions, negatively impacting on cardiomyocyte function and survival as well as on cardiac fibrosis. Statins and several other established (in part non-cardiac) and experimental pharmacological treatments inhibit these negative effects, positively influencing immunometabolism of the heart. ATGL, adipose triglyceride lipase; GLP1-RA, glucagon-like peptide-1 receptor agonist; MRA, mineralocorticoid receptor antagonist; SGLT2i, sodium–glucose cotransporter 2 inhibitor.

ATP production²⁰⁴ (Table 1). Lower levels of coenzyme Q10 are associated with worse HF symptoms, lower ejection fraction and higher NT-proBNP.^{205,206} However, association between its values and mortality remains controversial.²⁰⁷ Coenzyme Q10 supplementation improved HF symptoms and reduced major cardiovascular outcomes in patients with HF in a randomized controlled trial.²⁰⁸ However, a more recent meta-analysis, due to the low quality of available data, concluded for no convincing evidence supporting the use of coenzyme 10 in patients with HF.²⁰⁹

In addition, beneficial effects on cardiac metabolism can be achieved with the direct inhibition of oxidative phosphorylation by shifting energy production from fatty acid to glucose oxidation by selective block of mitochondrial long chain 3-ketoacyl coenzyme A thiolase (3-KAT) activity, the last enzyme involved in beta-oxidation²¹⁰ (Table 1). This approach has been shown to preserve phosphocreatine and ATP intracellular levels in the failing heart and to exert significant beneficial effects in patients with ischaemic and non-ischaemic left ventricular dysfunction, especially in HF patients who also have diabetes.^{211–213} More recently, partial FFA oxidation inhibition with trimetazidine has been shown to attenuate macrophage infiltration and pro-inflammatory responses in sepsis-induced myocardial dysfunction. These effects are achieved by normalizing the inflammatory response pathway in macrophages.²¹⁴ Similar anti-inflammatory effects have been observed with ranolazine, another partial FFA oxidation inhibitor, which has also been shown to inhibit the cellular late inward sodium current.^{215–217} In particular, the demonstration that functional Na⁺ shuttling is required for a full cellular response to inflammation and that inhibition of Na⁺–Ca²⁺ exchange during

inflammation by ranolazine reduces the inflammatory response in human endothelial cells *in vitro*, in a mouse atherosclerotic disease model and in human patients, opens a new potential immunometabolic therapeutic target.²¹⁵

Moderate caloric restriction could also represent an attractive therapeutic tool targeting the immunometabolic crosstalk in HF. It could improve transcriptional reprogramming in adipose tissue through pathways regulating mitochondrial bioenergetics and anti-inflammatory responsiveness,^{218,219} possibly by a conveyed reduction of sympathetic activity.²²⁰ Caloric restriction exerts potent anti-inflammatory effects in different pathological conditions by supporting the function of memory T cells and inducing a metabolic switch from glycolysis and fatty acid synthesis to fatty acid oxidation and lipolysis,^{221,222} with mechanisms related to increased ketone bodies availability.^{27,223–226} Additional studies are warranted to evaluate the clinical role of nutritional intervention in different pathologic conditions, including HF.^{227,228}

Targeting other sources of inflammation

Other drugs than evidence-based pharmacotherapy are frequently used in patients with HF due to the frequent coexistence of multiple comorbidities. In randomized controlled trials, statins did not clearly show a prognostic benefit in patients with HFrEF.^{229,230} A possible selection bias due to the exclusion of patients with strong primary indication for statins cannot be excluded. Two large observational studies suggested a possible benefit with statins in HFrEF with an ischaemic aetiology, where they are indicated

regardless of HF,^{231,232} whereas in HFpEF the observed benefit was independent of the aetiology. It was hypothesized that in HFpEF the underlying effect on systemic inflammation or on microvascular dysfunction might be implicated (Table 1, Figure 5).²³³ Statins are mainly involved in vasculoprotective and anti-atherosclerotic effects; however, they were also shown to mediate direct effects on cardiac tissue, including cardiac endothelium, cardiomyocytes and connective tissue, as for example anti-oxidative, anti-hypertrophic and anti-fibrotic effects, among others through regulating small GTP-binding proteins.^{234–236}

White adipose tissue serves as endocrine organ which systemically secretes adipokines and cytokines.²³⁷ Adipokines are involved in metabolic processes and regulate insulin signalling, glucose uptake, FFA oxidation and other energy-producing processes; cytokines mediate regulation of inflammation and adaptive angiogenesis. Lean, healthy adipose tissue secretes anti-inflammatory adipokines and cytokines, whereas obese adipose tissue switches to an inflammatory phenotype, secreting both locally and systemically pro-inflammatory cytokines and chemo-attractants, and is characterized by immune cell infiltration.²³⁷ Obese adipose tissue inflammation negatively impacts remote organ function, including the heart.^{103,237} Interestingly, epicardial adipose tissue may be involved in chronic inflammatory disorders, especially those leading to HFpEF, by switching to an inflammatory adipocyte phenotype which, by secreting pro-inflammatory adipokines, may cause atrial and ventricular fibrosis.¹⁰⁸ Thus, epicardial adipose tissue may be a transducer of the adverse effects of systemic inflammation and metabolic disorders on the heart, and may therefore represent an interesting therapeutic target. Drugs such as statins, metformin, sodium–glucose cotransporter 2 inhibitors (SGLT2i) or mineralocorticoid receptor antagonists may inhibit epicardial adipose tissue accumulation and inflammation¹⁰⁸ (Table 1, Figure 5). GLP-1 receptor agonists mitigate inflammation in visceral fat adipocytes, exert pleiotropic anti-inflammatory properties in different tissues and may modulate the immune system, with potential beneficial effects on cardiac immunometabolism^{238,239} (Table 1, Figure 5).

Neutrophils secrete myeloperoxidase, which has been implicated in the comorbidity-inflammation-microvascular dysfunction paradigm for HFpEF, and inhibition of myeloperoxidase with mitipiperstat (AZD4831) appears promising in HFpEF.^{110,240}

Low-grade inflammation of adipose tissue promotes the ectopic accumulation of lipids in other organs, including the heart, and catecholamine-derived increased lipolytic activity in adipose tissue results in elevated plasma lipids. Elevated lipid uptake and insufficient lipid oxidation in cardiomyocytes may cause mitochondrial dysfunction and oxidative stress, and consequently cardiac apoptosis, necrosis, fibrosis and inflammation.²⁴¹ Novel experimental therapies, such as adipose triglyceride lipase inhibitors, leading to improved adipose tissue storage capacities and reduced lipolytic activity as well as to anti-apoptotic, anti-fibrotic, and anti-inflammatory actions in cardiac tissue, showed promising results on cardiac metabolism and cardiac dysfunction in experimental disease models, and may therefore represent new promising pharmacological tools to treat pathological metabolic phenotypes of HF²⁴¹ (Table 1, Figure 5).

Finally, iron is essential for cardiovascular processes, including bioenergetics, electrical activity, and programmed cell death. Iron deficiency leads to impaired immune function and metabolism, while iron overload causes oxidative stress, chronic inflammation, and increased susceptibility to infections and metabolic disorders.²⁴² Therefore, maintaining iron homeostasis is crucial for both immune health and overall metabolic balance. Specifically, several studies have shown that a dysregulated systemic iron homeostasis can determine and/or worsen HF.²⁴³ The evidence of the beneficial effects of iron-modulating therapeutics in HF demonstrate the importance of maintaining iron homeostasis in the cardiovascular system.²⁴⁴

The role of metabolic substrate availability

Traditionally, metabolic drug development for HF aimed to stimulate glucose oxidation or inhibit FFA oxidation.² An alternative approach to restoring cardiac energetics may lie in providing the heart with alternative fuel sources that can bypass the roadblocks in cardiac metabolism that occur in failing cardiomyocytes.^{2,245,246}

Ketone bodies are efficient fuel that are easily oxidized by both failing and non-failing hearts.^{225,246,247} Failing hearts have a significant capacity to further increase ketone oxidation, suggesting that strategies to increase ketone delivery to the heart may serve as a viable treatment for HF.^{225,247–249} SGLT2i have been shown to induce a state of mild ketosis and improve myocardial energetics, and it has been proposed that the beneficial effects of SGLT2i may at least partially depend on enhanced myocardial ketone oxidation.^{250–252} Moreover, ketone treatment has also been shown to ameliorate cardiac dysfunction and restore myocardial energetics in small and large animal models of HFpEF and HFrEF.^{225,249,253} Short-term ketone infusion or ingestion strategies also resulted in acute improvements in cardiac function in healthy volunteers and patients with HF.²²⁵ In a preliminary small-sized randomized controlled trial, short-term treatment with ketone ester improved rest and exercise haemodynamics in HFrEF.²⁵⁴ Multiple clinical trials are currently ongoing to determine the effects of acute and chronic ketone treatment strategies on clinical endpoints in HF.²²⁵

Short-chain fatty acids (SCFA) are another promising alternative energy source for the failing heart due to their ability to bypass the down-regulation of the CPT1 shuttle.²⁵⁵ Supplementing isolated perfused hearts with the SCFA butyrate resulted in robust improvements in cardiac performance, surpassing the benefits of ketone bodies in this model.²⁴⁷ The potential of butyrate as a cardiac fuel provides a strong rationale for exploring SCFA as metabolic treatments for HF, although it may be challenging to reach sufficient circulating concentrations to serve as fuels.

Ketone bodies and SCFAs have also been shown to modulate cellular signalling and suppress inflammation through a variety of pathways. Ketone bodies inhibit NLRP3 inflammasome formation and reduced pro-inflammatory cytokine-triggered mitochondrial dysfunction and fibrosis in a murine model of HFpEF.²⁵³ β -hydroxybutyrate also reduces NLRP3 activity in macrophages isolated from diabetic patients.²⁵⁶ SCFAs, in particular butyrate,

share many of these anti-inflammatory and signalling properties with ketone bodies, but appear to do so at low concentrations.^{245,257} This suggests that ketone bodies and SCFAs may offer cardioprotective benefits beyond fuelling the heart by reducing inflammation and fibrosis.

Benefits of exercise

Physical inactivity is linked to the development of many chronic diseases through a chronic low-grade inflammation state.²⁵⁸ In chronic HF, exercise deprivation has been specifically related to further deterioration of left ventricular remodelling.²⁵⁹ Exercise elicits a strong anti-inflammatory response and systemic effects on immune function, inflammation, and host metabolism independently of weight loss and can be a useful non-pharmacologic strategy to counteract low-grade inflammation conditions, including HF.²⁶⁰ Physical exercise inhibits the inflammatory response by various molecular mechanisms.^{261,262} Accumulating data from the field of exercise immunology indicate that exercise affects immune function via cellular metabolism.²⁶³ Apart from its role on skeletal and cardiac muscle metabolism, exercise interferes with the regulation and function of leucocyte metabolism.²⁶⁴ T cells and macrophages show high sensitivity to changes in their metabolic environment, which indirectly or directly affects their central functions.²⁶⁵ Understanding the interactions between the level of physical activity and the metabolic status of immune cells could be helpful to target the dysregulated immune system in primary and secondary prevention of HF. In these contexts, future research shall eventually dissect out the differential effects of acute exercise and chronic exercise training on immunometabolism, and potential differences in the effects of exercise between immune cell types and subsets.²⁶⁶

Conclusions

Heart failure is an ongoing epidemic. Despite remarkable therapeutic progress, hospitalizations for HF remain both frequent and costly. In this context, the complete understanding of alternative mechanisms of disease progression may allow a different and more complete phenotyping of the disease and the identification of novel therapeutic targets. Persistent latent myocardial inflammation in HF probably represents an attempt to restore tissue homeostasis in the failing heart, where cardiomyocytes and immune cells undergo metabolic reprogramming, which allow them to deal with decreased availability of nutrients and oxygen. This status can trigger a metabolic crosstalk between immune cells and cardiomyocytes which can perpetuate the maladaptive remodelling of the heart. Therefore, the interrelation between immune activation and metabolism is gaining recognition as a potential therapeutic framework. The traditional cornerstones of HF treatment are mainly directed at counteracting the neurohormonal activation induced by haemodynamic incompetence and the abnormal metabolic and immune adaptations can be mitigated indirectly as a consequence of reverse remodelling and improved mechanical function. Nevertheless, chronic inflammatory/immune activation remains associated with the progression of HF and indicates that this dysregulated immunologic state is not merely the result of the overall HF

process but an active player which could be tackled with specific treatments. On the other hand, improving the metabolic efficiency of the failing circulatory system might itself alleviate the pathological inflammation breaking the vicious circle that contributes to the progression of the syndrome. We are still far from completely understanding the shape and extent of cardiac and systemic metabolic alterations in HF impacting on immune function. However, it is becoming increasingly clear that the investigation of this conundrum—metabolism and immunity—in HF, is emerging as a priority research target for HF pathophysiology and therapeutic strategies. On these premises, future studies addressing novel HF treatments should attempt to evaluate the potential therapeutic role of direct metabolic and immunological crosstalk modulation.

Conflict of interest: G.F. has received honoraria for lectures by Servier international. D.S. has received honoraria for lectures and fees for expert advice from AstraZeneca, Novo Nordisk, Novartis, Janssen, Dompè, Gossamer Bio and Merck; all unrelated to the current article. A.B.G. reports personal fees from/or participated in boards for Abbott, Amgen, AstraZeneca, Bayer, Boehringer Ingelheim, Cytokinetics, Edwards, Medtronic, Novartis, Recor, Servier and Bayer, Roche Diagnostics, Vifor during the conduct of the study. O.C. reports Servier support for ESC Congress 2022 and 2023. S.H. receives personal fees for scientific pre-clinical advice from AstraZeneca, CSL Behring and Ribocure; research contract with AstraZeneca and CSL Behring. P.S.J. reports speaker fees from AstraZeneca, Novartis, Alkem Metabolics, ProAdWise Communications, Sun Pharmaceuticals, Intas Pharmaceuticals; advisory board fees from AstraZeneca, Boehringer Ingelheim, Novartis; research funding from AstraZeneca, Boehringer Ingelheim, Analog Devices Inc; P.S.J.'s employer the University of Glasgow has also been remunerated for clinical trial work from AstraZeneca, Bayer AG, Novartis, and Novo Nordisk; Director, Global Clinical Trial Partners. B.S.L. reports consulting fees from Janssen Research and Development and from Idorsia. G.D.L. receives research support from Boehringer Ingelheim and Applied Therapeutics, and is a shareholder in Metabolic Modulators Research Ltd. L.H.L. is supported by Karolinska Institutet, the Swedish Research Council [grant 523-2014-2336], the Swedish Heart Lung Foundation [grants 20190310, 20220540], and the Stockholm County Council [grants 20190525 and FoUJ-974 688]; relationships with industry: grants, consulting, honoraria: Alleviant, AstraZeneca, Bayer, Biopeutics, Boehringer Ingelheim, Edwards, Merck/MSD, Novartis, Novo Nordisk, Owkin, Pharmacosmos; stock ownership: AnaCardio. M.P. has received personal fees from Abbott Vascular, AstraZeneca, Boehringer Ingelheim, Novartis, Roche Diagnostics and Vifor Pharma. C.G.T. is supported by two grants from the Italian Ministry of Health (PNRR-MAD-2022-12376632 and RF-2016-02362988), and reports honoraria or consultation fees from VivaLyfe, Univers Formazione, Solaris, Summeet, Astra Zeneca, Myocardial Solutions, Medtronic; funding from Amgen and MSD, and is listed as an inventor of two patents related to heart failure, outside the submitted work. P.v.d.M. reports that UMCG, his employer, received consultancy fees and/or grants from Novartis, Pharmacosmos, Vifor Pharma, AstraZeneca, Pfizer, Pharma Nord, BridgeBio, Novo Nordisk, Daiichi Sankyo, Boehringer Ingelheim and Ionis. S.V.L. was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, SFB-1470-A07). B.D.W. has received consulting fees from Boehringer Ingelheim, AstraZeneca and Novartis as well as research fees from Siemens. M.M. has received consulting honoraria from Abbott structural, AstraZeneca, Bayer, Edwards Lifesciences, Pharmacosmos, Roche Diagnostics and Novo Nordisk in the last 3 years. G.S. reports grants and personal fees from Vifor, Boehringer Ingelheim, AstraZeneca, Novartis, Cytokinetics, Pharmacosmos, personal fees from

Servier, Medtronic, Abbott, TEVA, INTAS, Edwards Lifesciences, grants from Boston Scientific, Merck, outside the submitted work. All other authors have nothing to disclose.

References

- O'Neill LA, Kishton RJ, Rathmell J. A guide to immunometabolism for immunologists. *Nat Rev Immunol* 2016;**16**:553–565. <https://doi.org/10.1038/nri.2016.70>
- Lopaschuk GD, Karwi QG, Tian R, Wende AR, Abel ED. Cardiac energy metabolism in heart failure. *Circ Res* 2021;**128**:1487–1513. <https://doi.org/10.1161/CIRCRESAHA.121.318241>
- Bottomley PA, Panjath GS, Lai S, Hirsch GA, Wu K, Najjar SS, et al. Metabolic rates of ATP transfer through creatine kinase (CK flux) predict clinical heart failure events and death. *Sci Transl Med* 2013;**5**:215re213. <https://doi.org/10.1126/scitranslmed.3007328>
- Neubauer S. The failing heart – an engine out of fuel. *N Engl J Med* 2007;**356**:1140–1151. <https://doi.org/10.1056/NEJMra063052>
- Neubauer S, Horn M, Cramer M, Harre K, Newell JB, Peters W, et al. Myocardial phosphocreatine-to-ATP ratio is a predictor of mortality in patients with dilated cardiomyopathy. *Circulation* 1997;**96**:2190–2196. <https://doi.org/10.1161/01.cir.96.7.2190>
- Ingwall JS, Weiss RG. Is the failing heart energy starved? On using chemical energy to support cardiac function. *Circ Res* 2004;**95**:135–145. <https://doi.org/10.1161/01.RES.0000137170.41939.d9>
- Zhou B, Tian R. Mitochondrial dysfunction in pathophysiology of heart failure. *J Clin Invest* 2018;**128**:3716–3726. <https://doi.org/10.1172/JCI120849>
- Knowlton AA, Chen L, Malik ZA. Heart failure and mitochondrial dysfunction: The role of mitochondrial fission/fusion abnormalities and new therapeutic strategies. *J Cardiovasc Pharmacol* 2014;**63**:196–206. <https://doi.org/10.1097/01.fjc.0000432861.55968.a6>
- Kumar AA, Kelly DP, Chirinos JA. Mitochondrial dysfunction in heart failure with preserved ejection fraction. *Circulation* 2019;**139**:1435–1450. <https://doi.org/10.1161/CIRCULATIONAHA.118.036259>
- Song M, Mihara K, Chen Y, Scorrano L, Dorn GW 2nd. Mitochondrial fission and fusion factors reciprocally orchestrate mitophagic culling in mouse hearts and cultured fibroblasts. *Cell Metab* 2015;**21**:273–286. <https://doi.org/10.1016/j.cmet.2014.12.011>
- Keceli G, Gupta A, Sourdon J, Gabr R, Schar M, Dey S, et al. Mitochondrial creatine kinase attenuates pathologic remodeling in heart failure. *Circ Res* 2022;**130**:741–759. <https://doi.org/10.1161/CIRCRESAHA.121.319648>
- Zhang L, Jaswal JS, Ussher JR, Sankaralingam S, Wagg C, Zaugg M, et al. Cardiac insulin-resistance and decreased mitochondrial energy production precede the development of systolic heart failure after pressure-overload hypertrophy. *Circ Heart Fail* 2013;**6**:1039–1048. <https://doi.org/10.1161/CIRCHEARTFAILURE.112.000228>
- Zhabiyev P, Gandhi M, Mori J, Basu R, Kassiri Z, Clanachan A, et al. Pressure-overload-induced heart failure induces a selective reduction in glucose oxidation at physiological afterload. *Cardiovasc Res* 2013;**97**:676–685. <https://doi.org/10.1093/cvr/cvs424>
- Schroeder MA, Lau AZ, Chen AP, Gu Y, Nagendran J, Barry J, et al. Hyperpolarized ¹³C magnetic resonance reveals early- and late-onset changes to in vivo pyruvate metabolism in the failing heart. *Eur J Heart Fail* 2013;**15**:130–140. <https://doi.org/10.1093/eurjhf/hfs192>
- Fernandez-Caggiano M, Kamygina A, Francois AA, Prysazhna O, Eykyn TR, Krasemann S, et al. Mitochondrial pyruvate carrier abundance mediates pathological cardiac hypertrophy. *Nat Metab* 2020;**2**:1223–1231. <https://doi.org/10.1038/s42255-020-00276-5>
- Thakker GD, Frangogiannis NG, Zymek PT, Sharma S, Raya JL, Barger PM, et al. Increased myocardial susceptibility to repetitive ischemia with high-fat diet-induced obesity. *Obesity (Silver Spring)* 2008;**16**:2593–2600. <https://doi.org/10.1038/oby.2008.414>
- Holzner KM, Marmarstein JT, Madden EJ, Efimov IR. Diet-induced obesity promotes altered remodeling and exacerbated cardiac hypertrophy following pressure overload. *Physiol Rep* 2015;**3**:e12489. <https://doi.org/10.14814/phy2.12489>
- Mori J, Basu R, McLean BA, Das SK, Zhang L, Patel VB, et al. Agonist-induced hypertrophy and diastolic dysfunction are associated with selective reduction in glucose oxidation: A metabolic contribution to heart failure with normal ejection fraction. *Circ Heart Fail* 2012;**5**:493–503. <https://doi.org/10.1161/CIRCHEARTFAILURE.112.966705>
- Mori J, Alrob OA, Wagg CS, Harris RA, Lopaschuk GD, Oudit GY. ANG II causes insulin resistance and induces cardiac metabolic switch and inefficiency: A critical role of PDK4. *Am J Physiol Heart Circ Physiol* 2013;**304**:H1103–H1113. <https://doi.org/10.1152/ajpheart.00636.2012>
- Christe ME, Rodgers RL. Cardiac glucose and fatty acid oxidation in the streptozotocin-induced diabetic spontaneously hypertensive rat. *Hypertension* 1995;**25**:235–241. <https://doi.org/10.1161/01.hyp.25.2.235>
- Sankaralingam S, Abo Alrob O, Zhang L, Jaswal JS, Wagg CS, Fukushima A, et al. Lowering body weight in obese mice with diastolic heart failure improves cardiac insulin sensitivity and function: Implications for the obesity paradox. *Diabetes* 2015;**64**:1643–1657. <https://doi.org/10.2337/db14-1050>
- Lopaschuk GD, Folmes CD, Stanley WC. Cardiac energy metabolism in obesity. *Circ Res* 2007;**101**:335–347. <https://doi.org/10.1161/CIRCRESAHA.107.150417>
- Murashige D, Jang C, Neinast M, Edwards JJ, Cowan A, Hyman MC, et al. Comprehensive quantification of fuel use by the failing and nonfailing human heart. *Science* 2020;**370**:364–368. <https://doi.org/10.1126/science.abc8861>
- Mori J, Patel VB, Abo Alrob O, Basu R, Altamimi T, Desaulniers J, et al. Angiotensin 1-7 ameliorates diabetic cardiomyopathy and diastolic dysfunction in db/db mice by reducing lipotoxicity and inflammation. *Circ Heart Fail* 2014;**7**:327–339. <https://doi.org/10.1161/CIRCHEARTFAILURE.113.000672>
- Badolia R, Ramadurai DKA, Abel ED, Ferrin P, Taleb I, Shankar TS, et al. The role of nonglycolytic glucose metabolism in myocardial recovery upon mechanical unloading and circulatory support in chronic heart failure. *Circulation* 2020;**142**:259–274. <https://doi.org/10.1161/CIRCULATIONAHA.119.044452>
- McCommis KS, Kovacs A, Weinheimer CJ, Shew TM, Koves TR, Ilkayeva OR, et al. Nutritional modulation of heart failure in mitochondrial pyruvate carrier-deficient mice. *Nat Metab* 2020;**2**:1232–1247. <https://doi.org/10.1038/s42255-020-00296-1>
- Zhang H, Tang K, Ma J, Zhou L, Liu J, Zeng L, et al. Ketogenesis-generated beta-hydroxybutyrate is an epigenetic regulator of CD8⁺ T-cell memory development. *Nat Cell Biol* 2020;**22**:18–25. <https://doi.org/10.1038/s41556-019-0440-0>
- Barger PM, Kelly DP. Fatty acid utilization in the hypertrophied and failing heart: Molecular regulatory mechanisms. *Am J Med Sci* 1999;**318**:36–42. <https://doi.org/10.1097/0000441-199907000-00006>
- Perrone-Filardi P, Bacharach SL, Dilsizian V, Panza JA, Maurea S, Bonow RO. Regional systolic function, myocardial blood flow and glucose uptake at rest in hypertrophic cardiomyopathy. *Am J Cardiol* 1993;**72**:199–204. [https://doi.org/10.1016/0002-9149\(93\)90160-e](https://doi.org/10.1016/0002-9149(93)90160-e)
- Kato T, Niizuma S, Inuzuka Y, Kawashima T, Okuda J, Tamaki Y, et al. Analysis of metabolic remodeling in compensated left ventricular hypertrophy and heart failure. *Circ Heart Fail* 2010;**3**:420–430. <https://doi.org/10.1161/CIRCHEARTFAILURE.109.888479>
- Davila-Roman VG, Vedala G, Herrero P, de las Fuentes L, Rogers JG, Kelly DP, et al. Altered myocardial fatty acid and glucose metabolism in idiopathic dilated cardiomyopathy. *J Am Coll Cardiol* 2002;**40**:271–277. [https://doi.org/10.1016/s0735-1097\(02\)01967-8](https://doi.org/10.1016/s0735-1097(02)01967-8)
- Tuunanen H, Engblom E, Naum A, Nagren K, Hesse B, Airaksinen KE, et al. Free fatty acid depletion acutely decreases cardiac work and efficiency in cardiomyopathic heart failure. *Circulation* 2006;**114**:2130–2137. <https://doi.org/10.1161/CIRCULATIONAHA.106.645184>
- Neglia D, De Caterina A, Marraccini P, Natali A, Ciardetti M, Vecoli C, et al. Impaired myocardial metabolic reserve and substrate selection flexibility during stress in patients with idiopathic dilated cardiomyopathy. *Am J Physiol Heart Circ Physiol* 2007;**293**:H3270–H3278. <https://doi.org/10.1152/ajpheart.00887.2007>
- Sack MN, Rader TA, Park S, Bastin J, McCune SA, Kelly DP. Fatty acid oxidation enzyme gene expression is downregulated in the failing heart. *Circulation* 1996;**94**:2837–2842. <https://doi.org/10.1161/01.cir.94.11.2837>
- Osorio JC, Stanley WC, Linke A, Castellari M, Diep QN, Panchal AR, et al. Impaired myocardial fatty acid oxidation and reduced protein expression of retinoid X receptor- α in pacing-induced heart failure. *Circulation* 2002;**106**:606–612. <https://doi.org/10.1161/01.cir.0000023531.22727.c1>
- Heather LC, Cole MA, Lygate CA, Evans RD, Stuckey DJ, Murray AJ, et al. Fatty acid transporter levels and palmitate oxidation rate correlate with ejection fraction in the infarcted rat heart. *Cardiovasc Res* 2006;**72**:430–437. <https://doi.org/10.1016/j.cardiores.2006.08.020>
- Taylor M, Wallhaus TR, Degradó TR, Russell DC, Stanko P, Nickles RJ, et al. An evaluation of myocardial fatty acid and glucose uptake using PET with [¹⁸F]fluoro-6-thia-heptadecanoic acid and [¹⁸F]FDG in patients with congestive heart failure. *J Nucl Med* 2001;**42**:55–62.
- Funada J, Betts TR, Hodson L, Humphreys SM, Timperley J, Frayn KN, et al. Substrate utilization by the failing human heart by direct quantification using arterio-venous blood sampling. *PLoS One* 2009;**4**:e7533. <https://doi.org/10.1371/journal.pone.0007533>
- Voros G, Ector J, Garweg C, Droogne W, Van Cleemput J, Peersman N, et al. Increased cardiac uptake of ketone bodies and free fatty acids in human

- heart failure and hypertrophic left ventricular remodeling. *Circ Heart Fail* 2018;**11**:e004953. <https://doi.org/10.1161/CIRCHEARTFAILURE.118.004953>
40. Peterson LR, Herrero P, Schechtman KB, Racette SB, Waggoner AD, Kisrieva-Ware Z, et al. Effect of obesity and insulin resistance on myocardial substrate metabolism and efficiency in young women. *Circulation* 2004;**109**:2191–2196. <https://doi.org/10.1161/01.CIR.0000127959.28627.F8>
 41. Rijzewijk LJ, van der Meer RW, Lamb HJ, de Jong HW, Lubberink M, Romijn JA, et al. Altered myocardial substrate metabolism and decreased diastolic function in nonischemic human diabetic cardiomyopathy: Studies with cardiac positron emission tomography and magnetic resonance imaging. *J Am Coll Cardiol* 2009;**54**:1524–1532. <https://doi.org/10.1016/j.jacc.2009.04.074>
 42. Barouch LA, Berkowitz DE, Harrison RW, O'Donnell CP, Hare JM. Disruption of leptin signaling contributes to cardiac hypertrophy independently of body weight in mice. *Circulation* 2003;**108**:754–759. <https://doi.org/10.1161/01.CIR.0000083716.82622.FD>
 43. Christoffersen C, Bollano E, Lindegaard ML, Bartels ED, Goetze JP, Andersen CB, et al. Cardiac lipid accumulation associated with diastolic dysfunction in obese mice. *Endocrinology* 2003;**144**:3483–3490. <https://doi.org/10.1210/en.2003-0242>
 44. Horton JL, Davidson MT, Kurishima C, Vega RB, Powers JC, Matsuura TR, et al. The failing heart utilizes 3-hydroxybutyrate as a metabolic stress defense. *JCI Insight* 2019;**4**:e124079. <https://doi.org/10.1172/jci.insight.124079>
 45. Mann DL. Innate immunity and the failing heart: The cytokine hypothesis revisited. *Circ Res* 2015;**116**:1254–1268. <https://doi.org/10.1161/CIRCRESAHA.116.302317>
 46. Torre-Amione G, Kapadia S, Lee J, Durand JB, Bies RD, Young JB, et al. Tumor necrosis factor- α and tumor necrosis factor receptors in the failing human heart. *Circulation* 1996;**93**:704–711. <https://doi.org/10.1161/01.cir.93.4.704>
 47. Zhang Y, Bauersachs J, Langer HF. Immune mechanisms in heart failure. *Eur J Heart Fail* 2017;**19**:1379–1389. <https://doi.org/10.1002/ehf.942>
 48. Adamo L, Rocha-Resende C, Prabhu SD, Mann DL. Reappraising the role of inflammation in heart failure. *Nat Rev Cardiol* 2020;**17**:269–285. <https://doi.org/10.1038/s41569-019-0315-x>
 49. Medzhitov R. Origin and physiological roles of inflammation. *Nature* 2008;**454**:428–435. <https://doi.org/10.1038/nature07201>
 50. Mouton AJ, Hall JE. Novel roles of immunometabolism and nonmyocyte metabolism in cardiac remodeling and injury. *Am J Physiol Regul Integr Comp Physiol* 2020;**319**:R476–R484. <https://doi.org/10.1152/ajpregu.00188.2020>
 51. Marelli-Berg FM, Aksentijevic D. Immunometabolic cross-talk in the inflamed heart. *Cell Stress* 2019;**3**:240–266. <https://doi.org/10.15698/cst2019.08.194>
 52. Steffens S, Van Linthout S, Sluijter JPG, Tocchetti CG, Thum T, Madonna R. Stimulating pro-reparative immune responses to prevent adverse cardiac remodeling: Consensus document from the joint 2019 meeting of the ESC Working Groups of cellular biology of the heart and myocardial function. *Cardiovasc Res* 2020;**116**:1850–1862. <https://doi.org/10.1093/cvr/cvaa137>
 53. Norata GD, Caligiuri G, Chavakis T, Matarese G, Netea MG, Nicoletti A, et al. The cellular and molecular basis of translational immunometabolism. *Immunity* 2015;**43**:421–434. <https://doi.org/10.1016/j.immuni.2015.08.023>
 54. Zhang S, Weinberg S, DeBerge M, Gainullina A, Schipma M, Kinchen JM, et al. Efferocytosis fuels requirements of fatty acid oxidation and the electron transport chain to polarize macrophages for tissue repair. *Cell Metab* 2019;**29**:443–456.e5. <https://doi.org/10.1016/j.cmet.2018.12.004>
 55. Shi LZ, Wang R, Huang G, Vogel P, Neale G, Green DR, et al. HIF1 α -dependent glycolytic pathway orchestrates a metabolic checkpoint for the differentiation of TH17 and Treg cells. *J Exp Med* 2011;**208**:1367–1376. <https://doi.org/10.1084/jem.20110278>
 56. Wu J, Sun P, Chen Q, Sun Y, Shi M, Mang G, et al. Metabolic reprogramming orchestrates CD4⁺ T-cell immunological status and restores cardiac dysfunction in autoimmune induced-dilated cardiomyopathy mice. *J Mol Cell Cardiol* 2019;**135**:134–148. <https://doi.org/10.1016/j.yjmcc.2019.08.002>
 57. Dehghan M, Mente A, Zhang X, Swaminathan S, Li W, Mohan V, et al.; Prospective Urban Rural Epidemiology (PURE) Study Investigators. Associations of fats and carbohydrate intake with cardiovascular disease and mortality in 18 countries from five continents (PURE): A prospective cohort study. *Lancet* 2017;**390**(10107):2050–2062. [https://doi.org/10.1016/S0140-6736\(17\)32252-3](https://doi.org/10.1016/S0140-6736(17)32252-3)
 58. Brandhorst S, Longo VD. Dietary restrictions and nutrition in the prevention and treatment of cardiovascular disease. *Circ Res* 2019;**124**:952–965. <https://doi.org/10.1161/CIRCRESAHA.118.313352>
 59. Kitzman DW, Brubaker P, Morgan T, Haykowsky M, Hundley G, Kraus WE, et al. Effect of caloric restriction or aerobic exercise training on peak oxygen consumption and quality of life in obese older patients with heart failure with preserved ejection fraction: A randomized clinical trial. *JAMA* 2016;**315**:36–46. <https://doi.org/10.1001/jama.2015.17346>
 60. Brubaker PH, Nicklas BJ, Houston DK, Hundley WG, Chen H, Molina AJA, et al. A randomized, controlled trial of resistance training added to caloric restriction plus aerobic exercise training in obese heart failure with preserved ejection fraction. *Circ Heart Fail* 2023;**16**:e010161. <https://doi.org/10.1161/CIRCHEARTFAILURE.122.010161>
 61. Janssen H, Kahles F, Liu D, Downey J, Koekkoek LL, Roudko V, et al. Monocytes re-enter the bone marrow during fasting and alter the host response to infection. *Immunity* 2023;**56**:783–796.e7. <https://doi.org/10.1016/j.immuni.2023.01.024>
 62. Tschope C, Ammirati E, Bozkurt B, Caforio ALP, Cooper LT, Felix SB, et al. Myocarditis and inflammatory cardiomyopathy: Current evidence and future directions. *Nat Rev Cardiol* 2021;**18**:169–193. <https://doi.org/10.1038/s41569-020-00435-x>
 63. Vilahur G, Juan-Babot O, Pena E, Onate B, Casani L, Badimon L. Molecular and cellular mechanisms involved in cardiac remodeling after acute myocardial infarction. *J Mol Cell Cardiol* 2011;**50**:522–533. <https://doi.org/10.1016/j.yjmcc.2010.12.021>
 64. Antipenko S, Mayfield N, Jinno M, Gunzer M, Ismail MA, Hamid T, et al. Neutrophils are indispensable for adverse cardiac remodeling in heart failure. *J Mol Cell Cardiol* 2024;**189**:1–11. <https://doi.org/10.1016/j.yjmcc.2024.02.005>
 65. Bansal SS, Ismail MA, Goel M, Patel B, Hamid T, Rokosh G, et al. Activated T lymphocytes are essential drivers of pathological remodeling in ischemic heart failure. *Circ Heart Fail* 2017;**10**:e003688. <https://doi.org/10.1161/CIRCHEARTFAILURE.116.003688>
 66. Varda-Bloom N, Leor J, Ohad DG, Hasin Y, Amar M, Fixler R, et al. Cytotoxic T lymphocytes are activated following myocardial infarction and can recognize and kill healthy myocytes in vitro. *J Mol Cell Cardiol* 2000;**32**:2141–2149. <https://doi.org/10.1006/jmcc.2000.1261>
 67. Zweck E, Karschnia M, Scheiber D, Heidecke H, Dechend R, Barthuber C, et al. Receptor autoantibodies: Associations with cardiac markers, histology, and function in human non-ischaemic heart failure. *ESC Heart Fail* 2023;**10**:1258–1269. <https://doi.org/10.1002/ehf2.14293>
 68. Mo F, Luo Y, Yan Y, Li J, Lai S, Wu W. Are activated B cells involved in the process of myocardial fibrosis after acute myocardial infarction? An in vivo experiment. *BMC Cardiovasc Disord* 2021;**21**:5. <https://doi.org/10.1186/s12872-020-01775-9>
 69. Mishra S, Kass DA. Cellular and molecular pathobiology of heart failure with preserved ejection fraction. *Nat Rev Cardiol* 2021;**18**:400–423. <https://doi.org/10.1038/s41569-020-00480-6>
 70. Suetomi T, Willeford A, Brand CS, Cho Y, Ross RS, Miyamoto S, et al. Inflammation and NLRP3 inflammasome activation initiated in response to pressure overload by Ca²⁺/calmodulin-dependent protein kinase II δ signaling in cardiomyocytes are essential for adverse cardiac remodeling. *Circulation* 2018;**138**:2530–2544. <https://doi.org/10.1161/CIRCULATIONAHA.118.034621>
 71. Mann DL, McMurray JJ, Packer M, Swedberg K, Borer JS, Colucci WS, et al. Targeted anticytokine therapy in patients with chronic heart failure: Results of the Randomized Etanercept Worldwide Evaluation (RENEWAL). *Circulation* 2004;**109**:1594–1602. <https://doi.org/10.1161/01.CIR.0000124490.27666.B2>
 72. Schiattarella GG, Rodolico D, Hill JA. Metabolic inflammation in heart failure with preserved ejection fraction. *Cardiovasc Res* 2021;**117**:423–434. <https://doi.org/10.1093/cvr/cvaa217>
 73. Iacobellis G. Epicardial adipose tissue in contemporary cardiology. *Nat Rev Cardiol* 2022;**19**:593–606. <https://doi.org/10.1038/s41569-022-00679-9>
 74. van Woerden G, van Veldhuisen DJ, Westenbrink BD, de Boer RA, Rienstra M, Gorter TM. Connecting epicardial adipose tissue and heart failure with preserved ejection fraction: Mechanisms, management and modern perspectives. *Eur J Heart Fail* 2022;**24**:2238–2250. <https://doi.org/10.1002/ehf.2741>
 75. Ko HJ, Zhang Z, Jung DY, Jun JY, Ma Z, Jones KE, et al. Nutrient stress activates inflammation and reduces glucose metabolism by suppressing AMP-activated protein kinase in the heart. *Diabetes* 2009;**58**:2536–2546. <https://doi.org/10.2337/db08-1361>
 76. Paulus WJ, Tschope C. A novel paradigm for heart failure with preserved ejection fraction: Comorbidities drive myocardial dysfunction and remodeling through coronary microvascular endothelial inflammation. *J Am Coll Cardiol* 2013;**62**:263–271. <https://doi.org/10.1016/j.jacc.2013.02.092>
 77. Christ A, Gunther P, Lauterbach MAR, Duewelling P, Biswas D, Pelka K, et al. Western diet triggers NLRP3-dependent innate immune reprogramming. *Cell* 2018;**172**:162–175.e14. <https://doi.org/10.1016/j.cell.2017.12.013>
 78. Olivetti G, Melissari M, Capasso JM, Anversa P. Cardiomyopathy of the aging human heart. Myocyte loss and reactive cellular hypertrophy. *Circ Res* 1991;**68**:1560–1568. <https://doi.org/10.1161/01.res.68.6.1560>
 79. Bergmann O, Bhardwaj RD, Bernard S, Zdunek S, Barnabe-Heider F, Walsh S, et al. Evidence for cardiomyocyte renewal in humans. *Science* 2009;**324**:98–102. <https://doi.org/10.1126/science.1164680>
 80. Lainscak M, Vitale C. Biological and chronological age in heart failure: Role of immunosenescence. *J Cardiovasc Med (Hagerstown)* 2016;**17**:857–859. <https://doi.org/10.2459/JCM.0000000000000448>

81. Nymo SH, Hulthe J, Ueland T, McMurray J, Wikstrand J, Askevold ET, et al. Inflammatory cytokines in chronic heart failure: Interleukin-8 is associated with adverse outcome. Results from CORONA. *Eur J Heart Fail* 2014;**16**:68–75. <https://doi.org/10.1093/eurjhf/hft125>
82. Walford RL. The immunologic theory of aging. *Gerontologist* 1964;**4**:195–197. <https://doi.org/10.1093/geront/4.4.195>
83. Seeger T, Haffez F, Fischer A, Koehl U, Leistner DM, Seeger FH, et al. Immunosenescence-associated microRNAs in age and heart failure. *Eur J Heart Fail* 2013;**15**:385–393. <https://doi.org/10.1093/eurjhf/hfs184>
84. Franceschi C, Bonafe M, Valensin S, Olivieri F, De Luca M, Ottaviani E, et al. Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann NY Acad Sci* 2000;**908**:244–254. <https://doi.org/10.1111/j.1749-6632.2000.tb06651.x>
85. Barcena ML, Aslam M, Pozdniakova S, Norman K, Ladilov Y. Cardiovascular inflammaging: Mechanisms and translational aspects. *Cells* 2022;**11**:1010. <https://doi.org/10.3390/cells11061010>
86. Bakula D, Scheibye-Knudsen M. Mitophagy: Mitophagy in aging and disease. *Front Cell Dev Biol* 2020;**8**:239. <https://doi.org/10.3389/fcell.2020.00239>
87. Chen G, Kroemer G, Kepp O. Mitophagy: An emerging role in aging and age-associated diseases. *Front Cell Dev Biol* 2020;**8**:200. <https://doi.org/10.3389/fcell.2020.00200>
88. Harman D. Aging: A theory based on free radical and radiation chemistry. *J Gerontol* 1956;**11**:298–300. <https://doi.org/10.1093/geronj/11.3.298>
89. Pinti M, Cevenini E, Nasi M, De Biasi S, Salvioli S, Monti D, et al. Circulating mitochondrial DNA increases with age and is a familial trait: Implications for “inflamm-aging”. *Eur J Immunol* 2014;**44**:1552–1562. <https://doi.org/10.1002/eji.201343921>
90. Dela Cruz CS, Kang MJ. Mitochondrial dysfunction and damage associated molecular patterns (DAMPs) in chronic inflammatory diseases. *Mitochondrion* 2018;**41**:37–44. <https://doi.org/10.1016/j.mito.2017.12.001>
91. Lu L, Guo J, Hua Y, Huang K, Magaye R, Cornell J, et al. Cardiac fibrosis in the ageing heart: Contributors and mechanisms. *Clin Exp Pharmacol Physiol* 2017;**44**:55–63. <https://doi.org/10.1111/1440-1681.12753>
92. Madonna R, Doria V, Minnucchi I, Pucci A, Pierdomenico DS, De Caterina R. Empagliflozin reduces the senescence of cardiac stromal cells and improves cardiac function in a murine model of diabetes. *J Cell Mol Med* 2020;**24**:12331–12340. <https://doi.org/10.1111/jcmm.15699>
93. Kulkarni AS, Gubbi S, Barzilai N. Benefits of metformin in attenuating the hallmarks of aging. *Cell Metab* 2020;**32**:15–30. <https://doi.org/10.1016/j.cmet.2020.04.001>
94. Park SJ, Ahmad F, Philp A, Baar K, Williams T, Luo H, et al. Resveratrol ameliorates aging-related metabolic phenotypes by inhibiting cAMP phosphodiesterases. *Cell* 2012;**148**:421–433. <https://doi.org/10.1016/j.cell.2012.01.017>
95. Ochoa JJ, Quiles JL, Huertas JR, Mataix J. Coenzyme Q10 protects from aging-related oxidative stress and improves mitochondrial function in heart of rats fed a polyunsaturated fatty acid (PUFA)-rich diet. *J Gerontol A Biol Sci Med Sci* 2005;**60**:970–975. <https://doi.org/10.1093/gerona/60.8.970>
96. Nilsson MI, Bourgeois JM, Nederveen JP, Leite MR, Hettlinga BP, Bujak AL, et al. Lifelong aerobic exercise protects against inflammaging and cancer. *PLoS One* 2019;**14**:e0210863. <https://doi.org/10.1371/journal.pone.0210863>
97. Bautmans I, Salimans L, Njemi R, Beyer I, Lietsen S, Liberman K. The effects of exercise interventions on the inflammatory profile of older adults: A systematic review of the recent literature. *Exp Gerontol* 2021;**146**:11236. <https://doi.org/10.1016/j.exger.2021.11236>
98. Jaiswal S, Fontanillas P, Flannick J, Manning A, Grauman PV, Mar BG, et al. Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med* 2014;**371**:2488–2498. <https://doi.org/10.1056/NEJMoa1408617>
99. Wang Y, Sano S, Yura Y, Ke Z, Sano M, Oshima K, et al. Tet2-mediated clonal hematopoiesis in nonconditioned mice accelerates age-associated cardiac dysfunction. *JCI Insight* 2020;**5**:e135204. <https://doi.org/10.1172/jci.insight.135204>
100. Schiattarella GG, Alcaide P, Condorelli G, Gillette TG, Heymans S, Jones EAV, et al. Immunometabolic mechanisms of heart failure with preserved ejection fraction. *Nat Cardiovasc Res* 2022;**1**:211–222. <https://doi.org/10.1038/s44161-022-00032-w>
101. Karlsson T, Rask-Andersen M, Pan G, Högglund J, Wadelius C, Ek WE, et al. Contribution of genetics to visceral adiposity and its relation to cardiovascular and metabolic disease. *Nat Med* 2019;**25**:1390–1395. <https://doi.org/10.1038/s41591-019-0563-7>
102. Powell-Wiley TM, Poirier P, Burke LE, Despres JP, Gordon-Larsen P, Lavie CJ, et al.; American Heart Association Council on Lifestyle and Cardiometabolic Health; Council on Cardiovascular and Stroke Nursing; Council on Clinical Cardiology; Council on Epidemiology and Prevention; and Stroke Council. Obesity and cardiovascular disease: A scientific statement from the American Heart Association. *Circulation* 2021;**143**:e984–e1010. <https://doi.org/10.1161/CIR.0000000000000973>
103. Oikonomou EK, Antoniadou C. The role of adipose tissue in cardiovascular health and disease. *Nat Rev Cardiol* 2019;**16**:83–99. <https://doi.org/10.1038/s41569-018-0097-6>
104. Hotamisligil GS. Foundations of immunometabolism and implications for metabolic health and disease. *Immunity* 2017;**47**:406–420. <https://doi.org/10.1016/j.immuni.2017.08.009>
105. Osborn O, Olefsky JM. The cellular and signaling networks linking the immune system and metabolism in disease. *Nat Med* 2012;**18**:363–374. <https://doi.org/10.1038/nm.2627>
106. McNelis JC, Olefsky JM. Macrophages, immunity, and metabolic disease. *Immunity* 2014;**41**:36–48. <https://doi.org/10.1016/j.immuni.2014.05.010>
107. Saltiel AR, Olefsky JM. Inflammatory mechanisms linking obesity and metabolic disease. *J Clin Invest* 2017;**127**:1–4. <https://doi.org/10.1172/JCI92035>
108. Packer M. Epicardial adipose tissue may mediate deleterious effects of obesity and inflammation on the myocardium. *J Am Coll Cardiol* 2018;**71**:2360–2372. <https://doi.org/10.1016/j.jacc.2018.03.509>
109. D’Amario D, Laborante R, Bianchini E, Ciliberti G, Pagliarini DA, Galli M, et al. Impact of coronary microvascular dysfunction in heart failure with preserved ejection fraction: A meta-analysis. *ESC Heart Fail* 2024;**11**:2063–2075. <https://doi.org/10.1002/ehf2.14626>
110. Michaelsson E, Lund LH, Hage C, Shah SJ, Voors AA, Saraste A, et al. Myeloperoxidase inhibition reverses biomarker profiles associated with clinical outcomes in HFpEF. *JACC Heart Fail* 2023;**11**:775–787. <https://doi.org/10.1016/j.jchf.2023.03.002>
111. Venkateshvaran A, Faxen UL, Hage C, Michaelsson E, Svedlund S, Saraste A, et al. Association of epicardial adipose tissue with proteomics, coronary flow reserve, cardiac structure and function, and quality of life in heart failure with preserved ejection fraction: Insights from the PROMIS-HFpEF study. *Eur J Heart Fail* 2022;**24**:2251–2260. <https://doi.org/10.1002/ehf.2709>
112. Sanders-van Wijk S, Tromp J, Beussink-Nelson L, Hage C, Svedlund S, Saraste A, et al. Proteomic evaluation of the comorbidity-inflammation paradigm in heart failure with preserved ejection fraction: Results from the PROMIS-HFpEF study. *Circulation* 2020;**142**:2029–2044. <https://doi.org/10.1161/CIRCULATIONAHA.120.045810>
113. Shah SJ, Lam CSP, Svedlund S, Saraste A, Hage C, Tan RS, et al. Prevalence and correlates of coronary microvascular dysfunction in heart failure with preserved ejection fraction: PROMIS-HFpEF. *Eur Heart J* 2018;**39**:3439–3450. <https://doi.org/10.1093/eurheartj/ehy531>
114. Harmancey R, Wilson CR, Taegtmeyer H. Adaptation and maladaptation of the heart in obesity. *Hypertension* 2008;**52**:181–187. <https://doi.org/10.1161/HYPERTENSIONAHA.108.110031>
115. Mathias S, Pena LA, Kolesnick RN. Signal transduction of stress via ceramide. *Biochem J* 1998;**335**:465–480. <https://doi.org/10.1042/bj3350465>
116. Virtue S, Vidal-Puig A. Adipose tissue expandability, lipotoxicity and the metabolic syndrome – an allostatic perspective. *Biochim Biophys Acta* 2010;**1801**:338–349. <https://doi.org/10.1016/j.bbalip.2009.12.006>
117. Cosentino F, Grant PJ, Aboyans V, Bailey CJ, Ceriello A, Delgado V, et al.; ESC Scientific Document Group. 2019 ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD. *Eur Heart J* 2020;**41**:255–323. <https://doi.org/10.1093/eurheartj/ehz486>
118. Phang RJ, Ritchie RH, Hausenloy DJ, Lees JG, Lim SY. Cellular interplay between cardiomyocytes and non-myocytes in diabetic cardiomyopathy. *Cardiovasc Res* 2023;**119**:668–690. <https://doi.org/10.1093/cvr/cvac049>
119. Seferovic PM, Paulus WJ. Clinical diabetic cardiomyopathy: A two-faced disease with restrictive and dilated phenotypes. *Eur Heart J* 2015;**36**:1718–1727. <https://doi.org/10.1093/eurheartj/ehv134>
120. Aon MA, Tocchetti CG, Bhatt N, Paolucci N, Cortassa S. Protective mechanisms of mitochondria and heart function in diabetes. *Antioxid Redox Signal* 2015;**22**:1563–1586. <https://doi.org/10.1089/ars.2014.6123>
121. Marwick TH, Ritchie R, Shaw JE, Kaye D. Implications of underlying mechanisms for the recognition and management of diabetic cardiomyopathy. *J Am Coll Cardiol* 2018;**71**:339–351. <https://doi.org/10.1016/j.jacc.2017.11.019>
122. Tan Y, Zhang Z, Zheng C, Wintergerst KA, Keller BB, Cai L. Mechanisms of diabetic cardiomyopathy and potential therapeutic strategies: Preclinical and clinical evidence. *Nat Rev Cardiol* 2020;**17**:585–607. <https://doi.org/10.1038/s41569-020-0339-2>
123. Borghetti G, von Lewinski D, Eaton DM, Sourij H, Houser SR, Wallner M. Diabetic cardiomyopathy: Current and future therapies. Beyond glycemic control. *Front Physiol* 2018;**9**:1514. <https://doi.org/10.3389/fphys.2018.01514>
124. Rao X, Zhong J, Sun Q. The heterogenic properties of monocytes/macrophages and neutrophils in inflammatory response in diabetes. *Life Sci* 2014;**116**:59–66. <https://doi.org/10.1016/j.lfs.2014.09.015>
125. Zeng C, Shi X, Zhang B, Liu H, Zhang L, Ding W, et al. The imbalance of Th17/Th1/Tregs in patients with type 2 diabetes: Relationship with metabolic

- factors and complications. *J Mol Med (Berl)* 2012;**90**:175–186. <https://doi.org/10.1007/s00109-011-0816-5>
126. Nevers T, Salvador AM, Grodecki-Pena A, Knapp A, Velazquez F, Aronovitz M, et al. Left ventricular T-cell recruitment contributes to the pathogenesis of heart failure. *Circ Heart Fail* 2015;**8**:776–787. <https://doi.org/10.1161/CIRCHEARTFAILURE.115.002225>
 127. Frati G, Schirone L, Chimenti I, Yee D, Biondi-Zoccai G, Volpe M, et al. An overview of the inflammatory signalling mechanisms in the myocardium underlying the development of diabetic cardiomyopathy. *Cardiovasc Res* 2017;**113**:378–388. <https://doi.org/10.1093/cvr/cvx011>
 128. Li J, Wang L, Wang S, Zhu H, Ye P, Xie A, et al. The Treg/Th17 imbalance in patients with idiopathic dilated cardiomyopathy. *Scand J Immunol* 2010;**71**:298–303. <https://doi.org/10.1111/j.1365-3083.2010.02374.x>
 129. Triposkiadis F, Xanthopoulos A, Bargiata A, Kitai T, Katsiki N, Farmakis D, et al. Diabetes mellitus and heart failure. *J Clin Med* 2021;**10**:3682. <https://doi.org/10.3390/jcm10163682>
 130. Sousa GR, Pober D, Galderisi A, Lv H, Yu L, Pereira AC, et al. Glycemic control, cardiac autoimmunity, and long-term risk of cardiovascular disease in type 1 diabetes mellitus. *Circulation* 2019;**139**:730–743. <https://doi.org/10.1161/CIRCULATIONAHA.118.036068>
 131. Van Linthout S, Tschope C. Inflammation – cause or consequence of heart failure or both? *Curr Heart Fail Rep* 2017;**14**:251–265. <https://doi.org/10.1007/s11897-017-0337-9>
 132. Sandek A, Bauditz J, Swidinski A, Buhner S, Weber-Eibel J, von Haehling S, et al. Altered intestinal function in patients with chronic heart failure. *J Am Coll Cardiol* 2007;**50**:1561–1569. <https://doi.org/10.1016/j.jacc.2007.07.016>
 133. Tilg H, Zmora N, Adolph TE, Elinav E. The intestinal microbiota fuelling metabolic inflammation. *Nat Rev Immunol* 2020;**20**:40–54. <https://doi.org/10.1038/s41577-019-0198-4>
 134. Nagatomo Y, Tang WH. Intersections between microbiome and heart failure: Revisiting the gut hypothesis. *J Card Fail* 2015;**21**:973–980. <https://doi.org/10.1016/j.cardfail.2015.09.017>
 135. Tang WH, Wang Z, Fan Y, Levison B, Hazen JE, Donahue LM, et al. Prognostic value of elevated levels of intestinal microbe-generated metabolite trimethylamine-N-oxide in patients with heart failure: Refining the gut hypothesis. *J Am Coll Cardiol* 2014;**64**:1908–1914. <https://doi.org/10.1016/j.jacc.2014.02.617>
 136. Krack A, Richartz BM, Gastmann A, Greim K, Lotze U, Anker SD, et al. Studies on intragastric PCO₂ at rest and during exercise as a marker of intestinal perfusion in patients with chronic heart failure. *Eur J Heart Fail* 2004;**6**:403–407. <https://doi.org/10.1016/j.ejheart.2004.03.002>
 137. Pasini E, Aquilani R, Testa C, Baiardi P, Angioletti S, Boschi F, et al. Pathogenic gut flora in patients with chronic heart failure. *JACC Heart Fail* 2016;**4**:220–227. <https://doi.org/10.1016/j.jchf.2015.10.009>
 138. Chioncel O, Ambrosy AP. Trimethylamine N-oxide and risk of heart failure progression: Marker or mediator of disease. *Eur J Heart Fail* 2019;**21**:887–890. <https://doi.org/10.1002/ehf.1409>
 139. Zheng D, Liwinski T, Elinav E. Interaction between microbiota and immunity in health and disease. *Cell Res* 2020;**30**:492–506. <https://doi.org/10.1038/s41422-020-0332-7>
 140. Wilck N, Matus MG, Kearney SM, Olesen SW, Forslund K, Bartolomaeus H, et al. Salt-responsive gut commensal modulates T_H17 axis and disease. *Nature* 2017;**551**:585–589. <https://doi.org/10.1038/nature24628>
 141. Agus A, Denizot J, Thevenot J, Martinez-Medina M, Massier S, Sauvanet P, et al. Western diet induces a shift in microbiota composition enhancing susceptibility to adherent-invasive *E. coli* infection and intestinal inflammation. *Sci Rep* 2016;**6**:19032. <https://doi.org/10.1038/srep19032>
 142. Madan S, Mehra MR. Gut dysbiosis and heart failure: Navigating the universe within. *Eur J Heart Fail* 2020;**22**:629–637. <https://doi.org/10.1002/ehf.1792>
 143. Virtue AT, McCright SJ, Wright JM, Jimenez MT, Mowle WK, Kotzin JJ, et al. The gut microbiota regulates white adipose tissue inflammation and obesity via a family of microRNAs. *Sci Transl Med* 2019;**11**:eaav1892. <https://doi.org/10.1126/scitranslmed.aav1892>
 144. Ganesh N, van der Vorst EPC, Spiesshofer J, He S, Burgmaier M, Findeisen H, et al. Gut immune cells – a novel therapeutic target for cardiovascular disease? *Front Cardiovasc Med* 2022;**9**:943214. <https://doi.org/10.3389/fcvm.2022.943214>
 145. Parker CM, Cepek KL, Russell GJ, Shaw SK, Posnett DN, Schwarting R, et al. A family of beta 7 integrins on human mucosal lymphocytes. *Proc Natl Acad Sci USA* 1992;**89**:1924–1928. <https://doi.org/10.1073/pnas.89.5.1924>
 146. He S, Kahles F, Rattik S, Nairz M, McAlpine CS, Anzai A, et al. Gut intraepithelial T cells calibrate metabolism and accelerate cardiovascular disease. *Nature* 2019;**566**:115–119. <https://doi.org/10.1038/s41586-018-0849-9>
 147. Hattori A, Kawamura I, Yamada Y, Kanamori H, Aoyama T, Ushikoshi H, et al. Elevated plasma GLP-1 levels and enhanced expression of cardiac GLP-1 receptors as markers of left ventricular systolic dysfunction: A cross-sectional study. *BMJ Open* 2013;**3**:e003201. <https://doi.org/10.1136/bmjopen-2013-003201>
 148. Ussher JR, Drucker DJ. Glucagon-like peptide 1 receptor agonists: Cardiovascular benefits and mechanisms of action. *Nat Rev Cardiol* 2023;**20**:463–474. <https://doi.org/10.1038/s41569-023-00849-3>
 149. Sattar N, Lee MMY, Kristensen SL, Branch KRH, Del Prato S, Khurmi NS, et al. Cardiovascular, mortality, and kidney outcomes with GLP-1 receptor agonists in patients with type 2 diabetes: A systematic review and meta-analysis of randomised trials. *Lancet Diabetes Endocrinol* 2021;**9**:653–662. [https://doi.org/10.1016/S2213-8587\(21\)00203-5](https://doi.org/10.1016/S2213-8587(21)00203-5)
 150. Kosiborod MN, Abildstrom SZ, Borlaug BA, Butler J, Rasmussen S, Davies M, et al. STEP-HFpEF Trial Committees and Investigators. Semaglutide in patients with heart failure with preserved ejection fraction and obesity. *N Engl J Med* 2023;**389**:1069–1084. <https://doi.org/10.1056/NEJMoa2306963>
 151. Kosiborod MN, Petrie MC, Borlaug BA, Butler J, Davies MJ, Hovingh GK, et al. STEP-HFpEF DM Trial Committees and Investigators. Semaglutide in patients with obesity-related heart failure and type 2 diabetes. *N Engl J Med* 2024;**390**:1394–1407. <https://doi.org/10.1056/NEJMoa2313917>
 152. Butler J, Shah SJ, Petrie MC, Borlaug BA, Abildstrom SZ, Davies MJ, et al. STEP-HFpEF Trial Committees and Investigators. Semaglutide versus placebo in people with obesity-related heart failure with preserved ejection fraction: A pooled analysis of the STEP-HFpEF and STEP-HFpEF DM randomised trials. *Lancet* 2024;**403**:1635–1648. [https://doi.org/10.1016/S0140-6736\(24\)00469-0](https://doi.org/10.1016/S0140-6736(24)00469-0)
 153. Gil-Cruz C, Perez-Shibayama C, De Martin A, Ronchi F, van der Borgh K, Niederer R, et al. Microbiota-derived peptide mimics drive lethal inflammatory cardiomyopathy. *Science* 2019;**366**:881–886. <https://doi.org/10.1126/science.aav3487>
 154. Rosano GMC, Stolfo D, Anderson L, Abdelhamid M, Adamo M, Bauersachs J, et al. Differences in presentation, diagnosis and management of heart failure in women. A scientific statement of the Heart Failure Association of the ESC. *Eur J Heart Fail* 2024;**26**:1669–1686. <https://doi.org/10.1002/ehf.3284>
 155. Packer M, Lam CSP, Lund LH, Maurer MS, Borlaug BA. Characterization of the inflammatory-metabolic phenotype of heart failure with a preserved ejection fraction: A hypothesis to explain influence of sex on the evolution and potential treatment of the disease. *Eur J Heart Fail* 2020;**22**:1551–1567. <https://doi.org/10.1002/ehf.1902>
 156. Murphy E. Estrogen signaling and cardiovascular disease. *Circ Res* 2011;**109**:687–696. <https://doi.org/10.1161/CIRCRESAHA.110.236687>
 157. Tamis-Holland JE. Sex and outcomes after percutaneous coronary intervention: A cause for concern for young women and those with ST-segment elevation myocardial infarction? *J Am Heart Assoc* 2017;**6**:e005739. <https://doi.org/10.1161/JAHA.117.005739>
 158. Sicklinghe AA, Korporeal SJA, den Ruijter HM, Kessler EL. Estrogen contributions to microvascular dysfunction evolving to heart failure with preserved ejection fraction. *Front Endocrinol (Lausanne)* 2019;**10**:442. <https://doi.org/10.3389/fendo.2019.00442>
 159. Lau ES, Paniagua SM, Guseh JS, Bhamhani V, Zanni MV, Courchesne P, et al. Sex differences in circulating biomarkers of cardiovascular disease. *J Am Coll Cardiol* 2019;**74**:1543–1553. <https://doi.org/10.1016/j.jacc.2019.06.077>
 160. Klein SL, Flanagan KL. Sex differences in immune responses. *Nat Rev Immunol* 2016;**16**:626–638. <https://doi.org/10.1038/nri.2016.90>
 161. Pelliccia F, Kaski JC, Crea F, Camici PG. Pathophysiology of Takotsubo syndrome. *Circulation* 2017;**135**:2426–2441. <https://doi.org/10.1161/CIRCULATIONAHA.116.027121>
 162. Bauersachs J, Konig T, van der Meer P, Petrie MC, Hilfiker-Kleiner D, Mbakem A, et al. Pathophysiology, diagnosis and management of peripartum cardiomyopathy: A position statement from the Heart Failure Association of the European Society of Cardiology Study Group on peripartum cardiomyopathy. *Eur J Heart Fail* 2019;**21**:827–843. <https://doi.org/10.1002/ehf.1493>
 163. Fliegner D, Schubert C, Penkalla A, Witt H, Kararigas G, Dworatzek E, et al. Female sex and estrogen receptor-beta attenuate cardiac remodeling and apoptosis in pressure overload. *Am J Physiol Regul Integr Comp Physiol* 2010;**298**:R1597–R1606. <https://doi.org/10.1152/ajpregu.00825.2009>
 164. Regitz-Zagrosek V. Sex and gender differences in heart failure. *Int J Heart Fail* 2020;**2**:157–181. <https://doi.org/10.36628/ijhf.2020.0004>
 165. Kararigas G, Dworatzek E, Petrov G, Summer H, Schulze TM, Baczkowski I, et al. Sex-dependent regulation of fibrosis and inflammation in human left ventricular remodelling under pressure overload. *Eur J Heart Fail* 2014;**16**:1160–1167. <https://doi.org/10.1002/ehf.171>
 166. Chen JQ, Eshete M, Alworth WL, Yager JD. Binding of MCF-7 cell mitochondrial proteins and recombinant human estrogen receptors alpha and beta to human mitochondrial DNA estrogen response elements. *J Cell Biochem* 2004;**93**:358–373. <https://doi.org/10.1002/jcb.20178>

167. Rettberg JR, Yao J, Brinton RD. Estrogen: A master regulator of bioenergetic systems in the brain and body. *Front Neuroendocrinol* 2014;**35**:8–30. <https://doi.org/10.1016/j.yfrne.2013.08.001>
168. Miller VM, Duckles SP. Vascular actions of estrogens: Functional implications. *Pharmacol Rev* 2008;**60**:210–241. <https://doi.org/10.1124/pr.107.08002>
169. Humes HD, Aaronson KD, Buffington DA, Sabbah HN, Westover AJ, Yessayan LT, et al. Translation of immunomodulatory therapy to treat chronic heart failure: Preclinical studies to first in human. *PLoS One* 2023;**18**:e0273138. <https://doi.org/10.1371/journal.pone.0273138>
170. Palsen-McDermott EM, O'Neill LAJ. Targeting immunometabolism as an anti-inflammatory strategy. *Cell Res* 2020;**30**:300–314. <https://doi.org/10.1038/s41422-020-0291-z>
171. Fragasso G, Margonato A, Spoladore R, Lopaschuk GD. Metabolic effects of cardiovascular drugs. *Trends Cardiovasc Med* 2019;**29**:176–187. <https://doi.org/10.1016/j.tcm.2018.08.001>
172. Ridker PM, Rane M. Interleukin-6 signaling and anti-Interleukin-6 therapeutics in cardiovascular disease. *Circ Res* 2021;**128**:1728–1746. <https://doi.org/10.1161/CIRCRESAHA.121.319077>
173. Olsen MB, Gregersen I, Sandanger O, Yang K, Sokolova M, Halvorsen BE, et al. Targeting the inflammasome in cardiovascular disease. *JACC Basic Transl Sci* 2022;**7**:84–98. <https://doi.org/10.1016/j.jacbs.2021.08.006>
174. Petrie M, Borlaug B, Buchholtz K, Ducharme A, Hvelplund A, Ping C, et al. HERMES: Effects of ziltivekimab versus placebo on morbidity and mortality in patients with heart failure with mildly reduced or preserved ejection fraction and systemic inflammation. *J Card Fail* 2024;**30**:126. <https://doi.org/10.1016/j.cardfail.2023.10.024>
175. Vasamsetti SB, Florentin J, Coppin E, Stiekema LCA, Zheng KH, Nisar MU, et al. Sympathetic neuronal activation triggers myeloid progenitor proliferation and differentiation. *Immunity* 2018;**49**:93–106.e7. <https://doi.org/10.1016/j.immuni.2018.05.004>
176. Badoer E. Microglia: Activation in acute and chronic inflammatory states and in response to cardiovascular dysfunction. *Int J Biochem Cell Biol* 2010;**42**:1580–1585. <https://doi.org/10.1016/j.biocel.2010.07.005>
177. Wang M, Pan W, Xu Y, Zhang J, Wan J, Jiang H. Microglia-mediated neuroinflammation: A potential target for the treatment of cardiovascular diseases. *J Inflamm Res* 2022;**15**:3083–3094. <https://doi.org/10.2147/JIR.S350109>
178. Day JL. The metabolic consequences of adrenergic blockade: A review. *Metabolism* 1975;**24**:987–996. [https://doi.org/10.1016/0026-0495\(75\)90090-6](https://doi.org/10.1016/0026-0495(75)90090-6)
179. Ahlborg G, Juhlén-Dannfelt A. Effect of beta-receptor blockade on splanchnic and muscle metabolism during prolonged exercise in men. *J Appl Physiol (1985)* 1994;**76**:1037–1042. <https://doi.org/10.1152/jappl.1994.76.3.1037>
180. Mora-Rodriguez R, Hodgkinson BJ, Byerley LO, Coyle EF. Effects of beta-adrenergic receptor stimulation and blockade on substrate metabolism during submaximal exercise. *Am J Physiol Endocrinol Metab* 2001;**280**:E752–E760. <https://doi.org/10.1152/ajpendo.2001.280.5.E752>
181. Wallhaus TR, Taylor M, DeGrado TR, Russell DC, Stanko P, Nickles RJ, et al. Myocardial free fatty acid and glucose use after carvedilol treatment in patients with congestive heart failure. *Circulation* 2001;**103**:2441–2446. <https://doi.org/10.1161/01.cir.103.20.2441>
182. Bakris GL, Fonseca V, Katholi RE, McGill JB, Messerli FH, Phillips RA, et al.; GEMINI Investigators. Metabolic effects of carvedilol vs metoprolol in patients with type 2 diabetes mellitus and hypertension: A randomized controlled trial. *JAMA* 2004;**292**:2227–2236. <https://doi.org/10.1001/jama.292.18.2227>
183. Nielsen TT, Bagge JP, Thomassen A. Improved myocardial lactate extraction after propranolol in coronary artery disease: Effected by peripheral glutamate and free fatty acid metabolism. *Br Heart J* 1986;**55**:140–147. <https://doi.org/10.1136/hrt.55.2.140>
184. Pisarenko OI, Solomatina ES, Studneva IM, Ivanov VE, Kapelko VI, Smirnov VN. Protective effect of glutamic acid on cardiac function and metabolism during cardioplegia and reperfusion. *Basic Res Cardiol* 1983;**78**:534–543. <https://doi.org/10.1007/BF01906464>
185. Spoladore R, Fragasso G, Perseghin G, De Cobelli F, Esposito A, Maranta F, et al. Beneficial effects of beta-blockers on left ventricular function and cellular energy reserve in patients with heart failure. *Fundam Clin Pharmacol* 2013;**27**:455–464. <https://doi.org/10.1111/j.1472-8206.2012.01029.x>
186. Zderic TW, Schenk S, Davidson CJ, Byerley LO, Coyle EF. Manipulation of dietary carbohydrate and muscle glycogen affects glucose uptake during exercise when fat oxidation is impaired by beta-adrenergic blockade. *Am J Physiol Endocrinol Metab* 2004;**287**:E1195–E1201. <https://doi.org/10.1152/ajpendo.00302.2004>
187. Raptis S, Rosenthal J, Welzel D, Moulouopoulos S. Effects of cardioselective and non-cardioselective beta-blockade on adrenaline-induced metabolic and cardiovascular responses in man. *Eur J Clin Pharmacol* 1981;**20**:17–22. <https://doi.org/10.1007/BF00554661>
188. Gruber-Olipitz M, Stevenson R, Olipitz W, Wagner E, Gesslbauer B, Kungl A, et al. Transcriptional pattern analysis of adrenergic immunoregulation in mice. Twelve hours norepinephrine treatment alters the expression of a set of genes involved in monocyte activation and leukocyte trafficking. *J Neuroimmunol* 2004;**155**:136–142. <https://doi.org/10.1016/j.jneuroim.2004.07.003>
189. Ohtsuka T, Hamada M, Hiasa G, Sasaki O, Suzuki M, Hara Y, et al. Effect of beta-blockers on circulating levels of inflammatory and anti-inflammatory cytokines in patients with dilated cardiomyopathy. *J Am Coll Cardiol* 2001;**37**:412–417. [https://doi.org/10.1016/s0735-1097\(00\)01121-9](https://doi.org/10.1016/s0735-1097(00)01121-9)
190. Yang SP, Ho LJ, Cheng SM, Hsu YL, Tsao TP, Chang DM, et al. Carvedilol differentially regulates cytokine production from activated human peripheral blood mononuclear cells. *Cardiovasc Drugs Ther* 2004;**18**:183–188. <https://doi.org/10.1023/B:CARD.0000033638.37830.f7>
191. Elenkov IJ, Wilder R, Chrousos G, Vizi E. The sympathetic nerve – an integrative interface between two supersystems: The brain and the immune system. *Pharmacol Rev* 2000;**52**:595–638. [https://doi.org/10.1016/S0031-6997\(24\)01470-4](https://doi.org/10.1016/S0031-6997(24)01470-4)
192. Alfieri AB, Briceno L, Fragasso G, Spoladore R, Pallosi A, Bassanelli G, et al. Differential long-term effects of carvedilol on proinflammatory and antiinflammatory cytokines, asymmetric dimethylarginine, and left ventricular function in patients with heart failure. *J Cardiovasc Pharmacol* 2008;**52**:49–54. <https://doi.org/10.1097/FJC.0b013e31817e0edd>
193. Pellieux C, Aasum E, Larsen TS, Montessuit C, Papageorgiou I, Pedrazzini T, et al. Overexpression of angiotensinogen in the myocardium induces downregulation of the fatty acid oxidation pathway. *J Mol Cell Cardiol* 2006;**41**:459–466. <https://doi.org/10.1016/j.yjmcc.2006.06.004>
194. Dai DF, Johnson S, Villarin J, Chin M, Nieves M, Chen T, et al. Mitochondrial oxidative stress mediates angiotensin II-induced cardiac hypertrophy and Gq overexpression-induced heart failure. *Circ Res* 2011;**108**:837–846. <https://doi.org/10.1161/CIRCRESAHA.110.232306>
195. Henriksen EJ, Jacob S. Modulation of metabolic control by angiotensin converting enzyme (ACE) inhibition. *J Cell Physiol* 2003;**196**:171–179. <https://doi.org/10.1002/jcp.10294>
196. Dietze GJ, Henriksen EJ. Angiotensin-converting enzyme in skeletal muscle: Sentinel of blood pressure control and glucose homeostasis. *J Renin Angiotensin Aldosterone Syst* 2008;**9**:75–88. <https://doi.org/10.3317/jraas.2008.011>
197. Jin HM, Pan Y. Angiotensin type-1 receptor blockade with losartan increases insulin sensitivity and improves glucose homeostasis in subjects with type 2 diabetes and nephropathy. *Nephrol Dial Transplant* 2007;**22**:1943–1949. <https://doi.org/10.1093/ndt/gfm049>
198. Cabassi A, Coghi P, Govoni P, Barouhiel E, Speroni E, Cavazzini S, et al. Sympathetic modulation by carvedilol and losartan reduces angiotensin II-mediated lipolysis in subcutaneous and visceral fat. *J Clin Endocrinol Metab* 2005;**90**:2888–2897. <https://doi.org/10.1210/jc.2004-1995>
199. Fillmore N, Mori J, Lopaschuk GD. Mitochondrial fatty acid oxidation alterations in heart failure, ischaemic heart disease and diabetic cardiomyopathy. *Br J Pharmacol* 2014;**171**:2080–2090. <https://doi.org/10.1111/bph.12475>
200. Awad K, Zaki MM, Mohammed M, Lewek J, Lavie CJ, Banach M; Lipid and Blood Pressure Meta-analysis Collaboration Group. Effect of the renin-angiotensin system inhibitors on inflammatory markers: A systematic review and meta-analysis of randomized controlled trials. *Mayo Clin Proc* 2022;**97**:1808–1823. <https://doi.org/10.1016/j.mayocp.2022.06.036>
201. Oosthuizen D, Sturrock ED. Exploring the impact of ACE inhibition in immunity and disease. *J Renin Angiotensin Aldosterone Syst* 2022;**2022**:9028969. <https://doi.org/10.1155/2022/9028969>
202. Paradies G, Paradies V, De Benedictis V, Ruggiero FM, Petrosillo G. Functional role of cardiolipin in mitochondrial bioenergetics. *Biochim Biophys Acta* 2014;**1837**:408–417. <https://doi.org/10.1016/j.bbabi.2013.10.006>
203. Bentinger M, Brismar K, Dallner G. The antioxidant role of coenzyme Q. *Mitochondrion* 2007;**7**:S41–S50. <https://doi.org/10.1016/j.mito.2007.02.006>
204. Sharma A, Fonarow GC, Butler J, Ezekowitz JA, Felker GM. Coenzyme Q10 and heart failure: A state-of-the-art review. *Circ Heart Fail* 2016;**9**:e002639. <https://doi.org/10.1161/CIRCHEARTFAILURE.115.002639>
205. Folkers K, Vadhavavik S, Mortensen SA. Biochemical rationale and myocardial tissue data on the effective therapy of cardiomyopathy with coenzyme Q10. *Proc Natl Acad Sci USA* 1985;**82**:901–904. <https://doi.org/10.1073/pnas.82.3.901>
206. Molyneux SL, Florkowski CM, George PM, Pilbrow AP, Frampton CM, Lever M, et al. Coenzyme Q10: An independent predictor of mortality in chronic heart failure. *J Am Coll Cardiol* 2008;**52**:1435–1441. <https://doi.org/10.1016/j.jacc.2008.07.044>
207. McMurray JJ, Dunselman P, Wedel H, Cleland JG, Lindberg M, Hjalmarsson A, et al.; CORONA Study Group. Coenzyme Q10, rosuvastatin, and clinical outcomes in heart failure: A pre-specified substudy of CORONA (controlled rosuvastatin multinational study in heart failure). *J Am Coll Cardiol* 2010;**56**:1196–1204. <https://doi.org/10.1016/j.jacc.2010.02.075>

208. Mortensen SA, Rosenfeldt F, Kumar A, Dolliner P, Filipiak KJ, Pella D, et al.; Q-SYMBIO Study Investigators. The effect of coenzyme Q10 on morbidity and mortality in chronic heart failure: Results from Q-SYMBIO: A randomized double-blind trial. *JACC Heart Fail* 2014;**2**:641–649. <https://doi.org/10.1016/j.jchf.2014.06.008>
209. Al Saadi T, Assaf Y, Farwati M, Turkmani K, Al-Mouakeh A, Shebli B, et al. Coenzyme Q10 for heart failure. *Cochrane Database Syst Rev* 2021;**2**:CD008684. <https://doi.org/10.1002/14651858.CD008684.pub3>
210. Kantor PF, Lucien A, Kozak R, Lopaschuk GD. The antianginal drug trimetazidine shifts cardiac energy metabolism from fatty acid oxidation to glucose oxidation by inhibiting mitochondrial long-chain 3-ketoacyl coenzyme A thiolase. *Circ Res* 2000;**86**:580–588. <https://doi.org/10.1161/01.res.86.5.580>
211. Fragasso G, Perseghin G, De Cobelli F, Esposito A, Pallosi A, Lattuada G, et al. Effects of metabolic modulation by trimetazidine on left ventricular function and phosphocreatine/adenosine triphosphate ratio in patients with heart failure. *Eur Heart J* 2006;**27**:942–948. <https://doi.org/10.1093/eurheartj/ehi816>
212. Fragasso G, Pallosi A, Puccetti P, Silipigni C, Rossodivita A, Pala M, et al. A randomized clinical trial of trimetazidine, a partial free fatty acid oxidation inhibitor, in patients with heart failure. *J Am Coll Cardiol* 2006;**48**:992–998. <https://doi.org/10.1016/j.jacc.2006.03.060>
213. Monti LD, Setola E, Fragasso G, Camisasca RP, Lucotti P, Galluccio E, et al. Metabolic and endothelial effects of trimetazidine on forearm skeletal muscle in patients with type 2 diabetes and ischemic cardiomyopathy. *Am J Physiol Endocrinol Metab* 2006;**290**:E54–E59. <https://doi.org/10.1152/ajpendo.00083.2005>
214. Chen J, Lai J, Yang L, Ruan G, Chaugai S, Ning Q, et al. Trimetazidine prevents macrophage-mediated septal myocardial dysfunction via activation of the histone deacetylase sirtuin 1. *Br J Pharmacol* 2016;**173**:545–561. <https://doi.org/10.1111/bph.13386>
215. Lenz M, Salzmann M, Ciotu CI, Kaun C, Krychtiuk KA, Rehberger Likoza A, et al. Pharmacologic modulation of intracellular Na⁺ concentration with ranolazine impacts inflammatory response in humans and mice. *Proc Natl Acad Sci USA* 2022;**119**:e2207020119. <https://doi.org/10.1073/pnas.2207020119>
216. McCormack JG, Barr RL, Wolff AA, Lopaschuk GD. Ranolazine stimulates glucose oxidation in normoxic, ischemic, and reperfused ischemic rat hearts. *Circulation* 1996;**93**:135–142. <https://doi.org/10.1161/01.cir.93.1.135>
217. Antzelevitch C, Belardinelli L, Zygmunt AC, Burashnikov A, Di Diego JM, Fish JM, et al. Electrophysiological effects of ranolazine, a novel antianginal agent with antiarrhythmic properties. *Circulation* 2004;**110**:904–910. <https://doi.org/10.1161/01.CIR.0000139333.83620.5D>
218. Spadaro O, Youm Y, Shchukina I, Ryu S, Sidorov S, Ravussin A, et al. Caloric restriction in humans reveals immunometabolic regulators of health span. *Science* 2022;**375**:671–677. <https://doi.org/10.1126/science.abg7292>
219. Niemann B, Chen Y, Issa H, Silber RE, Rohrbach S. Caloric restriction delays cardiac ageing in rats: Role of mitochondria. *Cardiovasc Res* 2010;**88**:267–276. <https://doi.org/10.1093/cvr/cvq273>
220. Niemann B, Li L, Simm A, Molenda N, Kockskemper J, Boening A, et al. Caloric restriction reduces sympathetic activity similar to beta-blockers but conveys additional mitochondrio-protective effects in aged myocardium. *Sci Rep* 2021;**11**:1931. <https://doi.org/10.1038/s41598-021-81438-7>
221. Collins N, Han SJ, Enamorado M, Link VM, Huang B, Moseman EA, et al. The bone marrow protects and optimizes immunological memory during dietary restriction. *Cell* 2019;**178**:1088–1101.e15. <https://doi.org/10.1016/j.cell.2019.07.049>
222. Bruss MD, Khambatta CF, Ruby MA, Aggarwal I, Hellerstein MK. Calorie restriction increases fatty acid synthesis and whole body fat oxidation rates. *Am J Physiol Endocrinol Metab* 2010;**298**:E108–E116. <https://doi.org/10.1152/ajpendo.00524.2009>
223. Adam C, Paolini L, Gueguen N, Mabileau G, Preisser L, Blanchard S, et al. Acetoacetate protects macrophages from lactic acidosis-induced mitochondrial dysfunction by metabolic reprogramming. *Nat Commun* 2021;**12**:7115. <https://doi.org/10.1038/s41467-021-27426-x>
224. Bae HR, Kim DH, Park MH, Lee B, Kim MJ, Lee EK, et al. Beta-hydroxybutyrate suppresses inflammasome formation by ameliorating endoplasmic reticulum stress via AMPK activation. *Oncotarget* 2016;**7**:66444–66454. <https://doi.org/10.18632/oncotarget.12119>
225. Matsuura TR, Puchalska P, Crawford PA, Kelly DP. Ketones and the heart: Metabolic principles and therapeutic implications. *Circ Res* 2023;**132**:882–898. <https://doi.org/10.1161/CIRCRESAHA.123.321872>
226. Shimazu T, Hirschey MD, Newman J, He W, Shirakawa K, Le Moan N, et al. Suppression of oxidative stress by beta-hydroxybutyrate, an endogenous histone deacetylase inhibitor. *Science* 2013;**339**:211–214. <https://doi.org/10.1126/science.1227166>
227. Procaccini C, de Candia P, Russo C, De Rosa G, Lepore MT, Colamattéo A, et al. Caloric restriction for the immunometabolic control of human health. *Cardiovasc Res* 2024;**119**:2787–2800. <https://doi.org/10.1093/cvr/cvad035>
228. Forte M, Rodolico D, Ameri P, Catalucci D, Chimenti C, Crotti L, et al.; Italian Society of Cardiology Working Group on Cellular, Molecular Biology of the Heart. Molecular mechanisms underlying the beneficial effects of exercise and dietary interventions in the prevention of cardiometabolic diseases. *J Cardiovasc Med (Hagerstown)* 2023;**24**:e3–e14. <https://doi.org/10.2459/JCM.0000000000001397>
229. Kjekshus J, Apetrei E, Barrios V, Bohm M, Cleland JG, Cornel JH, et al.; CORONA Group. Rosuvastatin in older patients with systolic heart failure. *N Engl J Med* 2007;**357**:2248–2261. <https://doi.org/10.1056/NEJMoa0706201>
230. Tavazzi L, Maggioni AP, Marchioli R, Barlera S, Franzosi MG, Latini R, et al.; GISSI-HF Investigators. Effect of rosuvastatin in patients with chronic heart failure (the GISSI-HF trial): A randomised, double-blind, placebo-controlled trial. *Lancet* 2008;**372**:1231–1239. [https://doi.org/10.1016/S0140-6736\(08\)61240-4](https://doi.org/10.1016/S0140-6736(08)61240-4)
231. Alehagen U, Benson L, Edner M, Dahlstrom U, Lund LH. Association between use of statins and outcomes in heart failure with reduced ejection fraction: Prospective propensity score matched cohort study of 21 864 patients in the Swedish Heart Failure Registry. *Circ Heart Fail* 2015;**8**:252–260. <https://doi.org/10.1161/CIRCHEARTFAILURE.114.001730>
232. Vrints C, Andreotti F, Koskinas KC, Rossello X, Adamo M, Ainslie J, et al.; ESC Scientific Document Group. 2024 ESC Guidelines for the management of chronic coronary syndromes. *Eur Heart J* 2024;**45**:3415–3537. <https://doi.org/10.1093/eurheartj/ehae177>
233. Alehagen U, Benson L, Edner M, Dahlstrom U, Lund LH. Association between use of statins and mortality in patients with heart failure and ejection fraction of $\geq 50\%$. *Circ Heart Fail* 2015;**8**:862–870. <https://doi.org/10.1161/CIRCHEARTFAILURE.115.002143>
234. Oesterle A, Laufs U, Liao JK. Pleiotropic effects of statins on the cardiovascular system. *Circ Res* 2017;**120**:229–243. <https://doi.org/10.1161/CIRCRESAHA.116.308537>
235. Mital S, Liao JK. Statins and the myocardium. *Semin Vasc Med* 2004;**4**:377–384. <https://doi.org/10.1055/s-2004-869594>
236. Gorabi AM, Kiaie N, Bianconi V, Pirro M, Jamalrahmani T, Sahebkar A. Statins attenuate fibrotic manifestations of cardiac tissue damage. *Curr Mol Pharmacol* 2021;**14**:782–797. <https://doi.org/10.2174/1874467214666210210123206>
237. Kawai T, Autieri MV, Scalia R. Adipose tissue inflammation and metabolic dysfunction in obesity. *Am J Physiol Cell Physiol* 2021;**320**:C375–C391. <https://doi.org/10.1152/ajpcell.00379.2020>
238. Martins FF, Marinho TS, Cardoso LEM, Barbosa-da-Silva S, Souza-Mello V, Aguilu MB, et al. Semaglutide (GLP-1 receptor agonist) stimulates browning on subcutaneous fat adipocytes and mitigates inflammation and endoplasmic reticulum stress in visceral fat adipocytes of obese mice. *Cell Biochem Funct* 2022;**40**:903–913. <https://doi.org/10.1002/cbf.3751>
239. Bendotti G, Montefusco L, Lunati ME, Usuell V, Pastore I, Lazzaroni E, et al. The anti-inflammatory and immunological properties of GLP-1 receptor agonists. *Pharmacol Res* 2022;**182**:106320. <https://doi.org/10.1016/j.phrs.2022.106320>
240. Lam CSP, Lund LH, Shah SJ, Voors AA, Erlinge D, Saraste A, et al. Myeloperoxidase inhibition in heart failure with preserved or mildly reduced ejection fraction: SATELLITE trial results. *J Card Fail* 2024;**30**:104–110. <https://doi.org/10.1016/j.cardfail.2023.04.003>
241. Smeir E, Kintscher U, Foryst-Ludwig A. Adipose tissue-heart crosstalk as a novel target for treatment of cardiometabolic diseases. *Curr Opin Pharmacol* 2021;**60**:249–254. <https://doi.org/10.1016/j.coph.2021.07.017>
242. Oppenheimer SJ. Iron and its relation to immunity and infectious disease. *J Nutr* 2001;**131**:616S–633S. <https://doi.org/10.1093/jn/131.2.616S>
243. von Haehling S, Ebner N, Evertz R, Ponikowski P, Anker SD. Iron deficiency in heart failure: An overview. *JACC Heart Fail* 2019;**7**:36–46. <https://doi.org/10.1016/j.jchf.2018.07.015>
244. Ponikowski P, Mentz RJ, Hernandez AF, Butler J, Khan MS, van Veldhuisen DJ, et al. Efficacy of ferric carboxymaltose in heart failure with iron deficiency: An individual patient data meta-analysis. *Eur Heart J* 2023;**44**:5077–5091. <https://doi.org/10.1093/eurheartj/ehad586>
245. Palm CL, Nijholt KT, Bakker BM, Westenbrink BD. Short-chain fatty acids in the metabolism of heart failure – rethinking the fat stigma. *Front Cardiovasc Med* 2022;**9**:915102. <https://doi.org/10.3389/fcvm.2022.915102>
246. Yurista SR, Chong CR, Badimon JJ, Kelly DP, de Boer RA, Westenbrink BD. Therapeutic potential of ketone bodies for patients with cardiovascular disease: JACC state-of-the-art review. *J Am Coll Cardiol* 2021;**77**:1660–1669. <https://doi.org/10.1016/j.jacc.2020.12.065>
247. Ho KL, Zhang L, Wagg C, Al Batran R, Gopal K, Levasseur J, et al. Increased ketone body oxidation provides additional energy for the failing heart without improving cardiac efficiency. *Cardiovasc Res* 2019;**115**:1606–1616. <https://doi.org/10.1093/cvr/cvz045>

248. Ho KL, Karwi QG, Wagg C, Zhang L, Vo K, Altamimi T, et al. Ketones can become the major fuel source for the heart but do not increase cardiac efficiency. *Cardiovasc Res* 2021;**117**:1178–1187. <https://doi.org/10.1093/cvr/cvaa1435>
249. Yurista SR, Matsuura TR, Sillje HHW, Nijholt KT, McDauid KS, Shewale SV, et al. Ketone ester treatment improves cardiac function and reduces pathologic remodeling in preclinical models of heart failure. *Circ Heart Fail* 2021;**14**:e007684. <https://doi.org/10.1161/CIRCHEARTFAILURE.120.007684>
250. Selvaraj S, Fu Z, Jones P, Kwee LC, Windsor SL, Ilkayeva O, et al.; DEFINE-HF Investigators. Metabolomic profiling of the effects of dapagliflozin in heart failure with reduced ejection fraction: DEFINE-HF. *Circulation* 2022;**146**:808–818. <https://doi.org/10.1161/CIRCULATIONAHA.122.060402>
251. Yurista SR, Sillje HHW, Oberdorf-Maass SU, Schouten EM, Pavez Giani MG, Hillebrands JL, et al. Sodium-glucose co-transporter 2 inhibition with empagliflozin improves cardiac function in non-diabetic rats with left ventricular dysfunction after myocardial infarction. *Eur J Heart Fail* 2019;**21**:862–873. <https://doi.org/10.1002/ehf.1473>
252. Santos-Gallego CG, Requena-Ibanez JA, San Antonio R, Ishikawa K, Watanabe S, Picatoste B, et al. Empagliflozin ameliorates adverse left ventricular remodeling in nondiabetic heart failure by enhancing myocardial energetics. *J Am Coll Cardiol* 2019;**73**:1931–1944. <https://doi.org/10.1016/j.jacc.2019.01.056>
253. Deng Y, Xie M, Li Q, Xu X, Ou W, Zhang Y, et al. Targeting mitochondria-inflammation circuit by beta-hydroxybutyrate mitigates HFpEF. *Circ Res* 2021;**128**:232–245. <https://doi.org/10.1161/CIRCRESAHA.120.317933>
254. Berg-Hansen K, Gopalasingam N, Christensen KH, Ladefoged B, Andersen MJ, Poulsen SH, et al. Cardiovascular effects of oral ketone ester treatment in patients with heart failure with reduced ejection fraction: A randomized, controlled, double-blind trial. *Circulation* 2024;**149**:1474–1489. <https://doi.org/10.1161/CIRCULATIONAHA.123.067971>
255. Challa AA, Lewandowski ED. Short-chain carbon sources: Exploiting pleiotropic effects for heart failure therapy. *JACC Basic Transl Sci* 2022;**7**:730–742. <https://doi.org/10.1016/j.jacbs.2021.12.010>
256. Kim SR, Lee SG, Kim SH, Kim JH, Choi E, Cho W, et al. SGLT2 inhibition modulates NLRP3 inflammasome activity via ketones and insulin in diabetes with cardiovascular disease. *Nat Commun* 2020;**11**:2127. <https://doi.org/10.1038/s41467-020-15983-6>
257. Chriett S, Dabek A, Wojtala M, Vidal H, Balcerczyk A, Pirola L. Prominent action of butyrate over beta-hydroxybutyrate as histone deacetylase inhibitor, transcriptional modulator and anti-inflammatory molecule. *Sci Rep* 2019;**9**:742. <https://doi.org/10.1038/s41598-018-36941-9>
258. Petersen AM, Pedersen BK. The anti-inflammatory effect of exercise. *J Appl Physiol (1985)* 2005;**98**:1154–1162. <https://doi.org/10.1152/jappphysiol.00164.2004>
259. Kraigher-Krainer E, Lyass A, Massaro JM, Lee DS, Ho JE, Levy D, et al. Association of physical activity and heart failure with preserved vs. reduced ejection fraction in the elderly: The Framingham Heart Study. *Eur J Heart Fail* 2013;**15**:742–746. <https://doi.org/10.1093/eurjhf/hft025>
260. Padilha CS, Von Ah Morano AE, Kruger K, Rosa-Neto JC, Lira FS. The growing field of immunometabolism and exercise: Key findings in the last 5 years. *J Cell Physiol* 2022;**237**:4001–4020. <https://doi.org/10.1002/jcp.30866>
261. Kim SD, Yeun YR. Effects of resistance training on C-reactive protein and inflammatory cytokines in elderly adults: A systematic review and meta-analysis of randomized controlled trials. *Int J Environ Res Public Health* 2022;**19**:3434. <https://doi.org/10.3390/ijerph19063434>
262. Feng L, Li G, An J, Liu C, Zhu X, Xu Y, et al. Exercise training protects against heart failure via expansion of myeloid-derived suppressor cells through regulating IL-10/STAT3/S100A9 pathway. *Circ Heart Fail* 2022;**15**:e008550. <https://doi.org/10.1161/CIRCHEARTFAILURE.121.008550>
263. Hodgman CF, Hunt RM, Crane JC, Elzayat MT, LaVoy EC. A scoping review on the effects of physical exercise and fitness on peripheral leukocyte energy metabolism in humans. *Exerc Immunol Rev* 2023;**29**:54–87.
264. Dorneles GP, Lira FS, Romao PRT, Kruger K, Rosa-Neto JC, Peres A, et al. Levels of cardiorespiratory fitness in men exerts strong impact on lymphocyte function after mitogen stimulation. *J Appl Physiol (1985)* 2021;**130**:1133–1142. <https://doi.org/10.1152/jappphysiol.01051.2020>
265. Rosa-Neto JC, Lira FS, Little JP, Landells G, Islam H, Chazaud B, et al. Immunometabolism-fit: How exercise and training can modify T cell and macrophage metabolism in health and disease. *Exerc Immunol Rev* 2022;**28**:29–46.
266. Weng TP, Huang SC, Chuang YF, Wang JS. Effects of interval and continuous exercise training on CD4 lymphocyte apoptotic and autophagic responses to hypoxic stress in sedentary men. *PLoS One* 2013;**8**:e80248. <https://doi.org/10.1371/journal.pone.0080248>